

COHOMOLOGICAL REPRESENTATIONS OF PARAHORIC SUBGROUPS

CHARLOTTE CHAN AND ALEXANDER IVANOV

ABSTRACT. We generalize a cohomological construction of representations due to Lusztig from the hyperspecial case to arbitrary parahoric subgroups of a reductive group over a local field which splits over an unramified extension. We compute the character of these representations on certain very regular elements.

1. INTRODUCTION

Let k be a non-archimedean local field with finite residue field. Let G be a reductive group over k , and let $T \subseteq G$ be a maximal torus defined over k and split over an unramified extension of k . Let P be a parahoric model of G , defined over the integers \mathcal{O}_k . We denote the schematic closure of T in P again by T . We will construct and study a tower of varieties over an algebraic closure of the residue field \mathbb{F}_q of k whose cohomology realizes interesting representations of $P(\mathcal{O}_k)$ parametrized by characters of $T(\mathcal{O}_k)$. This construction generalizes classical Deligne–Lusztig theory [DL76] (for reductive groups over finite fields), as well as the work of Lusztig [Lus04] and Stasinski [Sta09] (for reductive groups over henselian rings). Further, we give an explicit formula for the character on certain very regular elements, generalizing a special case of the character formula for representations of reductive groups over finite fields [DL76, Theorem 4.2].

More precisely, choose a Borel subgroup of G containing T (defined over some unramified extension of k) with unipotent radical U . Fix a Moy–Prasad filtration quotient \mathbb{G} of P , regarded as a smooth affine group scheme of finite type over \mathbb{F}_q . As such, one has a Frobenius $\sigma: \mathbb{G} \rightarrow \mathbb{G}$ and the corresponding Lang map $\mathbb{G} \rightarrow \mathbb{G}$, $g \mapsto g^{-1}\sigma(g)$. In \mathbb{G} we have the subgroups \mathbb{T} and \mathbb{U} , corresponding to the closures of T and U in P . Consider the subscheme $S_{T,U} \subset \mathbb{G}$ defined as the preimage of \mathbb{U} under the Lang map. By construction, $S_{T,U}$ has a natural action of $P(\mathcal{O}_k) \times T(\mathcal{O}_k)$ given by left and right multiplication. For a smooth character $\theta: T(\mathcal{O}_k) \rightarrow \overline{\mathbb{Q}}_\ell^\times$, we define $R_{T,U}^\theta$ to be the θ -isotypic component of the alternating sum of the cohomology groups of $S_{T,U}$ with $\overline{\mathbb{Q}}_\ell$ -coefficients. This is a virtual $P(\mathcal{O}_k)$ -representation.

Theorem 1.1 (cf. Corollary 4.7). *If θ is sufficiently generic, then $R_{T,U}^\theta$ is independent of the choice of U . Moreover, if the stabilizer of θ in the Weyl group of the special fiber of P is trivial, then $\pm R_{T,U}^\theta$ is an irreducible representation of $P(\mathcal{O}_k)$.*

The proof of Theorem 1.1 mainly follows the original method of Lusztig [Lus04], which treated the special case where P is reductive over \mathcal{O}_k . Some technical issues arise in the general setting; these are treated in Sections 2 and 3, especially Sections 3.3, 3.4 and 3.5.

Our second result is the computation of traces of *unramified very regular* elements of $P(\mathcal{O}_k)$ acting on $R_{T,U}^\theta$ (Definition 5.1). The proof is based on the Deligne–Lusztig fixed point formula [DL76, Theorem 3.2] and adapts ideas of [DL76, Theorem 4.2].

Theorem 1.2 (cf. Theorem 5.3). *For any character $\theta: T(\mathcal{O}_k) \rightarrow \overline{\mathbb{Q}}_\ell^\times$ and any unramified very regular element $g \in P(\mathcal{O}_k)$,*

$$\mathrm{Tr}(g, R_{T,U}^\theta) = \sum_{w \in W_{\mathbf{x}}(T, Z^\circ(g))^\sigma} (\theta \circ \mathrm{Ad}(w^{-1}))(g),$$

where the sum ranges over the finite set of σ -invariant elements in the principal homogeneous space $W_{\mathbf{x}}(T, Z^\circ(g))$ under the Weyl group of the special fiber of T in the special fiber of P (Section 2.8).

When G is any inner form of GL_n over k and T is an unramified maximal elliptic torus, we prove in [CI18] that the semi-infinite Deligne–Lusztig set of Lusztig [Lus79] is a scheme and its cohomology realizes the compact induction to $G(k)$ of (an extension of) the $P(\mathcal{O}_k)$ -representations $R_{T,U}^\theta$. Already in this setting, it is not enough to study $R_{T,U}^\theta$ for reductive P ; for example, when G is the anisotropic modulo center inner form of GL_n , the relevant parahoric is an Iwahori subgroup. This can occur even if G is split: if $G = \mathrm{Sp}_4$, then there is a conjugacy class of maximal elliptic tori in G , such that the relevant P is non-reductive, with the reductive quotient of the special fiber being isomorphic to $\mathrm{SL}_2 \times \mathrm{SL}_2$.

As such, we expect this work to be closely related to the problem of geometrically constructing representations of p -adic groups in general. More specifically, we expect that if T is elliptic and $\theta: T(k) \rightarrow \overline{\mathbb{Q}}_\ell^\times$ is a sufficiently generic character, then the compact induction to $G(k)$ of (an extension of) the $P(\mathcal{O}_k)$ -representation $R_{T,U}^\theta$ is related to the supercuspidal representations constructed by Yu [Yu01]. Both the irreducibility of and the character formula for $R_{T,U}^\theta$ are crucial ingredients to understanding the corresponding $G(k)$ -representation within the context of the local Langlands correspondence.

Acknowledgements. The first author was partially supported by an NSF Postdoctoral Research Fellowship (DMS-1802905) and by the DFG via the Leibniz Prize of Peter Scholze. The second author was supported by the DFG via the Leibniz Preis of Peter Scholze.

2. PRELIMINARIES

2.1. Notation. We denote by k a non-archimedean local field with residue field \mathbb{F}_q of prime characteristic p , and by \check{k} the completion of a maximal unramified extension of k . We denote by $\mathcal{O}_k, \mathfrak{p}_k$ (resp. $\mathcal{O}, \mathfrak{p}$) the integers and the maximal ideal of k (resp. \check{k}). The residue field of \check{k} is an algebraic closure $\overline{\mathbb{F}}_q$ of \mathbb{F}_q . We write σ for the Frobenius automorphism of \check{k} , which is the unique k -automorphism of \check{k} , lifting the \mathbb{F}_q -automorphism $x \mapsto x^q$ of $\overline{\mathbb{F}}_q$. Finally, we denote by ϖ a uniformizer of k (and hence of \check{k}) and by $\mathrm{ord} = \mathrm{ord}_{\check{k}}$ the valuation of \check{k} , normalized such that $\mathrm{ord}(\varpi) = 1$.

If k has positive characteristic, we let \mathbb{W} denote the ring scheme over \mathbb{F}_q where for any \mathbb{F}_q -algebra A , $\mathbb{W}(A) = A[[\varpi]]$. If k has mixed characteristic, we let \mathbb{W} denote the k -ramified Witt ring scheme over \mathbb{F}_q so that $\mathbb{W}(\mathbb{F}_q) = \mathcal{O}_k$ and $\mathbb{W}(\overline{\mathbb{F}}_q) = \mathcal{O}$. As the Witt vectors are only well behaved on perfect \mathbb{F}_q -algebras, algebro-geometric considerations when k has mixed characteristic are taken up to perfection. We fix the following convention.

Convention. If k has mixed characteristic, whenever we speak of a scheme over its residue field \mathbb{F}_q , we mean a *perfect scheme*, that is a functor a set-valued functor on perfect \mathbb{F}_q -algebras.

For results on perfect schemes we refer to [Zhu17, BS17]. Note that passing to perfection does not affect the ℓ -adic étale cohomology; thus for purposes of this paper, we could in principle

pass to perfection in all cases. However, in the equal characteristic case working on non-perfect rings does not introduce complications, and we prefer to work in this slightly greater generality.

Fix a prime $\ell \neq p$ and an algebraic closure $\overline{\mathbb{Q}}_\ell$ of \mathbb{Q}_ℓ . The field of coefficients of all representations is assumed to be $\overline{\mathbb{Q}}_\ell$ and all cohomology groups throughout are compactly supported ℓ -adic étale cohomology groups.

2.2. Group-theoretic data. We let G be a connected reductive group over k , such that the base change $G_{\check{k}}$ to \check{k} is split. Let T be a k -rational, \check{k} -split maximal torus in G . Let $\mathcal{B}_{\check{k}}$ and \mathcal{B}_k denote the Bruhat–Tits building of the adjoint group of G over \check{k} and over k , and let $\mathcal{A}_{T,\check{k}} \subseteq \mathcal{B}_{\check{k}}$ denote the apartment of T . Note that there is a natural action of $\text{Aut}(\check{k}/k) = \langle \sigma \rangle$ on $\mathcal{B}_{\check{k}}$ and on $\mathcal{A}_{T,\check{k}}$, and that $\mathcal{B}_k = \mathcal{B}_{\check{k}}^{(\sigma)}$.

Let $X^*(T)$ and $X_*(T)$ denote the group of characters and cocharacters of T . We denote by $\langle \cdot, \cdot \rangle: X^*(T) \times X_*(T) \rightarrow \mathbb{Z}$ the natural \mathbb{Z} -linear pairing between them. We extend it to the uniquely determined \mathbb{R} -linear pairing $\langle \cdot, \cdot \rangle: X^*(T)_{\mathbb{R}} \times X_*(T)_{\mathbb{R}} \rightarrow \mathbb{R}$, where we write $M_{\mathbb{R}} = M \otimes_{\mathbb{Z}} \mathbb{R}$ for a \mathbb{Z} -module M .

Denote by Φ the set of roots of T in $G_{\check{k}}$ and for a root $\alpha \in \Phi$ let $U_\alpha \subseteq G_{\check{k}}$ denote the corresponding root subgroup. There is an action of $\langle \sigma \rangle$ on Φ . Fix a Chevalley system $u_\alpha: \mathbb{G}_a \xrightarrow{\sim} U_\alpha$ for $G_{\check{k}}$ (cf. e.g. [BT84, 4.1.3]). To any root $\alpha \in \Phi$ we can attach the valuation $\varphi_\alpha: U_\alpha(\check{k}) \rightarrow \mathbb{Z}$ given by $\varphi_\alpha(u_\alpha(y)) = \text{ord}(y)$. The set of valuations $\{\varphi_\alpha\}_{\alpha \in \Phi}$ defines a point \mathbf{x}_0 in the apartment $\mathcal{A}_{T,\check{k}}$. Moreover $\mathcal{A}_{T,\check{k}}$ is an affine space under $X_*(T)_{\mathbb{R}}$ and the point $\mathbf{x}_0 + v \in \mathcal{A}_{T,\check{k}}$ for $v \in X_*(T)_{\mathbb{R}}$ corresponds to the valuations $\{\tilde{\varphi}_\alpha\}_{\alpha \in \Phi}$ of the root datum given by $\tilde{\varphi}_\alpha(u) = \varphi_\alpha(u) + \langle \alpha, v \rangle$ (see [BT72, 6.2]).

We let U, U^- be the unipotent radicals of two opposite \check{k} -rational Borel subgroups of $G_{\check{k}}$ containing T .

2.3. Affine roots and filtration on the torus. We have the set Φ_{aff} of affine roots of T in $G_{\check{k}}$. It is the set of affine functions of $\mathcal{A}_{T,\check{k}}$ defined as

$$\Phi_{\text{aff}} = \{\mathbf{x} \mapsto \alpha(\mathbf{x} - \mathbf{x}_0) + m : \alpha \in \Phi, m \in \mathbb{Z}\}.$$

Denote the affine root $(\alpha, m): \mathbf{x} \mapsto \alpha(\mathbf{x} - \mathbf{x}_0) + m$ and call α its vector part. We have the affine root subgroups $\check{U}_{\alpha,m} \subseteq U_\alpha(\check{k})$, defined by

$$\check{U}_{\alpha,m} = \{u \in U_\alpha(\check{k}) : u = 1 \text{ or } \varphi_\alpha(u) \geq m\}$$

They define a descending separated filtration of $U_\alpha(\check{k})$. There is a natural action of Frobenius σ on the set of affine roots. We make it explicit:

Lemma 2.1. *Let $(\alpha, m) \in \Phi_{\text{aff}}$. Then $\sigma(\alpha, m) = (\sigma(\alpha), m - \langle \alpha, \sigma(\mathbf{x}_0) - \mathbf{x}_0 \rangle)$.*

Proof. We have $\sigma(\alpha, m) = (\sigma(\alpha), m')$ for some $m' \in \mathbb{Z}$. The evaluation of the affine-linear form (α, m) on the apartment $\mathcal{A}_{T,\check{k}}$ is σ -linear, thus we have for all $\mathbf{x} \in \mathcal{A}_{T,\check{k}}$:

$$\sigma(\alpha, m)(\mathbf{x}) = (\alpha, m)(\sigma^{-1}(\mathbf{x})) = \langle \alpha, \sigma^{-1}(\mathbf{x}) - \mathbf{x}_0 \rangle + m = \langle \sigma(\alpha), \mathbf{x} - \mathbf{x}_0 \rangle + m - \langle \sigma(\alpha), \sigma(\mathbf{x}_0) - \mathbf{x}_0 \rangle.$$

On the other side, $(\sigma(\alpha), m')(\mathbf{x}) = \langle \sigma(\alpha), \mathbf{x} - \mathbf{x}_0 \rangle + m'$, whence the lemma. \square

Let $\tilde{\mathbb{R}} = \mathbb{R} \cup \{r+ : r \in \mathbb{R}\} \cup \{\infty\}$ denote the ordered monoid as in [BT72, 6.4.1]. Let $\check{T}^0 \subseteq T(\check{k})$ be the maximal bounded subgroup. For $r \in \tilde{\mathbb{R}}_{\geq 0} \setminus \{\infty\}$, we have a descending

separated filtration of \check{T}^0 given by

$$\check{T}^r = \{t \in \check{T}^0 : \text{ord}(\chi(t) - 1) \geq r \ \forall \chi \in X^*(T)\}.$$

2.4. Parahoric subgroups, Moy–Prasad filtration and integral models. Fix a point $\mathbf{x} \in \mathcal{A}_{T,\check{k}}$. Following Bruhat and Tits [BT84, 5.2.6], there is a parahoric group scheme $P_{\mathbf{x}}$ over \mathcal{O} attached to \mathbf{x} , with generic fiber G , and with connected special fiber. The group $\check{P}_{\mathbf{x}} := P_{\mathbf{x}}(\mathcal{O})$ is generated by \check{T}^0 and $\check{U}_{\alpha,m}$ for all $(\alpha, m) \in \Phi_{\text{aff}}$ such that $\langle \alpha, \mathbf{x} - \mathbf{x}_0 \rangle \geq -m$ (that is, $(\alpha, m)(\mathbf{x}) \geq 0$). The schematic closure of T in $P_{\mathbf{x}}$ is the connected Néron model of T . We denote it again by T . We have $T(\mathcal{O}) = \check{T}^0$. (As $G_{\check{k}}$ is split, condition (T) of [Yu02, 8.1] is satisfied. The claim about the closure of T in $P_{\mathbf{x}}$ follows e.g. from [Yu02, Corollary 8.6(ii)]. Again, because $G_{\check{k}}$ is split, it also follows ([BT84, 4.6.1]) that the connected Néron model of T is equal to the maximal subgroup scheme of finite type of the lft model of T . The \mathcal{O} -points of the latter are equal to \check{T}^0 , hence we indeed have $T(\mathcal{O}) = \check{T}^0$.)

The Moy–Prasad filtration on $\check{P}_{\mathbf{x}}$ is given by the series of normal subgroups $\check{P}_{\mathbf{x}}^r \subseteq \check{P}_{\mathbf{x}}$ ($r \in \mathbb{R}_{\geq 0} \setminus \{\infty\}$), generated by \check{T}^r and $\check{U}_{(\alpha,m)}$ for all $(\alpha, m) \in \Phi_{\text{aff}}$ such that $\langle \alpha, \mathbf{x} - \mathbf{x}_0 \rangle \geq r - m$. By [Yu02, 8.6 Corollary], there is a unique smooth \mathcal{O} -model $P_{\mathbf{x}}^r$ of G , such that $P_{\mathbf{x}}^r(\mathcal{O}) = \check{P}_{\mathbf{x}}^r$. Moreover, part (ii) of the same corollary describes the schematic closures of U_{α} , T in $P_{\mathbf{x}}^r$, and in particular, we have

$$\check{P}_{\mathbf{x}}^r \cap U_{\alpha}(\check{k}) = \check{U}_{\alpha, [r - \langle \alpha, \mathbf{x} - \mathbf{x}_0 \rangle]} \quad \text{and} \quad \check{P}_{\mathbf{x}}^r \cap T(\check{k}) = \check{T}^r. \quad (2.1)$$

Note that for $r \in \mathbb{R}_{\geq 0}$, we have $\check{P}_{\mathbf{x}}^{r+} = \bigcup_{s \in \mathbb{R}, s > r} \check{P}_{\mathbf{x}}^s$. For further properties of the Moy–Prasad filtration we refer to [MP94, §2.6], and for further properties of the smooth models $P_{\mathbf{x}}^r$ we refer to [Yu02].

Assume now that $\mathbf{x} \in \mathcal{A}_{T,\check{k}} \cap \mathcal{B}_k$. Then all group schemes $P_{\mathbf{x}}$, $P_{\mathbf{x}}^r$ descend to smooth group schemes over \mathcal{O}_k , again denoted by $P_{\mathbf{x}}$, $P_{\mathbf{x}}^r$ (cf. [Yu02, §9.1]). In particular, all groups $\check{P}_{\mathbf{x}}^r$ ($r \geq 0$) are σ -stable (this can also be deduced from Lemma 2.1, which shows that σ maps $\check{U}_{\alpha, [r - \langle \alpha, \mathbf{x} - \mathbf{x}_0 \rangle]}$ isomorphically onto $\check{U}_{\sigma(\alpha), [r - \langle \sigma(\alpha), \mathbf{x} - \mathbf{x}_0 \rangle]}$), and $P_{\mathbf{x}}(\mathcal{O}_k) = \check{P}_{\mathbf{x}}^{\sigma}$ and $P_{\mathbf{x}}^r(\mathcal{O}_k) = (\check{P}_{\mathbf{x}}^r)^{\sigma}$.

2.5. Moy–Prasad quotients. For a scheme \mathfrak{X} over \mathcal{O}_k (resp. over \mathcal{O}), the functor of positive loops $L^+\mathfrak{X}$ is the functor on \mathbb{F}_q -algebras (resp. $\overline{\mathbb{F}}_q$ -algebras) given by

$$L^+\mathfrak{X}(R) = \mathfrak{X}(\mathbb{W}(R))$$

If \mathfrak{X} is affine and of finite type, then $L^+\mathfrak{X}$ is represented by an affine scheme.

Let $\mathbf{x} \in \mathcal{A}_{T,\check{k}} \cap \mathcal{B}_k$ be as in Section 2.4. We have the infinite-dimensional affine \mathbb{F}_q -group scheme $L^+P_{\mathbf{x}}$, and will now introduce convenient finite-dimensional quotients of it. Let $r \in \mathbb{Z}_{\geq 1}$. We consider the fpqc quotient sheaf $\mathbb{G}_r := L^+P_{\mathbf{x}}/L^+P_{\mathbf{x}}^{(r-1)+}$. By [CI18, Proposition 4.2(ii)] it is representable by a smooth affine group scheme over \mathbb{F}_q of finite type, which we again denote by \mathbb{G}_r . From [Yu02, Theorem 8.8], along with the fact that $L^+P_{\mathbf{x}}^{(r-1)+}$ is pro-unipotent, it follows by taking Galois cohomology,

$$\check{\mathbb{G}}_r := \mathbb{G}_r(\overline{\mathbb{F}}_q) = \check{P}_{\mathbf{x}}/\check{P}_{\mathbf{x}}^{(r-1)+} \quad \text{and} \quad \check{\mathbb{G}}_r^{\sigma} = \mathbb{G}_r(\mathbb{F}_q) = (\check{P}_{\mathbf{x}}/\check{P}_{\mathbf{x}}^{(r-1)+})^{\sigma}.$$

For $r \geq s \geq 1$ we have natural surjections of \mathbb{F}_q -groups $L^+P_{\mathbf{x}} \rightarrow \mathbb{G}_r \rightarrow \mathbb{G}_s$. We write $\mathbb{G}_r^s = \ker(\mathbb{G}_r \rightarrow \mathbb{G}_s)$ and $\check{\mathbb{G}}_r^s := \mathbb{G}_r^s(\overline{\mathbb{F}}_q)$. Moreover, we also have natural surjections $\mathbb{G}_2 \rightarrow P_{\mathbf{x}} \otimes_{\mathcal{O}_k} \mathbb{F}_q \rightarrow (P_{\mathbf{x}} \otimes_{\mathcal{O}_k} \mathbb{F}_q)^{\text{red}} = \mathbb{G}_1$ identifying \mathbb{G}_1 with the reduced quotient of the special fiber of $P_{\mathbf{x}}$ and of each \mathbb{G}_r .

2.6. Subgroups of \mathbb{G}_r . Let $H \subseteq G_{\check{k}}$ be a closed subgroup scheme. Let $r \in \mathbb{Z}_{\geq 1}$. We will attach to H the subgroup $\mathbb{H}_r \subseteq \mathbb{G}_{r, \overline{\mathbb{F}}_q}$ as follows. The schematic closure $H_{\mathbf{x}}$ of H in $P_{\mathbf{x}, \mathcal{O}}$ is flat (by [BT72, 1.2.6] as \mathcal{O} -flat is equivalent to \mathcal{O} -torsion free). It follows that $H_{\mathbf{x}}$ is a closed subgroup scheme of $P_{\mathbf{x}, \mathcal{O}}$ ([BT72, 1.2.7]). Apply L_r^+ to the inclusion $H_{\mathbf{x}} \subseteq P_{\mathbf{x}, \mathcal{O}}$ to obtain the subgroup scheme $L_r^+ H_{\mathbf{x}} \subseteq L_r^+ P_{\mathbf{x}, \mathcal{O}}$. The last inclusion is a closed immersion (e.g. by [Gre61, Corollary 2 on p. 639]). We define the closed subgroup scheme $\mathbb{H}_r \subseteq \mathbb{G}_{r, \overline{\mathbb{F}}_q}$ as the image of $L_r^+ H_{\mathbf{x}}$ under $L_r^+ P_{\mathbf{x}, \mathcal{O}} \rightarrow \mathbb{G}_{r, \overline{\mathbb{F}}_q}$. We write $\mathbb{H}_r^s := \ker(\mathbb{H}_r \rightarrow \mathbb{H}_s)$ and $\mathbb{H}_r^{s,*} := \mathbb{H}_r^s \setminus \mathbb{H}_r^{s+1}$.

Suppose now additionally that $H_{\mathbf{x}}$ is smooth. Then $L_r^+ H_{\mathbf{x}}$ is reduced (one could e.g. use [Gre63, Corollary 2 on p. 264]), and hence \mathbb{H}_r is too. If H is already defined over the finite subextension of \check{k}/k of degree d , then $H_{\mathbf{x}}$ is defined over the integers of this subextension. This implies that $\mathbb{H}_r(\overline{\mathbb{F}}_q)$ is stable under the action of σ^d . Hence \mathbb{H}_r is defined over \mathbb{F}_{q^d} (here we use that \mathbb{H}_r is reduced separated scheme of finite type over \mathbb{F}_q).

Using the procedure described above we obtain the closed \mathbb{F}_q -subgroup $\mathbb{T}_r \subseteq \mathbb{G}_r$ attached to $T \subseteq G$. Analogously, we have the subgroups $\mathbb{U}_r, \mathbb{U}_r^- \subseteq \mathbb{G}_{r, \overline{\mathbb{F}}_q}$ corresponding to $U, U^- \subseteq G_{\check{k}}$ and for any root $\alpha \in \Phi$ the subgroup $\mathbb{U}_{r, \alpha} \subseteq \mathbb{G}_{r, \overline{\mathbb{F}}_q}$ corresponding to U_{α} . Note that all these are reduced connected closed subgroups of $\mathbb{G}_{r, \overline{\mathbb{F}}_q}$. Moreover, $\mathbb{U}_{r, \alpha}$ is defined over \mathbb{F}_{q^d} where $d \in \mathbb{Z}_{\geq 1}$ is the smallest positive integer such that $\sigma^d(\alpha) = \alpha$ in Φ (indeed the group $U_{\alpha, \mathbf{x}}$ is smooth by [Yu02, 8.3 Theorem (ii)]), and a similar statement holds for $\mathbb{U}_r, \mathbb{U}_r^-$.

For any reduced $\overline{\mathbb{F}}_q$ -subscheme $\mathbb{X}_r \subseteq \mathbb{G}_{r, \overline{\mathbb{F}}_q}$, we define $\check{X}_r := \mathbb{X}_r(\overline{\mathbb{F}}_q) \subseteq \mathbb{G}_r(\overline{\mathbb{F}}_q) = \check{G}_r$. Thus for example we write $\check{U}_{\alpha, r}^a = \mathbb{U}_{\alpha, r}^a(\overline{\mathbb{F}}_q)$ for $\alpha \in \Phi$ and $1 \leq a \leq r-1$. Following Lusztig, we denote by \mathcal{T} the groups \mathbb{T}_r^{r-1} . For $\alpha \in \Phi$, let $T^{\alpha} \subset T_{\check{k}} \subset G_{\check{k}}$ be the unique 1-dimensional torus contained in the subgroup of $G_{\check{k}}$ generated by U_{α} and $U_{-\alpha}$; let \mathbb{T}_r^{α} be the corresponding subgroup scheme of $\mathbb{G}_{r, \overline{\mathbb{F}}_q}$ and write $\mathcal{T}^{\alpha} := \mathbb{T}_r^{\alpha, r-1}$.

Lemma 2.2. *Let $r \in \mathbb{Z}_{\geq 1}$ and $1 \leq a \leq r-1$.*

- (i) *The group \mathbb{G}_r is generated by \mathbb{T}_r and all $\mathbb{U}_{\alpha, r}$ ($\alpha \in \Phi$).*
- (ii) *The group \mathbb{G}_r^{a+1} is generated by \mathbb{T}_r^{a+1} and all $\mathbb{U}_{\beta, r}^{a+1}$ ($\alpha \in \Phi$).*

Proof. As $\mathbb{G}_r, \mathbb{G}_r^a$ are smooth affine \mathbb{F}_q -groups, the assertions can be checked on $\overline{\mathbb{F}}_q$ -points. Now both cases follow from [Yu02, Theorem 8.3] applied to the smooth models $P_{\mathbf{x}}$ and $P_{\mathbf{x}}^{a+1}$ of G respectively (note that with notations as in *loc. cit.*, the group $G(k)_{\mathbf{x}, f}$ is by definition the one generated by all $U_a(k)_{\mathbf{x}, f(a)}$). \square

Remark 2.3. Let U' be the unipotent radical of some other Borel subgroup of $G_{\check{k}}$ containing T . Although U and U' are conjugate by an element of $G(\check{k})$, the groups $\mathbb{U}_r(\overline{\mathbb{F}}_q)$ and $\mathbb{U}'_r(\overline{\mathbb{F}}_q)$ need not be isomorphic. For example, let G be the anisotropic modulo center inner form of GL_3 (it splits over \check{k} and its k -points are isomorphic to the units of a division algebra over k). Let \mathbf{x} be the unique point in \mathcal{B}_k . Then $\mathbb{G}_1 = \mathbb{T}_1$ is a torus and (after an appropriate choice of \mathbf{x}_0) one has $\mathbb{G}_2(\overline{\mathbb{F}}_q) = \begin{pmatrix} \mathbb{W}_2(\overline{\mathbb{F}}_q)^{\times} & \overline{\mathbb{F}}_q & \overline{\mathbb{F}}_q \\ \varpi \overline{\mathbb{F}}_q & \mathbb{W}_2(\overline{\mathbb{F}}_q)^{\times} & \overline{\mathbb{F}}_q \\ \varpi \overline{\mathbb{F}}_q & \varpi \overline{\mathbb{F}}_q & \mathbb{W}_2(\overline{\mathbb{F}}_q)^{\times} \end{pmatrix}$ with obvious multiplication. Now, let U and U' be the group of upper- and lower-triangular unipotent matrices in G . Then $\mathbb{U}_2 = \mathbb{U}_2^1$ is non-abelian, whereas $\mathbb{U}'_2 = \mathbb{U}'_2^1$ is abelian. \diamond

2.7. The groups $\mathbb{U}_{\alpha, r}$. We now give explicit formulas for $\mathbb{U}_{\alpha, r} \subseteq \mathbb{G}_r$.

Definition 2.4. Let $\mathbf{x} \in \mathcal{A}_{T, \check{k}}$. We call a root $\alpha \in \Phi$

reductive	if $\langle \alpha, \mathbf{x} - \mathbf{x}_0 \rangle \in \mathbb{Z}$
non-reductive	otherwise.

For any $\alpha \in \Phi$, we may uniquely write $\langle \alpha, \mathbf{x} - \mathbf{x}_0 \rangle = -m_\alpha + \varepsilon_\alpha$ with $m_\alpha \in \mathbb{Z}$ and $0 \leq \varepsilon_\alpha < 1$. We have $m_\alpha = -\lfloor \langle \alpha, \mathbf{x} - \mathbf{x}_0 \rangle \rfloor$.

Note that $\alpha \in \Phi$ is reductive if and only if $\mathbb{U}_{\alpha, 1} \neq 1$.

Lemma 2.5. Let $\mathbf{x} \in \mathcal{A}_{T, \check{k}}$ and let $r \in \mathbb{Z}_{\geq 1}$. Let $\alpha \in \Phi$. We have

$$m_\alpha + m_{-\alpha} = \begin{cases} 0 & \text{if } \alpha \text{ is reductive} \\ 1 & \text{otherwise.} \end{cases}$$

Moreover, the natural map $\check{P}_{\mathbf{x}} \rightarrow \mathbb{G}_r(\overline{\mathbb{F}}_q)$ induces

$$\mathbb{U}_{\alpha, r}(\overline{\mathbb{F}}_q) = \begin{cases} \check{U}_{\alpha, m_\alpha} / \check{U}_{\alpha, m_\alpha + r} & \text{if } \alpha \text{ reductive,} \\ \check{U}_{\alpha, m_\alpha} / \check{U}_{\alpha, m_\alpha + r - 1} & \text{otherwise.} \end{cases}$$

Thus for $a \in \mathbb{Z}$, $r \geq a \geq 1$, the same map induces

$$\mathbb{U}_{\alpha, r}^a(\overline{\mathbb{F}}_q) = \begin{cases} \check{U}_{\alpha, m_\alpha + a} / \check{U}_{\alpha, m_\alpha + r} & \text{if } \alpha \text{ reductive,} \\ \check{U}_{\alpha, m_\alpha + a - 1} / \check{U}_{\alpha, m_\alpha + r - 1} & \text{otherwise.} \end{cases}$$

Finally, we have $\mathbb{T}_r(\overline{\mathbb{F}}_q) = \check{T}^0 / \check{T}^r$ and $\mathbb{T}_r^a = \check{T}^a / \check{T}^r$.

Proof. Noting that $\lceil -s \rceil = -\lfloor s \rfloor$ for $s \in \mathbb{R}$, the lemma follows immediately from (2.1) and the definitions of $\mathbb{U}_{\alpha, r}$, $\mathbb{U}_{\alpha, r}^a$ and \mathbb{G}_r . \square

We have the following elementary lemma will be useful later.

Lemma 2.6. Let $\alpha, \beta \in \Phi$ and assume that $p, q \in \mathbb{Z}_{\geq 1}$, such that $p\alpha + q\beta \in \Phi$. Then $pm_\alpha + qm_\beta - m_{p\alpha + q\beta} = p\varepsilon_\alpha + q\varepsilon_\beta - \varepsilon_{p\alpha + q\beta} = \lfloor p\varepsilon_\alpha + q\varepsilon_\beta \rfloor$. In particular, $pm_\alpha + qm_\beta - m_{p\alpha + q\beta} \geq 0$.

Proof. The first equality is immediate. In particular, $p\varepsilon_\alpha + q\varepsilon_\beta - \varepsilon_{p\alpha + q\beta}$ is an integer. This, along with the fact that $0 \leq \varepsilon_{p\alpha + q\beta} < 1$ by definition, implies the second equality. \square

2.8. Weyl groups. Bruhat–decomposition. We have the group

$$W_{\mathbf{x}}(T) := (N_G(T)(\check{k}) \cap \check{P}_{\mathbf{x}}^0) / \check{T}^0$$

(cf. [HR08, Proposition 8]), and it coincides with the Weyl group $W(\mathbb{T}_1, \mathbb{G}_1)$ of the torus \mathbb{T}_1 in the special fiber \mathbb{G}_1 of $P_{\mathbf{x}}$ ([HR08, Proposition 12]). It follows that both natural maps in the composition

$$W_{\mathbf{x}}(T) \rightarrow N_{\mathbb{G}_r}(\mathbb{T}_r)(\overline{\mathbb{F}}_q) / \mathbb{T}_r(\overline{\mathbb{F}}_q) \rightarrow N_{\mathbb{G}_1}(\mathbb{T}_1)(\overline{\mathbb{F}}_q) / \mathbb{T}_1(\overline{\mathbb{F}}_q)$$

are isomorphisms. Here $N_G(H)$ denotes the scheme-theoretic normalizer of the subgroup H of a group G (note that it might be non-reduced, but we have $N_G(H)(\overline{\mathbb{F}}_q) = N_G(H)_{\text{red}}(\overline{\mathbb{F}}_q) = \{g \in G(\overline{\mathbb{F}}_q) : gHg^{-1} = H\}$). We also note that $W_{\mathbf{x}}(T)$ coincides with the subgroup of the Weyl group $W = W(T, G)$ of T in G generated by the vector parts of all affine roots $\psi \in \Phi_{\text{aff}}$ satisfying $\psi(\mathbf{x}) = 0$ (cf. [Tit79, 1.9, 3.5.1]). It depends only on the facet of $\mathcal{B}_{\check{k}}$ in which \mathbf{x} lies, not on \mathbf{x} itself.

We will need a second k -rational, \check{k} -split maximal torus T' of G whose apartment $\mathcal{A}_{T',\check{k}}$ in $\mathcal{B}_{\check{k}}$ passes through the point \mathbf{x} . Let $N_G(T, T') = \{g \in G : gTg^{-1} = T'\}$ be the transporter from T to T' and analogously, let $N_{\mathbb{G}_r}(\mathbb{T}_r, \mathbb{T}'_r)$ be the transporter from \mathbb{T}_r to \mathbb{T}'_r . (Again, these need not be reduced, but we are interested in $\overline{\mathbb{F}}_q$ -points only.) We then have the principal homogeneous space

$$W_{\mathbf{x}}(T, T') := \check{T}^0 \backslash (N_G(T, T')(\check{k}) \cap \check{P}_{\mathbf{x}}^0) = \mathbb{T}_r(\overline{\mathbb{F}}_q) \backslash N_{\mathbb{G}_r}(\mathbb{T}_r, \mathbb{T}'_r)(\overline{\mathbb{F}}_q).$$

under $W_{\mathbf{x}}(T)$. Indeed, this follows as T and T' are conjugate by an element of $P_{\mathbf{x}}(\mathcal{O})$.

Let $r \geq 1$. For each $w \in W_{\mathbf{x}}(T, T')$ choose a representative $\dot{w} \in N_{\mathbb{G}_r}(\mathbb{T}_r, \mathbb{T}'_r)(\overline{\mathbb{F}}_q)$, and denote its image in \mathbb{G}_1 again by \dot{w} . We have the Bruhat decomposition $\mathbb{G}_1 = \bigsqcup_{w \in W_{\mathbf{x}}(T, T')} \mathbb{G}_{1,w}$ of the reductive quotient, where $\mathbb{G}_{1,w} = \mathbb{U}_1 \dot{w} \mathbb{T}'_1 \mathbb{U}'_1$. For $r \geq 1$, define $\mathbb{G}_{r,w}$ to be the pullback of $\mathbb{G}_{1,w}$ along the natural projection $\mathbb{G}_r \rightarrow \mathbb{G}_1$. Thus $\mathbb{G}_r = \bigsqcup_{w \in W_{\mathbf{x}}(T, T')} \mathbb{G}_{r,w}$. Let $\mathbb{K}_r := \mathbb{U}_r^- \cap \dot{w} \mathbb{U}'_r \dot{w}^{-1}$ and $\mathbb{K}_r^1 := \mathbb{K}_r \cap \mathbb{G}_r^1$.

Lemma 2.7. *For $r \geq 1$, we have $\mathbb{G}_{r,w} = \mathbb{U}_r \mathbb{K}_r^1 \dot{w} \mathbb{T}'_r \mathbb{U}'_r$.*

Proof. We compute

$$\begin{aligned} \mathbb{G}_{r,w} &= \mathbb{U}_r \dot{w} \mathbb{T}'_r \mathbb{G}_r^1 \mathbb{U}'_r = \mathbb{U}_r \dot{w} \mathbb{T}'_r ((\mathbb{G}_r^1 \cap \mathbb{T}'_r)(\mathbb{G}_r^1 \cap \mathbb{U}'_r)(\mathbb{G}_r^1 \cap \mathbb{U}'_r)) \mathbb{U}'_r \\ &= \mathbb{U}_r \dot{w} \mathbb{T}'_r (\mathbb{G}_r^1 \cap \mathbb{U}'_r) \mathbb{U}'_r = \mathbb{U}_r (\dot{w} (\mathbb{G}_r^1 \cap \mathbb{U}'_r) \dot{w}^{-1}) \dot{w} \mathbb{T}'_r \mathbb{U}'_r \\ &= \mathbb{U}_r (\mathbb{U}_r^- \cap \dot{w} (\mathbb{G}_r^1 \cap \mathbb{U}'_r) \dot{w}^{-1}) \dot{w} \mathbb{T}'_r \mathbb{U}'_r = \mathbb{U}_r \mathbb{K}_r^1 \dot{w} \mathbb{T}'_r \mathbb{U}'_r, \end{aligned}$$

where the second equality follows from [BT72, 6.4.48]. \square

2.9. Commutation relations. For two subgroups H_1, H_2 of an abstract group H , we denote by $[H_1, H_2]$ their commutator. For $x, y \in H$, we write $[x, y] := x^{-1}y^{-1}xy$.

For $\alpha \in \Phi$, let $T^\alpha \subseteq T$ denote the image of the coroot corresponding to α . It is a one-dimensional subtorus. We also write $\check{T}^{\alpha,r} = T^\alpha(\check{k}) \cap \check{T}^r$.

Lemma 2.8. (i) *Let $\alpha \in \Phi$ and $r, m \in \widetilde{\mathbb{R}}$. Then $[\check{T}^r, \check{U}_{\alpha,m}] \subseteq \check{U}_{\alpha,m+r}$.*

(ii) *If $\alpha, \beta \in \Phi$, $\alpha \neq -\beta$, and $m_1, m_2 \in \mathbb{Z}$, then $[\check{U}_{\alpha,m_1}, \check{U}_{\beta,m_2}]$ is contained in the group generated by $\check{U}_{p\alpha+q\beta, pm_1+qm_2}$ for all $p, q \in \mathbb{Z}_{\geq 1}$, such that $p\alpha + q\beta \in \Phi$.*

(iii) *Let $\alpha \in \Phi$ and $m_1, m_2 \in \mathbb{Z}$. Then $[\check{U}_{\alpha,m_1}, \check{U}_{-\alpha,m_2}] \subseteq \check{T}^{\alpha, m_1+m_2}$. For any element $x \in \check{U}_{-\alpha, m_2} \setminus \check{U}_{-\alpha, m_2+1}$, the map $\xi \mapsto [\xi, x]$ induces an isomorphism (of abelian groups)*

$$\lambda_x: \check{U}_{\alpha, m_1} / \check{U}_{\alpha, m_1+1} \xrightarrow{\sim} \check{T}^{\alpha, m_1+m_2} / \check{T}^{\alpha, m_1+m_2+1}.$$

Proof. (ii) follows from [BT72, (6.2.1)]. (i), (iii): By considering a morphism from SL_2 to $G_{\check{k}}$, whose image is generated by $U_{\pm\alpha}$ (as in [BT72, (6.2.3) b]), and pulling back the valuation of the root datum along this morphism, it suffices to prove the same statement for $\mathrm{SL}_2(\check{k})$. This is an immediate computation. \square

For two smooth (connected) closed subgroups $\mathbb{H}_1, \mathbb{H}_2$ of a connected linear algebraic group \mathbb{G} over a field, we denote by $[\mathbb{H}_1, \mathbb{H}_2]$ their commutator “in the sense of group varieties” as in [Bor91, §2.3] (it would be more precise to consider the scheme-theoretic commutator, but for our purposes this suffices).

Lemma 2.9. *Let $r \geq 2$ and $1 \leq a \leq r-1$. Let $\alpha \in \Phi$.*

(a) *If α is non-reductive, then $[\mathbb{G}_r^{a+1}, \mathbb{U}_{\alpha,r}^{-a}] = 1$.*

(b) *If α is reductive, then $[\mathbb{G}_r^a, \mathbb{U}_{r,\alpha}^{-a}] = 1$.*

Proof. It suffices to prove the claims on $\overline{\mathbb{F}}_q$ -points. (a): By Lemma 2.2 it suffices to show to show that $[\check{T}_r^{a+1}, \check{U}_{\alpha,r}^{r-a}] = 1$ and that $[\check{U}_{\beta,r}^{a+1}, \check{U}_{\alpha,r}^{r-a}] = 1$ ($\forall \beta \in \Phi$) in \mathbb{G}_r . By Lemma 2.5 \check{T}_r^{a+1} is the image in \check{G}_r of \check{T}^{a+1} , $\check{U}_{\alpha,r}^{r-a}$ is the image of $\check{U}_{\alpha,m_\alpha+r-a-1}$, and similar claims hold for all $\beta \in \Phi$. But $[\check{T}^{a+1}, \check{U}_{\alpha,m_\alpha+r-a-1}] \subseteq \check{U}_{\alpha,r+m_\alpha}$ by Lemma 2.8(i), and $\check{U}_{\alpha,r+m_\alpha}$ maps to 1 in \mathbb{G}_r , so $[\check{T}_r^{a+1}, \check{U}_{\alpha,r}^{r-a}] = 1$ follows. Now assume that $\beta = -\alpha$. Then $-\alpha$ is non-reductive as α is, and by Lemma 2.8(iii), $[\check{U}_{-\alpha,m_{-\alpha}+a}, \check{U}_{\alpha,m_\alpha+r-a-1}] \subseteq \check{T}^{\alpha,r+m_\alpha+m_{-\alpha}-1} = \check{T}^{\alpha,r}$ maps to 1 in \mathbb{G}_r . This shows $[\check{U}_{-\alpha,r}^{a+1}, \check{U}_{\alpha,r}^{r-a}] = 1$. Thus we can assume $\beta \in \Phi$, $\beta \neq -\alpha$. We have two cases.

Case: β is reductive. Then by Lemma 2.5, $\check{U}_{\beta,r}^{a+1}$ is the image in \check{G}_r of $\check{U}_{\beta,m_\beta+a+1}$ and by Lemma 2.8(ii) we have

$$[\check{U}_{\beta,m_\beta+a+1}, \check{U}_{\alpha,m_\alpha+r-a-1}] \subseteq \prod_{\substack{p,q \in \mathbb{Z}_{\geq 1} \\ p\alpha+q\beta \in \Phi}} \check{U}_{p\alpha+q\beta, p(m_\alpha+r-a-1)+q(m_\beta+a+1)}.$$

To ensure that this product maps to 1 in \check{G}_r , it suffices to show that for all $p, q \in \mathbb{Z}_{\geq 1}$ with $p\alpha + q\beta \in \Phi$, one has $p(m_\alpha + r - a - 1) + q(m_\beta + a + 1) \geq m_{p\alpha+q\beta} + r$, or equivalently,

$$pm_\alpha + qm_\beta - m_{p\alpha+q\beta} + (p-1)(r-a-1) + (q-1)(a+1) \geq 0.$$

But this follows from Lemma 2.6.

Case: β is non-reductive. By Lemma 2.5, $\check{U}_{\beta,r}^{a+1}$ is the image in \check{G}_r of $\check{U}_{\beta,m_\beta+a}$ and by Lemma 2.8(ii) we have

$$[\check{U}_{\beta,m_\beta+a}, \check{U}_{\alpha,m_\alpha+r-a-1}] \subseteq \prod_{\substack{p,q \in \mathbb{Z}_{\geq 1} \\ p\alpha+q\beta \in \Phi}} \check{U}_{p\alpha+q\beta, p(m_\alpha+r-a-1)+q(m_\beta+a)}.$$

To show that the image of this product vanishes in \mathbb{G}_r , we have to show that each single term does. Assume that $p\alpha + q\beta$ occurs in the product and is non-reductive. Then vanishing of $\check{U}_{p\alpha+q\beta, p(m_\alpha+r-a-1)+q(m_\beta+a)}$ in \check{G}_r amounts to the inequality

$$pm_\alpha + qm_\beta - m_{p\alpha+q\beta} + (p-1)(r-a-1) + (q-1)a \geq 0,$$

which holds true by Lemma 2.6. Assume finally that $p\alpha + q\beta$ occurs in the product and is reductive. Then vanishing of $\check{U}_{p\alpha+q\beta, p(m_\alpha+r-a-1)+q(m_\beta+a)}$ in \check{G}_r amounts to the inequality

$$pm_\alpha + qm_\beta - m_{p\alpha+q\beta} + (p-1)(r-a-1) + (q-1)a \geq 1,$$

or equivalently,

$$[p\varepsilon_\alpha + q\varepsilon_\beta] + (p-1)(r-a-1) + (q-1)a \geq 1,$$

i.e. it suffices to show that $p\varepsilon_\alpha + q\varepsilon_\beta \geq 1$. But as $p\alpha + q\beta$ is reductive,

$$\mathbb{Z} \ni \langle p\alpha + q\beta, \mathbf{x} - \mathbf{x}_0 \rangle = p\langle \alpha, \mathbf{x} - \mathbf{x}_0 \rangle + q\langle \beta, \mathbf{x} - \mathbf{x}_0 \rangle = -pm_\alpha - qm_\beta + p\varepsilon_\alpha + q\varepsilon_\beta. \quad (2.2)$$

As $-pm_\alpha - qm_\beta \in \mathbb{Z}$, we deduce $p\varepsilon_\alpha + q\varepsilon_\beta \in \mathbb{Z}$. On the other side $\varepsilon_\alpha, \varepsilon_\beta > 0$ (as α, β non-reductive), and hence $p\varepsilon_\alpha + q\varepsilon_\beta > 0$. Thus, $p\varepsilon_\alpha + q\varepsilon_\beta \geq 1$. This finishes the proof of (a).

(b): We have $[\check{T}_r^a, \check{U}_{\alpha,m_\alpha+r-a}] \subseteq \check{U}_{\alpha,m_\alpha+r}$ by Lemma 2.8(i), and the latter group maps to 1 in \check{G}_r . Thus $[\check{T}_r^a, \check{U}_{\alpha,r}^{r-a}] = 1$. Further, Lemma 2.8(iii) shows

$$[\check{U}_{-\alpha,m_{-\alpha}+a}, \check{U}_{\alpha,m_\alpha+r-a}] \subseteq \check{T}^{\alpha,m_\alpha+m_{-\alpha}+r} = \check{T}^{\alpha,r},$$

which maps to 1 in \check{G}_r . Thus $[\check{U}_{-\alpha,r}^a, \check{U}_{\alpha,r}^{r-a}] = 1$. Finally, let $\beta \in \Phi$, $\beta \neq -\alpha$. Again we have two cases.

Case: β is reductive. By Lemma 2.8(ii),

$$[\check{U}_{\beta,m_\beta+a}, \check{U}_{\alpha,m_\alpha+r-a}] \subseteq \prod_{\substack{p,q \in \mathbb{Z}_{\geq 1} \\ p\alpha+q\beta \in \Phi}} \check{U}_{p\alpha+q\beta, p(m_\alpha+r-a)+q(m_\beta+a)},$$

Now, by Lemma 2.6 we have

$$p(m_\alpha + r - a) + q(m_\beta + a) \geq m_{p\alpha+q\beta} + r.$$

So, regardless of whether $p\alpha + q\beta$ is reductive or not, it follows that $\check{U}_{p\alpha+q\beta, p(m_\alpha+r-a)+q(m_\beta+a)}$ maps to 1 in \check{G}_r , and hence $[\check{U}_{\beta,r}^a, \check{U}_{\alpha,r}^{r-a}] = 1$.

Case: β is non-reductive. By Lemma 2.8(ii),

$$[\check{U}_{\beta, m_\beta+a-1}, \check{U}_{\alpha, m_\alpha+r-a}] \subseteq \prod_{\substack{p,q \in \mathbb{Z}_{\geq 1} \\ p\alpha+q\beta \in \Phi}} \check{U}_{p\alpha+q\beta, p(m_\alpha+r-a)+q(m_\beta+a-1)},$$

and the proof can be finished exactly as in the “ β non-reductive”-case of part (a). \square

2.10. Regularity of characters. Recall the notation \mathcal{T} from Section 2.6. Consider the norm map $N_\sigma^m : \mathcal{T}(\overline{\mathbb{F}}_q)^{\sigma^m} \rightarrow \mathcal{T}(\overline{\mathbb{F}}_q)^\sigma = \mathcal{T}(\mathbb{F}_q)$ given by $t \mapsto t\sigma(t) \cdots \sigma^{m-1}(t)$. Let $r \in \mathbb{Z}_{\geq 1}$ be fixed. Following Lusztig [Lus04, 1.5], we say a character $\chi : \mathcal{T}(\mathbb{F}_q) \rightarrow \overline{\mathbb{Q}}_\ell^\times$ is *regular* if for any $\alpha \in \Phi$ and any $m \geq 1$ such that $\sigma^m(\alpha) = \alpha$, the restriction of $\chi \circ N_\sigma^m$ to $\mathcal{T}^\alpha(\overline{\mathbb{F}}_q)^{\sigma^m}$ is non-trivial. A character χ of \check{T}_r^σ is called *regular* if its restriction $\chi|_{\mathcal{T}(\mathbb{F}_q)}$ is regular.

Let $\theta : T(k) \rightarrow \overline{\mathbb{Q}}_\ell^\times$ be a character of level $r - 1$; that is, θ is trivial on $\check{T}^{(r-1)+} \cap T(k)$ but nontrivial on $\check{T}^{(r-2)+}$. Its restriction to $\check{T}^0 \cap T(k)$ can be viewed as a character χ of $\check{T}_r^\sigma = (\check{T}^0 / \check{T}^{(r-1)+})^\sigma$. We say θ is *regular* if χ is.

Remark 2.10. When G is an inner form of $\mathrm{GL}_n(K)$ and T is a maximal nonsplit unramified torus, then $T(k) \cong L^\times$, where L is the degree- n unramified extension of k . If $\theta : L^\times \rightarrow \overline{\mathbb{Q}}_\ell^\times$ is a smooth character trivial on $(\check{T}^r)^\sigma = U_L^r = 1 + \varpi^r \mathcal{O}_L$, then θ being *regular* is the same as being *primitive* in the sense of Boyarchenko–Weinstein [BW16, Section 7.1]. This is closely related to θ being *minimal admissible* in the sense of Bushnell–Henniart [BH05, Section 1.1]. We refer to [CI18, Remark 12.1] for a more precise comparison. \diamond

3. THE SCHEME Σ

We use notation from Section 2. We fix a point $\mathbf{x} \in \mathcal{B}_k$, an integer $r \geq 1$, and two maximal tori T, T' of G defined over k , split over \check{k} , and such that $\mathbf{x} \in \mathcal{A}_{T,\check{k}} \cap \mathcal{A}_{T',\check{k}}$. Further, we fix choose pairs of unipotent radicals of opposite Borels U, U^- (attached to T) and U', U'^- (attached to T') in $G_{\check{k}}$. The construction from Section 2.6, this gives the \mathbb{F}_q -groups $\mathbb{G}_r, \mathbb{T}_r, \mathbb{U}_r, \mathbb{U}_r^-, \mathbb{T}'_r, \mathbb{U}'_r, \mathbb{U}'_r^-$.

3.1. Definition of Σ, Σ_w . Attached to $(T, U), (T', U')$, we consider the following locally closed reduced subscheme of $\sigma(\mathbb{U}_r) \times \sigma(\mathbb{U}'_r) \times \mathbb{G}_r$ whose $\overline{\mathbb{F}}_q$ -points are given by

$$\Sigma(\overline{\mathbb{F}}_q) := \left\{ (x, x', y) \in \sigma(\check{U}_r) \times \sigma(\check{U}'_r) \times \check{G}_r : x\sigma(y) = yx' \right\}.$$

The scheme Σ decomposes into a disjoint union of locally closed subsets, $\Sigma = \coprod_{w \in W_{\mathbf{x}}(T, T')} \Sigma_w$, where Σ_w is the reduced subscheme of Σ , whose $\overline{\mathbb{F}}_q$ -points are given by

$$\Sigma_w(\overline{\mathbb{F}}_q) := \{(x, x', y) \in \Sigma(\overline{\mathbb{F}}_q) : y \in \mathbb{G}_{r,w}(\overline{\mathbb{F}}_q)\},$$

where $\mathbb{G}_{r,w}$ is as in Section 2.8. The group $\check{T}_r^\sigma \times \check{T}_r'^\sigma$ acts on Σ and each Σ_w by

$$(t, t') : (x, x', y) \mapsto (txt^{-1}, t'x't'^{-1}, tyt'^{-1}).$$

The following lemma is completely analogous to [Lus04, Lemma 1.4].

Lemma 3.1. *Let $r \geq 2$ and let $\theta : \check{T}_r^\sigma \rightarrow \overline{\mathbb{Q}}_\ell^\times$, $\theta' : \check{T}_r'^\sigma \rightarrow \overline{\mathbb{Q}}_\ell^\times$ be characters such that $H_c^j(\Sigma)_{\theta^{-1}, \theta'} \neq 0$ for some $j \in \mathbb{Z}$. Then there exist $n \geq 1$ and $g \in N_{\mathbb{G}_r}(\mathbb{T}'_r, \mathbb{T}_r)^{\sigma^n}$ such that $\text{Ad}(g)$ carries $\theta|_{\mathcal{T}^\sigma} \circ N_{\sigma}^{\sigma^n}$ to $\theta'|_{\mathcal{T}'^\sigma} \circ N_{\sigma}^{\sigma^n}$.*

Proof. The proof of [Lus04] applies. The only point where one must be careful is the claim that \mathcal{T} and \mathcal{T}' centralize \mathbb{G}_r^1 (this is used to extend the action of $\mathcal{T}(\mathbb{F}_q) \times \mathcal{T}'(\mathbb{F}_q)$ on a covering of Σ_w to an action of a connected group). Passing to $\overline{\mathbb{F}}_q$ -points, this is the claim that the subgroups $\check{T}^{(r-2)+}/\check{T}^{(r-1)+} = \check{T}^{(r-1)}/\check{T}^{(r-1)+}$ and $\check{T}'^{(r-2)+}/\check{T}'^{(r-1)+} = \check{T}'^{(r-1)}/\check{T}'^{(r-1)+}$ centralize $\check{P}_{\mathbf{x}}^{0+}/\check{P}_{\mathbf{x}}^{(r-1)+}$. By general properties of the Moy–Prasad filtration, $[P_{\mathbf{x}}^{0+}, P_{\mathbf{x}}^{(r-1)}] \subseteq P_{\mathbf{x}}^{(r-1)+}$, which verifies the claim. \square

3.2. Alternating sum of cohomology of Σ . Our main result about Σ is the following generalization of [Lus04, Lemma 1.9].

Theorem 3.2. *Let θ and θ' be characters of \check{T}_r^σ and $\check{T}_r'^\sigma$ respectively, and assume that θ is regular. Then*

$$\sum_{i \in \mathbb{Z}} \dim H_c^i(\Sigma, \overline{\mathbb{Q}}_\ell)_{\theta^{-1}, \theta'} = \#\{w \in W_{\mathbf{x}}(T, T')^\sigma : \theta \circ \text{Ad}(w) = \theta'\}.$$

Proof. Since $\Sigma = \bigsqcup_w \Sigma_w$, it is enough to show that $\sum_{i \in \mathbb{Z}} (-1)^i \dim H_c^i(\Sigma_w, \overline{\mathbb{Q}}_\ell)_{\theta^{-1}, \theta'}$ is 1 if $w \in W_{\mathbf{x}}(T, T')^\sigma$ and $\theta \circ \text{Ad}(w) = \theta'$, and is 0 otherwise. Fix a $w \in W_{\mathbf{x}}(T, T')$. Let $\widehat{\Sigma}_w$ be the locally closed reduced subscheme of $\sigma(\mathbb{U}_r) \times \sigma(\mathbb{U}'_r) \times \mathbb{U}_r \times \mathbb{U}'_r \times \mathbb{K}_r^1 \times \mathbb{T}'_r$ such that

$$\begin{aligned} \widehat{\Sigma}_w(\overline{\mathbb{F}}_q) &= \{(x, x', u, u', z, \tau') \in \sigma(\check{U}_r) \times \sigma(\check{U}'_r) \times \check{U}_r \times \check{U}'_r \times \check{K}_r^1 \times \check{T}'_r : \\ &\quad x\sigma(uz\dot{w}\tau'u') = uz\dot{w}\tau'u'x'\}, \end{aligned}$$

and define an action of $\check{T}_r^\sigma \times \check{T}_r'^\sigma$ by:

$$(t, t') : (x, x', u, u', z, \tau') \mapsto (txt^{-1}, t'x't'^{-1}, tut^{-1}, t'u't'^{-1}, tzt^{-1}, \dot{w}^{-1}t\dot{w}\tau't'^{-1}). \quad (3.1)$$

As the projection $\widehat{\Sigma}_w \rightarrow \Sigma_w$ is Zariski-locally trivial fibration (as being Zariski-locally trivial is preserved under base change, and $\Sigma_w = \Sigma \times_{\mathbb{G}_r} \mathbb{G}_{r,w}$, this follows from Lemma 2.7), the alternating sum of the cohomology does not change if we pass from Σ_w to $\widehat{\Sigma}_w$. Thus to finish the proof of the theorem it is enough to show that

$$\sum_{i \in \mathbb{Z}} (-1)^i \dim H_c^i(\widehat{\Sigma}_w, \overline{\mathbb{Q}}_\ell)_{\theta^{-1}, \theta'} = \begin{cases} 1 & \text{if } w \in W_{\mathbf{x}}(T, T')^\sigma \text{ and } \theta \circ \text{Ad}(w) = \theta', \\ 0 & \text{otherwise.} \end{cases} \quad (3.2)$$

We make the change of variables replacing $x\sigma(u)$ by x and $x'\sigma(u')^{-1}$ by x' , and rewrite $\widehat{\Sigma}_w(\overline{\mathbb{F}}_q)$ as

$$\widehat{\Sigma}_w(\overline{\mathbb{F}}_q) = \{(x, x', u, u', z, \tau') \in \sigma(\check{U}_r) \times \sigma(\check{U}'_r) \times \check{U}_r \times \check{U}'_r \times \check{K}_r^1 \times \check{T}'_r : x\sigma(z\dot{w}\tau') = uz\dot{w}\tau'u'x'\},$$

and the torus action is still given by (3.1). Define a partition into locally closed subsets $\widehat{\Sigma}_w = \widehat{\Sigma}'_w \sqcup \widehat{\Sigma}''_w$ by

$$\begin{aligned}\widehat{\Sigma}'_w(\overline{\mathbb{F}}_q) &= \{(x, x', u, u', z, \tau') \in \widehat{\Sigma}_w(\overline{\mathbb{F}}_q) : z \neq 1\}, \\ \widehat{\Sigma}''_w(\overline{\mathbb{F}}_q) &= \{(x, x', u, u', z, \tau') \in \widehat{\Sigma}_w(\overline{\mathbb{F}}_q) : z = 1\}.\end{aligned}$$

Both subsets are stable under the $\check{T}_r^\sigma \times \check{T}_r'^\sigma$ -action. Now,

$$\sum_{i \in \mathbb{Z}} (-1)^i \dim H_c^i(\widehat{\Sigma}''_w, \overline{\mathbb{Q}}_\ell)_{\theta^{-1}, \theta'} = \begin{cases} 1 & \text{if } w \in W_{\mathbf{x}}(T, T')^\sigma \text{ and } \theta \circ \text{Ad}(w) = \theta', \\ 0 & \text{otherwise,} \end{cases}$$

has literally the same proof as the claim (b) in the proof of [Lus04, Lemma 1.9]. Further, we show in Section 3.6 (after some preparations in Sections 3.3-3.5), under the assumption that θ is regular, that

$$\sum_{i \in \mathbb{Z}} (-1)^i \dim H_c^i(\widehat{\Sigma}'_w, \overline{\mathbb{Q}}_\ell)_{\theta^{-1}, \theta'} = 0, \quad (3.3)$$

so (3.2) holds. \square

3.3. Filtration of \mathbb{G}_{a+1}^a . The main difference of the present article to [Lus04], is that \mathbb{G}_2^1 is not abelian if (and only if) $P_{\mathbf{x}}$ is not reductive, i.e., if \mathbf{x} is not a hyperspecial point. To deal with this problem, we need a refinement of the filtration of \mathbb{G}_r^1 by its subgroups \mathbb{G}_r^a for $1 \leq a \leq r-1$. For $a \geq 1$, we define a filtration of \mathbb{G}_{a+1}^a as follows: let

$$H(1) := \text{subgroup of } \mathbb{G}_{a+1}^a \text{ generated } \mathbb{T}_{a+1}^a \text{ and } \mathbb{U}_{\alpha, a+1}^a \text{ for all reductive } \alpha \in \Phi,$$

and for all $0 \leq \varepsilon < 1$, let

$$H(\varepsilon) := \text{subgroup of } \mathbb{G}_{a+1}^a \text{ generated by } H(1) \text{ and all } \mathbb{U}_{\alpha, a+1}^a \text{ for } \alpha \in \Phi, \text{ satisfying } \varepsilon_\alpha \geq \varepsilon.$$

Note that $\mathbb{T}_{a+1}^a \subseteq H(1) \subseteq H(\varepsilon') \subseteq H(\varepsilon) \subseteq \mathbb{G}_{a+1}^a$ for all $1 > \varepsilon' \geq \varepsilon > 0$. Moreover, there are only finitely many values of ε (“jumps”) satisfying $H(\varepsilon) \supsetneq \bigcup_{\varepsilon' > \varepsilon} H(\varepsilon')$. We denote these jumps by $1 =: \varepsilon_{s+1} > \varepsilon_s > \dots > \varepsilon_1 > 0$ for some $s \geq 0$ (thus 1 is a jump by definition). The jumps are independent of a . We have $H(\varepsilon_1) = \mathbb{G}_{a+1}^a$. For $a \leq r-1$, let $p: \mathbb{G}_r^a \rightarrow \mathbb{G}_{a+1}^a$ be the natural projection, and for $s+1 \geq i \geq 1$, put

$$\mathbb{G}_r^{a,i} := p^{-1}(H(\varepsilon_i)).$$

For convenience, we put $\mathbb{G}_r^{a,s+2} := \mathbb{G}_r^{a+1}$. This defines a refinement $\{\mathbb{G}_r^{a,i}\}_{\substack{r-1 \geq a \geq 1 \\ s+2 \geq i \geq 1}}$ of the filtration $\{\mathbb{G}_r^a\}_{r-1 \geq a \geq 1}$ of \mathbb{G}_r^1 , decreasing with respect to the lexicographical ordering on pairs (a, i) . For $s+1 \geq i \geq 1$, let Φ_i be the set of roots “appearing” in $H(\varepsilon_i)/H(\varepsilon_{i+1})$:

$$\Phi_i := \begin{cases} \{\alpha \in \Phi : \varepsilon_\alpha = 0\} & \text{if } i = s+1, \\ \{\alpha \in \Phi : \varepsilon_\alpha = \varepsilon_i\} & \text{if } s \geq i \geq 1. \end{cases}$$

Lemma 3.3. *Let $r \geq 2$ and $r-1 \geq a \geq 1$.*

- (i) *Let $a \geq 2$. Then $\mathbb{G}_r^a/\mathbb{G}_r^{a+1} = \mathbb{G}_{a+1}^a$ is abelian, and in particular, for $s+1 \geq i \geq 1$, $\mathbb{G}_r^{a,i}/\mathbb{G}_r^{a,i+1}$ is abelian.*
- (ii) *Let $a = 1$ and $s+1 \geq i \geq 1$. Then $\mathbb{G}_r^{1,i}$ is normal in \mathbb{G}_r^1 and the quotient $\mathbb{G}_r^{1,i}/\mathbb{G}_r^{1,i+1}$ is abelian.*

Proof. It suffices to prove the assertions on $\overline{\mathbb{F}}_q$ -points. To show (i), notice that if $a \geq 2$, then $[\check{P}_x^{(a-1)+}, \check{P}_x^{(a-1)+}] \subseteq \check{P}_x^{2(a-1)+} \subseteq \check{P}_x^{a+}$, so it follows that $\check{G}_{a+1}^a = \check{P}_x^{(a-1)+} / \check{P}_x^{a+}$ is abelian. To establish (ii), it is enough to show that (with $a = 1$) for any $s + 1 \geq i \geq 1$, $H(\varepsilon_i)$ is normal in \mathbb{G}_2^1 and that $H(\varepsilon_i)/H(\varepsilon_{i+1})$ is abelian. We spend the rest of the proof establishing these two claims. Recall that for $s + 1 \geq i \geq 1$, $H(\varepsilon_i)$ is generated by \mathbb{T}_2^1 and all $\mathbb{U}_{\alpha,2}^1$ with $\alpha \in \bigsqcup_{j=i}^{s+1} \Phi_j$.

We start with $i = s + 1$, i.e. the case $H(\varepsilon_{s+1}) = H(1)$. By Lemma 2.8, $[\check{T}_2^1, \check{U}_{\alpha,2}^1] = 1$. Let $\alpha \in \Phi_{s+1}$ (thus α is reductive) and let $\beta \in \Phi$ be any non-reductive root. Then $[\check{U}_{\alpha,2}^1, \check{U}_{\beta,2}^1]$ is the image in \check{G}_2^1 of

$$[\check{U}_{\alpha, m_\alpha+1}, \check{U}_{\beta, m_\beta}] \subseteq \prod_{\substack{p, q \in \mathbb{Z}_{\geq 1} \\ p\alpha + q\beta \in \Phi}} \check{U}_{p\alpha + q\beta, p(m_\alpha+1) + qm_\beta}. \quad (3.4)$$

Using Lemma 2.6 along with $p \geq 1$, we see that $p(m_\alpha + 1) + qm_\beta \geq m_{p\alpha + q\beta} + 1$. Thus the contribution of $p\alpha + q\beta$ to the commutator lies in $\check{U}_{p\alpha + q\beta, m_{p\alpha + q\beta} + 1}$. From this we deduce $[\mathbb{U}_{\alpha,2}^1, \mathbb{U}_{\beta,2}^1] \subseteq H(1)$. Thus if $x \in \mathbb{U}_{\beta,2}^1$ for any $\beta \in \Phi$, and $y \in \mathbb{U}_{\alpha,2}^1$, then $xyx^{-1} = [x^{-1}, y^{-1}]y \in H(1)$, which shows that $H(1)$ is normal in \mathbb{G}_2^1 . A computation analogous to (3.4) for $\alpha, \beta \in \Phi^+$ both reductive, shows immediately that $[\mathbb{U}_{\alpha,2}^1, \mathbb{U}_{\beta,2}^1] = 1$ and $[\mathbb{T}_2^1, \mathbb{U}_{\alpha,2}^1] = 1$, so $H(1)$ is abelian.

Next, pick some $s \geq i \geq 1$. We show that $H(\varepsilon_i)$ is normal in \mathbb{G}_2^1 . Since we have already established that $H(\varepsilon_{s+1})$ is normal in \mathbb{G}_2^1 , it suffices to check as above that for all (non-reductive) $\alpha \in \Phi$ with $\varepsilon_\alpha \geq \varepsilon_i$ and all non-reductive $\beta \in \Phi$, we have $[\mathbb{U}_{\alpha,2}^1, \mathbb{U}_{\beta,2}^1] \subseteq H(\varepsilon_i)$. Now, $[\check{U}_{\alpha,2}^1, \check{U}_{\beta,2}^1]$ is the image in \check{G}_2^1 of

$$[\check{U}_{\alpha, m_\alpha}, \check{U}_{\beta, m_\beta}] \subseteq \prod_{\substack{p, q \in \mathbb{Z}_{\geq 1} \\ p\alpha + q\beta \in \Phi}} \check{U}_{p\alpha + q\beta, pm_\alpha + qm_\beta}.$$

Now, if $\varepsilon_{p\alpha + q\beta} \geq \varepsilon_i$, then the contribution of $p\alpha + q\beta$ to the commutator is contained in $\mathbb{U}_{p\alpha + q\beta, 2}^1 \subseteq H(\varepsilon_i)$. If $p\alpha + q\beta$ is reductive, the same computation as in (2.2) shows that $\check{U}_{p\alpha + q\beta, pm_\alpha + qm_\beta} \subseteq \check{U}_{p\alpha + q\beta, m_{p\alpha + q\beta}} \rightarrow \check{U}_{p\alpha + q\beta, 2}^1$. It remains to handle the case that $p\alpha + q\beta$ is non-reductive with $\varepsilon_{p\alpha + q\beta} < \varepsilon_i$. If $p\varepsilon_\alpha + q\varepsilon_\beta < 1$, then by Lemma 2.6, $p\varepsilon_\alpha + q\varepsilon_\beta - \varepsilon_{p\alpha + q\beta} = [p\varepsilon_\alpha + q\varepsilon_\beta] = 0$, i.e. $\varepsilon_i > \varepsilon_{p\alpha + q\beta} = p\varepsilon_\alpha + q\varepsilon_\beta \geq p\varepsilon_i$, which is a contradiction. Thus we must have $p\varepsilon_\alpha + q\varepsilon_\beta \geq 1$, whence $pm_\alpha + qm_\beta - m_{p\alpha + q\beta} = [p\varepsilon_\alpha + q\varepsilon_\beta] \geq 1$. Thus $\check{U}_{p\alpha + q\beta, pm_\alpha + qm_\beta} \subseteq \check{U}_{p\alpha + q\beta, m_{p\alpha + q\beta} + 1}$, whose image in \check{G}_2^1 vanishes. We may finally conclude that $[\mathbb{U}_{\alpha,2}^1, \mathbb{U}_{\beta,2}^1] \subseteq H(\varepsilon_i)$, which finishes the proof of normality of $H(\varepsilon_i)$ in \mathbb{G}_2^1 .

For α and β non-reductive with $\varepsilon_\alpha = \varepsilon_\beta = \varepsilon_i$, a similar computation shows that $[\mathbb{U}_{\alpha,2}^1, \mathbb{U}_{\beta,2}^1] \subseteq H(\varepsilon_{i+1})$. Thus $H(\varepsilon_i)/H(\varepsilon_{i+1})$ is abelian. \square

3.4. Pairings induced by the commutator. Let N, N^- be the unipotent radicals of any two opposite Borel subgroups of G which contain T and are defined over \check{k} . (We will specify N to suit our needs in Section 3.6.) For $r - 1 \geq a \geq 1$, let $\mathbb{N}_r, \mathbb{N}_r^-$ and $\mathbb{N}_r^a, \mathbb{N}_r^{-,a}$ be the corresponding subgroups of \mathbb{G}_r and \mathbb{G}_r^a . Let $\Phi^+ = \{\alpha \in \Phi : \mathbb{U}_{\alpha,r} \subseteq \mathbb{N}_r\}$ and $\Phi^- = \Phi \setminus \Phi^+ = \{\alpha \in \Phi : \mathbb{U}_{\alpha,r} \subseteq \mathbb{N}_r^-\}$. For $s + 1 \geq i \geq 1$, set $\Phi_i^+ = \Phi_i \cap \Phi^+$ and $\Phi_i^- = \Phi_i \cap \Phi^-$, and let $\mathbb{N}_r^{1,i} = \mathbb{G}_r^{1,i} \cap \mathbb{N}_r$. We study some pairings induced by the commutator map. Note that the targets of the maps in Lemma 3.4 are abelian by Lemma 3.3.

Lemma 3.4. *Let $r \geq 2$ and $r - 1 \geq a \geq 1$. Let $\alpha \in \Phi$ be a non-reductive root.*

(i) *Let $a \geq 2$. The commutator map induces a bilinear pairing of abelian groups,*

$$\mathbb{U}_{\alpha,r}^{r-a} / \mathbb{U}_{\alpha,r}^{r-a+1} \times \mathbb{N}_r^a / \mathbb{N}_r^{a+1} \rightarrow \mathbb{G}_r^{r-1}, \quad (\bar{\xi}, \bar{x}) \mapsto [\bar{\xi}, \bar{x}].$$

(ii) Let $a = 1$ and $s + 1 \geq i \geq 1$. Assume that $\varepsilon_{-\alpha} = \varepsilon_i$ (thus $\varepsilon_\alpha = 1 - \varepsilon_i$). We have $[\mathbb{U}_{\alpha,r}^{r-1}, \mathbb{N}_r^{1,i}] \subseteq \mathbb{G}_r^{r-1,s+1}$ and $[\mathbb{U}_{\alpha,r}^{r-1}, \mathbb{N}_r^{1,i+1}] = 1$. The commutator map induces a bilinear pairing of abelian groups,

$$\mathbb{U}_{\alpha,r}^{r-1} \times \mathbb{N}_r^{1,i}/\mathbb{N}_r^{1,i+1} \rightarrow \mathbb{G}_r^{r-1,s+1}, \quad (\xi, \bar{x}) \mapsto [\bar{\xi}, \bar{x}].$$

Proof. (i): By Lemma 2.9 applied three times, the commutator map $\mathbb{U}_{\alpha,r}^{r-a} \times \mathbb{N}_r^a \rightarrow \mathbb{G}_r$ induces the claimed pairing. It is linear in \bar{x} : if $x_1, x_2 \in \check{N}_r^a$, then

$$[\xi, x_1 x_2] = \xi^{-1} x_2^{-1} x_1^{-1} \xi x_1 x_2 = \xi^{-1} x_1^{-1} x_2^{-1} \xi x_2 x_1 = \xi^{-1} x_1^{-1} \xi [\xi, x_2] x_1 = [\xi, x_1] [\xi, x_2],$$

where the second equality follows from Lemma 2.9 and $\mathbb{N}_r^a/\mathbb{N}_r^{a+1}$ being abelian, and the fourth follows from Lemma 2.9 as $[\xi, x_2] \in \check{G}_r^{r-1}$, the assumption $a \geq 2$, and the subsequent fact that \mathbb{N}_{a+1}^a is generated by root subgroups contained in it. The linearity in $\bar{\xi}$ is shown similarly.

(ii): We work on $\overline{\mathbb{F}}_q$ -points. To show the first claim, we observe that $\mathbb{U}_{\alpha,r}^{r-1}$ commutes with \mathbb{N}_r^2 by Lemma 2.9. As $\mathbb{N}_r^{1,i+1}$ is generated by \mathbb{N}_r^2 along with $\mathbb{U}_{\beta,r}^1$ for all β which are either reductive or satisfy $\varepsilon_\beta \geq \varepsilon_i$, we have to show that $[\mathbb{U}_{\alpha,r}^{r-1}, \mathbb{U}_{\beta,r}^1] \subseteq \mathbb{G}_r^{r-1,s+1}$ for all such β . We have two cases:

Case: β is non-reductive. We have to show that $[\check{U}_{\alpha,m_\alpha+r-2}, \check{U}_{\beta,m_\beta}]$ maps to $\check{G}_r^{r-1,s+1}$ inside \check{G}_r . Using Lemma 2.8(ii), it is enough to show that for all $p, q \in \mathbb{Z}_{\geq 1}$ such that $p\alpha + q\beta \in \Phi$, $\check{U}_{p\alpha+q\beta, p(m_\alpha+r-2)+qm_\beta}$ maps to 1 in \check{G}_r if $p\alpha + q\beta$ is non-reductive and maps to $\check{U}_{p\alpha+q\beta, r}^{r-1}$ if $p\alpha + q\beta$ is reductive. In both cases, this amounts to the claim that

$$p(m_\alpha + r - 2) + qm_\beta \geq m_{p\alpha+q\beta} + r - 1,$$

which in turn by Lemma 2.6 is equivalent to

$$[p\varepsilon_\alpha + q\varepsilon_\beta] + (p-1)(r-2) \geq 1,$$

which is true as $\varepsilon_\beta \geq \varepsilon_i = \varepsilon_{-\alpha} = 1 - \varepsilon_\alpha$.

Case: β is reductive. This case is shown similarly (in fact, slightly simpler) to the above, and we omit the details. This finishes the proof of the first claim, i.e., $[\mathbb{U}_{\alpha,r}^{r-1}, \mathbb{N}_r^{1,i}] \subseteq \mathbb{G}_r^{r-1,s+1}$.

We now show the second claim, i.e., $[\mathbb{U}_{\alpha,r}^{r-1}, \mathbb{N}_r^{1,i+1}] = 1$. Proceeding analogously as in the proof of the first claim, we need only to show that for all $\beta \in \Phi$ either reductive or satisfying $\varepsilon_\beta \geq \varepsilon_{i+1}$, one has $[\mathbb{U}_{\alpha,r}^{r-1}, \mathbb{U}_{\beta,r}^1] = 1$. We again have two cases:

Case: β is non-reductive. We have to show that $[\check{U}_{\alpha,m_\alpha+r-2}, \check{U}_{\beta,m_\beta}]$ maps to 1 in \check{G}_r . Using Lemma 2.8(ii), it is enough to show that for all $p, q \in \mathbb{Z}_{\geq 1}$ such that $p\alpha + q\beta \in \Phi$, $\check{U}_{p\alpha+q\beta, p(m_\alpha+r-2)+qm_\beta}$ maps to 1 in \check{G}_r . If $p\alpha + q\beta$ is non-reductive, this follows from the similar statement in the proof of the first claim, as $\varepsilon_{i+1} \geq \varepsilon_i$. If $p\alpha + q\beta$ is reductive, it amounts to claim that

$$p(m_\alpha + r - 2) + qm_\beta \geq m_{p\alpha+q\beta} + r,$$

which by Lemma 2.6 is equivalent to

$$[p\varepsilon_\alpha + q\varepsilon_\beta] + (p-1)(r-2) \geq 2,$$

But this is true, as $[p\varepsilon_\alpha + q\varepsilon_\beta] \geq 2$. Indeed, as $p\alpha + q\beta$ is reductive, $\varepsilon_{p\alpha+q\beta} = 0$. Hence by Lemma 2.6 $[p\varepsilon_\alpha + q\varepsilon_\beta] = p\varepsilon_\alpha + q\varepsilon_\beta \geq \varepsilon_\alpha + \varepsilon_\beta > 1$. Being an integer, $[p\varepsilon_\alpha + q\varepsilon_\beta]$ must be ≥ 2 .

Case: β is reductive. This case is shown similarly (in fact, slightly simpler) to the above, and we omit the details. This finishes the proof of the second claim.

We are now ready to show that the claimed pairing is well-defined. Indeed, let $\xi \in \check{U}_{\alpha,r}^{r-1}$ and let $x, x' \in \check{N}_r^{1,i}$ with the same image $\bar{x} = \bar{x}' \in \check{N}_r^{1,i}/\check{N}_r^{1,i+1}$. Then there is an $y \in \check{N}_r^{1,i+1}$ such that $x' = xy$. We compute:

$$[\xi, x'] = [\xi, xy] = \xi^{-1}y^{-1}x^{-1}\xi xy = y^{-1}[\xi, x]y = [\xi, x],$$

where for the third equality we use that $[\mathbb{U}_{\alpha,r}^{r-1}, \mathbb{N}_r^{1,i+1}] = 1$ and for the last we use that $[\xi, x] \in \check{G}_r^{r-1,s+1}$ and $[\mathbb{G}_r^{r-1,s+1}, \mathbb{N}_r^1] = 1$ (indeed, for any reductive root γ we have $[\mathbb{U}_{\gamma,r}^{r-1}, \mathbb{N}_r^1] = 1$ by Lemma 2.9). Now we show that this pairing is linear in the second variable. Therefore, let $\xi \in \check{U}_{\alpha,r}^{r-1}$ and $x_1, x_2 \in \check{N}_r^{1,i}$. We compute:

$$\begin{aligned} [\xi, x_1x_2] &= \xi^{-1}x_2^{-1}x_1^{-1}\xi x_1x_2 = \xi^{-1}[x_2, x_1]x_1^{-1}x_2^{-1}\xi x_1x_2 \\ &= [x_2, x_1]\xi^{-1}x_1^{-1}x_2^{-1}\xi x_1x_2 = [x_2, x_1]\xi^{-1}x_1^{-1}x_2^{-1}\xi x_2x_1[x_1, x_2] \\ &= [x_2, x_1]\xi^{-1}x_1^{-1}\xi[\xi, x_2]x_1[x_1, x_2] = [x_2, x_1][\xi, x_1][\xi, x_2][x_1, x_2] \\ &= [\xi, x_1][\xi, x_2]. \end{aligned}$$

The third equality follows as $[x_2, x_1] \in \check{N}_r^{1,i+1}$ (as $\mathbb{N}_r^{1,i}/\mathbb{N}_r^{1,i+1}$ is abelian) and as $[\mathbb{U}_{\alpha,r}^{r-1}, \mathbb{N}_r^{1,i+1}] = 1$. The sixth equality follows as $[\xi, x_2] \in \check{G}_r^{r-1,s+1}$ commutes with $x_1 \in \check{N}_r^1$. The last equality follows as $[\xi, x_1], [\xi, x_2] \in \check{G}_r^{r-1,s+1}$ commute with $[x_1, x_2] \in \check{N}_r^1$, and as $[x_2, x_1][x_1, x_2] = 1$. An analogous (slightly simpler) computation shows the linearity in the first variable. \square

Remark 3.5. Lemma 2.8(ii) can certainly be generalized. As we will not use the following generalization, we state it without proof. As for any root $\alpha \in \Phi$, $-\alpha$ is a root too, and $\varepsilon_{-\alpha} = 1 - \varepsilon_\alpha$, we have a symmetry between the jumps ε_i . Concretely, we have $\varepsilon_i = 1 - \varepsilon_{s+1-i}$ for $1 \leq i \leq s$. For each $1 \leq a \leq r-1$, let $\mathbb{G}_r^{a,i}$ be the subgroup of \mathbb{G}_r^a generated by \mathbb{G}_r^{a+1} , \mathbb{T}_r^a , $\mathbb{U}_{\alpha,r}^a$ (α reductive or $\varepsilon_\alpha \geq \varepsilon_i$). Then Lemma 2.8 extends to the following general duality statement: Fix $1 \leq a \leq r-1$ and $1 \leq i \leq s$. Then the commutator induces a bilinear pairing,

$$\mathbb{G}_r^{r-a,s+1-i}/\mathbb{G}_r^{r-a,s+2-i} \times \mathbb{G}_r^{a,i}/\mathbb{G}_r^{a,i+1} \rightarrow \mathbb{G}_r^{r-1,s+1}. \quad \diamond$$

3.5. Stratification on (subgroups of) \mathbb{N}_r^1 . Recall that for any subgroup $H \subset G$ and associated subgroups $\mathbb{H}_r \subset \mathbb{G}_r$, we have the notation $\mathbb{H}_r^{a,*} = \mathbb{H}_r^a \setminus \mathbb{H}_r^{a+1}$ (open subscheme) and hence the corresponding set $\check{H}_r^{a,*}$ of $\bar{\mathbb{F}}_q$ -valued points.

Lemma 3.6. *Let $r \geq 2$ and let $r-1 \geq a \geq 1$. For $z \in \check{N}_r^{a,*}$, write $z = \prod_{\beta \in \Phi^+} x_\beta^z$ with $x_\beta^z \in \check{U}_{\beta,r}^a$ for a fixed (but arbitrary) order on Φ^+ . For $\beta \in \Phi^+$, let $a \leq a(\beta, z) \leq r$ be the integer such that $x_\beta^z \in \check{U}_{\beta,r}^{a(\beta,z),*}$.*

(i) *If $a \geq 2$, then the set*

$$A_z := \{\beta \in \Phi^+ : a(\beta, z) = a\}$$

is non-empty and independent of the chosen order on Φ^+ .

(ii) *Let $a = 1$ and let $s+1 \geq i \geq 1$ be such that $z \in \check{N}_r^{1,i,*}$. Then the set*

$$A_z := \{\beta \in \Phi_i^+ : a(\beta, z) = 1\}$$

is non-empty and independent of the chosen order on Φ^+ . Moreover, $a(\beta, z) > 1$ for all $\beta \in \bigcup_{j=1}^{i-1} \Phi_j^+$.

Proof. (i): As $a \geq 2$, the quotient $\mathbb{N}_r^a / \mathbb{N}_r^{a+1}$ is abelian by Lemma 3.3. Thus its $\overline{\mathbb{F}}_q$ -points are simply tuples $(\bar{x}_\beta)_{\beta \in \Phi^+}$ with $\bar{x}_\beta \in \check{U}_{\beta, a+1}^a$ with entry-wise multiplication. If $\bar{z} = (\bar{x}_\beta^z)$ is the image of z in this quotient, then A_z identifies with the set of those β for which $\bar{x}_\beta^z \neq 1$ (which is obviously independent of the order).

(ii): Assume that the last claim of (ii) is not true. Then let $1 \leq i_0 < i$ be the smallest integer such that $a(\beta, z) = 1$ for some $\beta \in \Phi_{i_0}^+$. Then from Lemma 3.3 it follows that $z \in \check{N}_r^{1, i_0, *}$, which contradicts the assumption. This shows the last claim. The first claim follows by the same argument as in (i). \square

Using Section 3.4 we can now prove the following generalization of [Lus04, Lemma 1.7].

Definition 3.7. For $\alpha \in \Phi^+$ define its *height* $\text{ht}(\alpha)$ (relative to N) to be the largest integer $m \geq 1$ such that $\alpha = \sum_{i=1}^m \alpha_i$ with $\alpha_i \in \Phi^+$.

Proposition 3.8. Let $r \geq 2$ and let $r-1 \geq a \geq 1$. Let $z = \prod_{\beta \in \Phi^+} x_\beta^z \in \check{N}_r^{a, *}$ for $x_\beta^z \in \check{U}_{\beta, r}^a$ and let A_z be as in Lemma 3.6.

(i) If A_z contains a non-reductive root, let $-\alpha \in A_z$ be a non-reductive root of maximal height and $\alpha \in \Phi^-$ its opposite. Then for any $\xi \in \mathbb{U}_{\alpha, r}^{r-a}$, we have $[\xi, z] \in \mathcal{T}^\alpha \mathbb{N}_r^{-, r-1}$. Moreover, projecting $[\xi, z]$ into \mathcal{T}^α induces an isomorphism

$$\lambda_z: \mathbb{U}_{\alpha, r}^{r-a} / \mathbb{U}_{\alpha, r}^{r-a+1} \xrightarrow{\sim} \mathcal{T}^\alpha$$

(ii) If A_z contains only reductive roots, let $-\alpha \in A_z$ be a root of maximal height and $\alpha \in \Phi^-$ its opposite. Then for any $\xi \in \mathbb{U}_{\alpha, r}^{r-a-1}$, we have $[\xi, z] \in \mathcal{T}^\alpha \mathbb{N}_r^{-, r-1}$. Moreover, projecting $[\xi, z]$ into \mathcal{T}^α induces an isomorphism

$$\lambda_z: \mathbb{U}_{\alpha, r}^{r-a-1} / \mathbb{U}_{\alpha, r}^{r-a} \xrightarrow{\sim} \mathcal{T}^\alpha$$

Proof. Parts (i) and (ii) can be proven in the same way. We give the full proof of (i) only.

Proof of (i) when $a \geq 2$. We work on $\overline{\mathbb{F}}_q$ -points. Assume that A_z contains a non-reductive root and let $-\alpha$ be such a root of maximal height and $\alpha \in \Phi^-$ its opposite. Let $\xi \in \check{U}_{\alpha, r}^{r-a}$ and let $\bar{\xi} \in \check{U}_{\alpha, r}^{r-a} / \check{U}_{\alpha, r}^{r-a+1}$ and $\bar{z} \in \check{N}_r^a / \check{N}_r^{a+1}$ be the images of ξ and z respectively. By Lemma 3.3 we may write

$$\bar{z} = \bar{x}_{-\alpha}^z \prod_{\beta \in \Phi^+ \text{ red.}} \bar{x}_\beta^z \cdot \prod_{\substack{\beta \in \Phi^+ \text{ non-red.}, \\ \text{ht}(\beta) \leq \text{ht}(-\alpha)}} \bar{x}_\beta^z,$$

where $\bar{x}_\beta^z \in \check{U}_{\beta, r}^a / \check{U}_{\beta, r}^{a+1}$ and where the products are taken in any order. Lemma 3.4 shows that $[\bar{\xi}, z]$ is the product of $[\bar{\xi}, \bar{x}_{-\alpha}^z]$ with all the $[\bar{\xi}, \bar{x}_\beta^z]$ for $\beta \in \Phi^+$, the product taken in any order. If β is reductive, then $[\bar{\xi}, \bar{x}_\beta^z] \in [\check{U}_{\alpha, r}^{r-a}, \check{U}_{\beta, r}^a] = 1$ by Lemma 2.9. If $\beta \neq -\alpha$ is non-reductive, then by assumption $\text{ht}(\beta) \leq \text{ht}(-\alpha)$. The commutator $[\bar{\xi}, \bar{x}_\beta^z]$ is the image of an element of

$$[\check{U}_{\alpha, m_\alpha + (r-a) - 1}, \check{U}_{\beta, m_\beta + a - 1}] \subseteq \prod_{\substack{p, q \in \mathbb{Z}_{\geq 1} \\ p\alpha + q\beta \in \Phi}} \check{U}_{p\alpha + q\beta, pm_\alpha + qm_\beta + p(r-a-1) + q(a-1)} \quad (3.5)$$

Lemma 3.9. The image of the right hand side of (3.5) in \check{G}_r lies in $\check{N}_r^{-, r-1}$.

Proof. It is enough to show that for each (p, q) occurring in the product, the corresponding factor is either contained in $\check{N}_r^{-, r-1}$ or vanishes in \check{G}_r . If $p \geq q$, then $\text{ht}(\beta) \leq \text{ht}(-\alpha)$ implies $p\alpha + q\beta \notin \Phi^+$. So, we may assume that $q > p$ and in particular $q \geq 2$. It is enough to show that

$$\check{U}_{p\alpha+q\beta, pm_\alpha+qm_\beta+p(r-a-1)+q(a-1)} \subseteq \begin{cases} \check{U}_{p\alpha+q\beta, m_{p\alpha+q\beta}+r} & \text{if } p\alpha + q\beta \text{ reductive} \\ \check{U}_{p\alpha+q\beta, m_{p\alpha+q\beta}+r-1} & \text{otherwise,} \end{cases}$$

as both map to 1 in \check{G}_r . Equivalently, we have to show that

$$pm_\alpha + qm_\beta - m_{p\alpha+q\beta} + p(r-a-1) + q(a-1) - (r-1) \geq \begin{cases} 1 & \text{if } p\alpha + q\beta \text{ reductive} \\ 0 & \text{otherwise.} \end{cases}$$

But this holds as by Lemma 2.6, $pm_\alpha + qm_\beta - m_{p\alpha+q\beta} = \lfloor p\varepsilon_\alpha + q\varepsilon_\beta \rfloor$ is ≥ 1 if $p\alpha + q\beta$ is reductive and is ≥ 0 otherwise, and as $q \geq 2$ and $a \geq 2$. \square

Finally, $[\bar{\xi}, \bar{x}_{-\alpha}^z] = [\xi, x_{-\alpha}^z] \in \mathcal{T}^\alpha(\overline{\mathbb{F}}_q)$ by Lemma 2.8(iii). Thus $[\xi, z] \in \mathcal{T}^\alpha(\overline{\mathbb{F}}_q)\check{N}_r^{-, r-1}$. Moreover, if we project onto $\mathcal{T}^\alpha(\overline{\mathbb{F}}_q)$, then only $[\bar{\xi}, \bar{x}_{-\alpha}^z]$ survives and Lemma 2.8(iii) proves the desired isomorphism λ_z . This finishes the proof of (i) in the case $a \geq 2$.

Proof of (i) when $a = 1$. Let $s \geq i \geq 1$ denote the integer such that $z \in \check{N}_r^{1, i, *}$. (Note that $i \neq s + 1$ as A_z contains a non-reductive root by assumption). We have $\xi \in \check{U}_{\alpha, r}^{r-1}$, and we let \bar{z} denote the image of z in $\check{N}_r^{1, i} / \check{N}_r^{1, i+1}$. By Lemma 3.3 we may write

$$\bar{z} = \bar{x}_{-\alpha}^z \left(\prod_{\substack{\beta \in \Phi_i^+ : \beta \neq -\alpha \\ \text{ht}(\beta) \leq \text{ht}(-\alpha)}} \bar{x}_\beta^z \right),$$

(product are taken in any order). By Lemma 3.4, $[\xi, \bar{z}]$ is the product of $[\xi, \bar{x}_{-\alpha}^z]$ with all the $[\xi, \bar{x}_\beta^z]$ taken in any order. By assumption $\varepsilon_\beta = \varepsilon_i = \varepsilon_{-\alpha} = 1 - \varepsilon_\alpha$. In particular, all β 's are non-reductive. Now, $[\xi, \bar{x}_\beta^z]$ is the image in $\check{G}_r^{r-1, s+1}$ of an element of

$$[\check{U}_{\alpha, m_\alpha+r-2}, \check{U}_{\beta, m_\beta}] \subseteq \prod_{\substack{p, q \in \mathbb{Z}_{\geq 1} \\ p\alpha+q\beta \in \Phi}} \check{U}_{p\alpha+q\beta, pm_\alpha+qm_\beta+p(r-2)} \quad (3.6)$$

Lemma 3.10. *The image of the right hand side of (3.6) in \check{G}_r lies in $\check{N}_r^{-, r-1}$.*

Proof. Note that the right hand side of (3.6) is contained in $\check{G}_r^{r-1, s+1}$ (exactly as in the proof of Lemma 3.4(ii)). Now the same arguments as in the proof Lemma 3.9 apply. If $p \geq q$, then $\text{ht}(\beta) \leq \text{ht}(-\alpha)$ implies $p\alpha + q\beta \notin \Phi^+$, thus the corresponding factor of the product is contained in $\check{N}_r^{-} \cap \check{G}_r^{r-1, s+1} \subseteq \check{N}_r^{-, r-1}$. Thus we may assume that $q > p$ and in particular $q \geq 2$. It is enough to show that

$$\check{U}_{p\alpha+q\beta, pm_\alpha+qm_\beta+p(r-2)} \subseteq \begin{cases} \check{U}_{p\alpha+q\beta, m_{p\alpha+q\beta}+r} & \text{if } p\alpha + q\beta \text{ is reductive} \\ \check{U}_{p\alpha+q\beta, m_{p\alpha+q\beta}+r-1} & \text{otherwise,} \end{cases}$$

as both map to 1 in \check{G}_r . Equivalently, we have to show that

$$pm_\alpha + qm_\beta - m_{p\alpha+q\beta} + p(r-2) - (r-1) \geq \begin{cases} 1 & \text{if } p\alpha + q\beta \text{ is reductive} \\ 0 & \text{otherwise.} \end{cases}$$

By Lemma 2.6, this follows from $\lfloor p\varepsilon_\alpha + q\varepsilon_\beta \rfloor \geq 2$ if $p\alpha + q\beta$ is reductive, resp. to $\lfloor p\varepsilon_\alpha + q\varepsilon_\beta \rfloor \geq 1$ if $p\alpha + q\beta$ is non-reductive. But in any case we have $p\varepsilon_\alpha + q\varepsilon_\beta \geq \varepsilon_\alpha + 2(1 - \varepsilon_\alpha) = 2 - \varepsilon_\alpha > 1$ by assumptions. In particular, we are done in the case when $p\alpha + q\beta$ is non-reductive. If $p\alpha + q\beta$ is reductive, then $p\varepsilon_\alpha + q\varepsilon_\beta$ must also be an integer (by Lemma 2.6) and hence ≥ 2 , and we are done in this case too. \square

Finally, $[\xi, \bar{x}_{-\alpha}^z] \in \mathcal{T}^\alpha(\overline{\mathbb{F}}_q)$ by Lemma 2.8(iii). Thus $[\xi, z] \in \mathcal{T}^\alpha(\overline{\mathbb{F}}_q)\check{N}_r^{-,r-1}$. Moreover, if we project onto $\mathcal{T}^\alpha(\overline{\mathbb{F}}_q)$, then only $[\bar{\xi}, \bar{x}_{-\alpha}^z]$ survives and Lemma 2.8(iii) proves the desired isomorphism λ_z . This finishes the proof of (i). \square

Remark 3.11. We note that in the proof of [Lus04, Lemma 1.7] there is an (easily correctable) mistake. It is claimed that whenever $-\alpha, \beta \in \Phi^+$ with $-\alpha \neq \beta$ and $\text{ht}(-\alpha) \geq \text{ht}(\beta)$, then $p\alpha + q\beta \notin \Phi^+$ for all $p, q \in \mathbb{Z}_{\geq 1}$. This is not true. For example, let Φ be of type C_2 , let ϵ_1, ϵ_2 denote a basis for $X^*(T)$ such that the $\Phi^+ = \{\epsilon_1 - \epsilon_2, \epsilon_1 + \epsilon_2, 2\epsilon_1, 2\epsilon_2\}$. Then taking $\alpha = -2\epsilon_1$, $\beta = \epsilon_1 + \epsilon_2$. Then $\text{ht}(-\alpha) = 3 > 2 = \text{ht}(\beta)$. But $\alpha + 2\beta = 2\epsilon_2 \in \Phi^+$. Observe here that $\alpha + \beta \notin \Phi^+$, which contradicts the parenthetical assertion at the end of the proof of [Lus04, Lemma 1.7].

Surely, the statement of [Lus04, Lemma 1.7] remains true. The place in its proof, where the abovementioned claim is used, can be corrected as follows: if $p\alpha + q\beta \in \Phi^+$ for some $p, q \in \mathbb{Z}_{\geq 1}$, then $q \geq 2$ and the part of the commutator (as in the proof of Proposition 3.8) inside $\mathbb{U}_{p\alpha+q\beta, r}$ vanishes, since all roots are reductive and $r \geq 2$. \diamond

Let $\mathbb{K}_r = \mathbb{U}_r^- \cap \mathbb{N}_r$. Let $\Phi' = \{\beta \in \Phi^+ : \mathbb{U}_{\beta, r} \subseteq \mathbb{K}_r\}$. Let \mathcal{X} denote the set of all non-empty subsets $I \subseteq \Phi'$ satisfying

- (i) the restriction of $\text{ht} : \Phi^+ \rightarrow \mathbb{Z}_{\geq 0}$ to I is constant, and
- (ii) I contains either only reductive or only non-reductive roots.

To $z \in \check{K}_r^1 \setminus \{1\}$ we attach a pair (a_z, I_z) with $1 \leq a_z \leq r - 1$ and $I_z \in \mathcal{X}$. Define a_z by $z \in \check{K}_r^{a_z, *}$. Let A_z be as in Lemma 3.6. If A_z contains a non-reductive root, let $I_z \subseteq A_z$ be the subset of all non-reductive roots of maximal height. If A_z contains only reductive roots, let $I_z \subseteq A_z$ be the subset of all roots of maximal height. We have a stratification into locally closed subsets

$$\mathbb{K}_r^1 \setminus \{1\} = \bigsqcup_{a, I} \mathbb{K}_r^{a, *, I}, \quad \text{where } \mathbb{K}_r^{a, *, I}(\overline{\mathbb{F}}_q) = \{z \in \check{K}_r^1 \setminus \{1\} : (a_z, I_z) = (a, I)\}. \quad (3.7)$$

3.6. Cohomology of $\widehat{\Sigma}'$. We now complete the proof of Theorem 3.2 by proving Claim (3.3). Using the stratification (3.7) and Proposition 3.8, Claim (3.3) is proven in exactly the same way as in [Lus04, Lemma 1.9]. For convenience, we sketch the arguments here. It is enough to show that $H_c^j(\widehat{\Sigma}'_w)_{\theta, \theta'} = 0$ for all $j \geq 0$. For a $\mathcal{T}'(\overline{\mathbb{F}}_q)^\sigma$ -module M and a character χ of $\mathcal{T}'(\overline{\mathbb{F}}_q)^\sigma$, write $M_{(\chi)}$ for the χ -isotypic component of M . Note that $\mathcal{T}'(\overline{\mathbb{F}}_q)^\sigma$ acts on $\widehat{\Sigma}'_w$ by

$$t' : (x, x', u, u', z, \tau') \mapsto (x, t'x't'^{-1}, u, t'u't'^{-1}, z, \tau't'^{-1}).$$

Hence $H_c^j(\widehat{\Sigma}'_w)$ is a $\mathcal{T}'(\overline{\mathbb{F}}_q)$ -module. It is enough to show that $H_c^j(\widehat{\Sigma}'_w)_{(\chi)} = 0$ for any regular character χ of $\mathcal{T}'(\overline{\mathbb{F}}_q)$. Fix such a χ . Set $N = \dot{w}U'^-\dot{w}^{-1}$, $N^- = \dot{w}U'\dot{w}^{-1}$. The stratification (3.7) of $\mathbb{K}_r^1 \setminus \{1\}$ induces a stratification of $\widehat{\Sigma}'_w$ into locally closed subsets indexed by $1 \leq a \leq r - 1$

and $I \in \mathcal{X}$:

$$\widehat{\Sigma}'_w = \bigsqcup_{a,I} \widehat{\Sigma}'_w{}^{a,I} \quad \text{where } \widehat{\Sigma}'_w{}^{a,I}(\overline{\mathbb{F}}_q) = \{(x, x', u, u', z, \tau') \in \widehat{\Sigma}'_w(\overline{\mathbb{F}}_q) : z \in \check{K}_r^{a,*,I}\}.$$

Note that each $\widehat{\Sigma}'_w{}^{a,I}$ is stable under $\mathcal{T}'(\mathbb{F}_q)$. Thus (3.3) follows from

$$H_c^j(\widehat{\Sigma}'_w{}^{a,I}, \overline{\mathbb{Q}}_\ell)_{(\chi)} = 0 \quad \text{for any fixed } a, I. \quad (3.8)$$

To show (3.8), choose a root α such that $-\alpha \in I$. Then $\mathbb{U}_{\alpha,r} \subseteq \mathbb{U}_r \cap \dot{w}\mathbb{U}'_r\dot{w}^{-1}$. By Proposition 3.8 for any $z \in \check{K}_r^{a,*,I}$, we have an isomorphism

$$\begin{aligned} \lambda_z : \mathbb{U}_{\alpha,r}^{r-a} / \mathbb{U}_{\alpha,r}^{r-a+1} &\xrightarrow{\sim} \mathcal{T}^\alpha, & \text{if } \alpha \text{ is non-reductive,} \\ \lambda_z : \mathbb{U}_{\alpha,r}^{r-a-1} / \mathbb{U}_{\alpha,r}^{r-a} &\xrightarrow{\sim} \mathcal{T}^\alpha, & \text{if } \alpha \text{ is reductive.} \end{aligned}$$

Let π denote the natural projection $\mathbb{U}_{\alpha,r}^{r-a} \rightarrow \mathbb{U}_{\alpha,r}^{r-a} / \mathbb{U}_{\alpha,r}^{r-a+1}$ if α is non-reductive and the natural projection $\mathbb{U}_{\alpha,r}^{r-a-1} \rightarrow \mathbb{U}_{\alpha,r}^{r-a-1} / \mathbb{U}_{\alpha,r}^{r-a}$ if α is reductive. Let ψ be a section to π such that $\pi\psi = 1$ and $\psi(1) = 1$. Let

$$\mathcal{H}' := \{t' \in \mathcal{T}' : t'^{-1}\sigma(t') \in \dot{w}^{-1}\mathcal{T}^\alpha\dot{w}\}.$$

This is a closed subgroup of \mathcal{T}' . For any $t' \in \mathcal{T}'$ define $f_{t'} : \widehat{\Sigma}'_w{}^{a,I} \rightarrow \widehat{\Sigma}'_w{}^{a,I}$ by

$$f_{t'}(x, x', u, u', z, \tau') = (x\sigma(\xi), \hat{x}', u, \sigma(t')^{-1}u'\sigma(t'), z, \tau'\sigma(t')),$$

where

$$\xi = \psi\lambda_z^{-1}(\dot{w}\sigma(t')^{-1}t'\dot{w}^{-1}) \in \begin{cases} \mathbb{U}_{\alpha,r}^{r-a-1} \subseteq \mathbb{U}_r \cap \dot{w}\mathbb{U}'_r\dot{w}^{-1} & \text{if } \alpha \text{ is reductive,} \\ \mathbb{U}_{\alpha,r}^{r-a} \subseteq \mathbb{U}_r \cap \dot{w}\mathbb{U}'_r\dot{w}^{-1} & \text{otherwise,} \end{cases}$$

and $\hat{x}' \in \mathbb{G}_r$ is defined by the condition that

$$x\sigma(\xi z \dot{w} \tau' \sigma(t')) \in u z \dot{w} \tau' \sigma(t') \sigma(t')^{-1} u' \sigma(t') \hat{x}'.$$

To check that $f_{t'}$ is well-defined we have to show $\hat{x}' \in \sigma(\mathbb{U}'_r)$. This is done with exactly the same computation as in the proof of [Lus04, Lemma 1.9], and we omit this. It is clear that $f_{t'} : \widehat{\Sigma}'_w{}^{a,I} \rightarrow \widehat{\Sigma}'_w{}^{a,I}$ is an isomorphism for any $t' \in \mathcal{H}'$. Moreover, since $\mathcal{T}'(\mathbb{F}_q) \subseteq \mathcal{H}'$ and since for any $t' \in \mathcal{T}'(\mathbb{F}_q)$ the map $f_{t'}$ coincides with the action of t' in the $\mathcal{T}'(\mathbb{F}_q)$ -action on $\widehat{\Sigma}'_w{}^{a,I}$ (we use $\psi(1) = 1$ here), it follows that we have constructed an action f of \mathcal{H}' on $\widehat{\Sigma}'_w{}^{a,I}$ extending the $\mathcal{T}'(\mathbb{F}_q)$ -action.

If a connected group acts on a scheme, the induced action in the cohomology is constant. Thus for any $t' \in \mathcal{H}'^\circ$, the induced map $f_{t'}^* : H_c^j(\widehat{\Sigma}'_w{}^{a,I}, \overline{\mathbb{Q}}_\ell) \rightarrow H_c^j(\widehat{\Sigma}'_w{}^{a,I}, \overline{\mathbb{Q}}_\ell)$ is constant when t' varies in \mathcal{H}'° . Hence $\mathcal{T}'(\mathbb{F}_q) \cap \mathcal{H}'^\circ$ acts trivially on $H_c^j(\widehat{\Sigma}'_w{}^{a,I}, \overline{\mathbb{Q}}_\ell)$.

We can find some $m \geq 1$ such that $\sigma^m(\dot{w}^{-1}\mathcal{T}^\alpha\dot{w}) = \dot{w}^{-1}\mathcal{T}^\alpha\dot{w}$. Then

$$t' \mapsto t'\sigma(t')\sigma^2(t') \cdots \sigma^{m-1}(t')$$

defines a morphism $\dot{w}^{-1}\mathcal{T}^\alpha\dot{w} \rightarrow \mathcal{H}'$. Since \mathcal{T}^α is connected, its image is also connected and hence contained in \mathcal{H}'° . If $t' \in (\dot{w}^{-1}\mathcal{T}^\alpha(\overline{\mathbb{F}}_q)\dot{w})^{\sigma^m}$, then $N_\sigma^{\sigma^m}(t') \in \mathcal{T}'(\overline{\mathbb{F}}_q)^\sigma$ and hence also $N_\sigma^{\sigma^m}(t') \in \mathcal{T}'(\overline{\mathbb{F}}_q)^\sigma \cap \mathcal{H}'^\circ(\overline{\mathbb{F}}_q)$. Thus the action of $N_\sigma^{\sigma^m}(t') \in \mathcal{T}'(\overline{\mathbb{F}}_q)^\sigma$ on $H_c^j(\widehat{\Sigma}'_w{}^{a,I})$ is trivial for any $t' \in (\dot{w}^{-1}\mathcal{T}^\alpha(\overline{\mathbb{F}}_q)\dot{w})^{\sigma^m}$.

Finally, observe that if $H_c^j(\widehat{\Sigma}'_w{}^{a,I}, \overline{\mathbb{Q}}_\ell)_{(\chi)} \neq 0$, then the above shows that $t' \mapsto \chi(N_\sigma^{\sigma^m}(t'))$ must be the trivial character, which contradicts the regularity assumption on χ . This establishes (3.8), and hence also (3.3), which was the last outstanding claim in the proof of Theorem 3.2.

4. REPRESENTATIONS OF PARAHORIC SUBGROUPS OF $G(k)$

4.1. **The schemes $S_{T,U}$.** Let the notation be as at the beginning of Section 3. In particular, fix $r \geq 1$, a k -rational \check{k} -split maximal torus T in G , a point $\mathbf{x} \in \mathcal{A}_{T,\check{k}} \cap \mathcal{B}_k$, and the unipotent radical U of a Borel subgroup of $G_{\check{k}}$, defined over \check{k} and containing $T_{\check{k}}$. Let d be the smallest positive integer such that $\sigma^d(U) = U$. To this data, we attach the \mathbb{F}_{q^d} -subscheme of \mathbb{G}_r

$$S_{\mathbf{x},T,U,r} := \{x \in \mathbb{G}_r : x^{-1}\sigma(x) \in \mathbb{U}_r\}.$$

To match the notation of [Lus04], we write $S_{T,U}$ for $S_{\mathbf{x},T,U,r}$.

Lemma 4.1. *$S_{T,U}$ is separated, smooth, and of finite type over \mathbb{F}_{q^d} . Moreover, $\dim S_{T,U} = (r-1)\#\Phi^+ + \#\Phi^{+,\text{red}}$, where Φ^+ and $\Phi^{+,\text{red}}$ are the roots and reductive roots of T in U .*

Proof. Indeed, $S_{T,U}$ is the pullback of \mathbb{U}_r under the finite étale Lang map $\mathbb{G}_r \rightarrow \mathbb{G}_r$, $x \mapsto x^{-1}\sigma(x)$, and $\dim \mathbb{U}_r = (r-1)\#\Phi^+ + \#\Phi^{+,\text{red}}$ is immediate. \square

The finite group $\check{G}_r^\sigma \times \check{T}_r^\sigma$ acts on $S_{T,U}$ by $(g,t): x \mapsto gxt$.

Remark 4.2. $S_{T,U}$ admits also a natural (free) action of $\mathbb{U}_r \cap \sigma^{-1}(\mathbb{U}_r)$ by right multiplication. If $r = 1$, the quotient of $S_{T,U}$ by this action is ($\overline{\mathbb{F}}_q$ -isomorphic to) a classical Deligne–Lusztig variety for the reductive \mathbb{F}_q -group \mathbb{G}_1 . \diamond

Lemma 4.3. *For (T,U) and (T',U') as in Section 3, the map*

$$\check{G}_r^\sigma \backslash (S_{T,U} \times S_{T',U'}) \xrightarrow{\sim} \Sigma, \quad (g,g') \mapsto (g^{-1}\sigma(g), g'^{-1}\sigma(g'), g^{-1}g'),$$

is a $\check{T}_r^\sigma \times \check{T}'^{\sigma}$ -equivariant isomorphism, where \check{G}_r^σ acts diagonally on $S_{T,U} \times S_{T',U'}$.

By functoriality of cohomology, the $\check{G}_r^\sigma \times \check{T}_r^\sigma$ -action on $S_{T,U}$ induces for each $i \in \mathbb{Z}$ a $\check{G}_r^\sigma \times \check{T}_r^\sigma$ -action on $H_c^i(S_{T,U}, \overline{\mathbb{Q}}_\ell)$. For a character $\theta: \check{T}_r^\sigma \rightarrow \overline{\mathbb{Q}}_\ell^\times$, let $H_c^i(S_{T,U}, \overline{\mathbb{Q}}_\ell)_\theta$ denote the θ -isotypic component. It is stable under the action of \check{G}_r^σ .

Definition 4.4. We define the virtual \check{G}_r^σ -representation with $\overline{\mathbb{Q}}_\ell$ -coefficients

$$R_{\mathbf{x},T,U,r}^\theta := \sum_{i \in \mathbb{Z}} (-1)^i H_c^i(S_{T,U}, \overline{\mathbb{Q}}_\ell)_\theta.$$

By pullback, we can also consider $R_{\mathbf{x},T,U,r}^\theta$ a virtual representation of the parahoric subgroup $\check{P}_{\mathbf{x}}^\sigma$ of $G(k)$. If \mathbf{x} is clear from the context, we write $R_{T,U,r}^\theta$ for $R_{\mathbf{x},T,U,r}^\theta$.

Moreover, by Corollary 4.7 below, $R_{T,U,r}^\theta$ does not depend on the choice of U , if θ is regular. In this case we denote $R_{T,U,r}^\theta$ by $R_{T,r}^\theta$. For the dependence on r see Section 4.2.

Recall the group $N_{\mathbb{G}_r}(\mathbb{T}_r, \mathbb{T}'_r)$ from Section 2.8. The next few results are all corollaries of Theorem 3.2, generalizing [Lus04, Propositions 2.2, 2.3, Corollary 2.4].

Proposition 4.5. *Assume that $r \geq 2$. Let (T,U) , (T',U') be as in Section 3. Further, let $\theta: \check{T}_r^\sigma \rightarrow \overline{\mathbb{Q}}_\ell^\times$, $\theta': \check{T}'^{\sigma} \rightarrow \overline{\mathbb{Q}}_\ell^\times$ be two characters.*

- (i) *Let $i, i' \in \mathbb{Z}$. Assume that an irreducible \check{G}_r^σ -representation appears in the dual space $(H_c^i(S_{T,U}, \overline{\mathbb{Q}}_\ell)_{\theta-1})^\vee$ of $H_c^i(S_{T,U}, \overline{\mathbb{Q}}_\ell)_\theta$ and in $H_c^{i'}(S_{T',U'}, \overline{\mathbb{Q}}_\ell)_{\theta'}$. Then there exists an integer $n \geq 1$ and a $g \in N_{\mathbb{G}_r}(\mathbb{T}'_r, \mathbb{T}_r)(\mathbb{F}_{q^n})$ such that the adjoint action of g carries $\theta \circ N_{\sigma^n}^\theta|_{\mathcal{T}(\overline{\mathbb{F}}_q)^{\sigma^n}}$ to $\theta' \circ N_{\sigma^n}^{\theta'}|_{\mathcal{T}'(\overline{\mathbb{F}}_q)^{\sigma^n}}$.*

- (ii) Assume that an irreducible \check{G}_r^σ -representation occurs in $R_{T,U,r}^\theta$ and $R_{T',U',r}^{\theta'}$. Then there exist some $n \geq 1$ and $g \in N_{\mathbb{G}_r}(\mathbb{T}'_r, \mathbb{T}_r)(\mathbb{F}_{q^n})$ such that the adjoint action of g carries $\theta \circ N_{\check{\sigma}}^{\sigma^n} |_{\mathcal{T}(\mathbb{F}_q)^{\sigma^n}}$ to $\theta' \circ N_{\check{\sigma}}^{\sigma^n} |_{\mathcal{T}'(\mathbb{F}_q)^{\sigma^n}}$.

Proof. The proof (using Lemma 4.3 and Lemma 3.1) is literally the same as the proof of [Lus04, Proposition 2.2]. We omit the details. \square

Proposition 4.6. *Assume that $r \geq 2$. Let (T, U, θ) and (T', U', θ') be two triples as in Proposition 4.5. Assume that θ or θ' is regular. Then*

$$\langle R_{T,U,r}^\theta, R_{T',U',r}^{\theta'} \rangle = \#\{w \in W_{\mathbf{x}}(T, T')^\sigma : \theta \circ \text{Ad}(w) = \theta'\}.$$

Proof. A standard computation using Lemma 4.3 and the Künneth formula shows that $\langle R_{T,U}^\theta, R_{T',U'}^{\theta'} \rangle = \sum_{i \in \mathbb{Z}} \dim H_c^i(\Sigma, \overline{\mathbb{Q}}_\ell)_{\theta^{-1}, \theta'}$. Now apply Theorem 3.2. \square

Corollary 4.7. *Assume that $r \geq 2$. Let $\theta: \check{T}_r^\sigma \rightarrow \overline{\mathbb{Q}}_\ell^\times$ be regular.*

- (i) $R_{T,U,r}^\theta$ is independent of the choice of U .
(ii) If additionally the stabilizer of θ in $W_{\mathbf{x}}(T, T)^\sigma$ is trivial, then $\pm R_{T,U,r}^\theta$ is an irreducible representation of \check{G}_r^σ and of $P_{\mathbf{x}}(\mathcal{O}_k)$.

Proof. See [Lus04, Corollary 2.4]. \square

4.2. Change of level. One could hope that if θ is a character of $T(\mathcal{O}_k) = (\check{T}^0)^\sigma$ which is trivial on $(\check{T}^r)^\sigma$, then the representations $R_{T,U,r}^\theta$ and $R_{T,U,s}^\theta$ for all $s \geq r$ coincide. In [CI18, Proposition 7.6], it is shown that this holds when G is an inner form of $\text{GL}_n(k)$ and T is an elliptic torus. We will show in subsequent work that for general G which split over \check{k} , this is true when T is elliptic. However this fails for general T . In some sense, the more T splits, the bigger is the discrepancy between $R_{T,U,r}^\theta$ and $R_{T,U,r+1}^\theta$. We will explain the failure in an example.

Assume that G is quasi-split over k and let $T \subseteq G$ be a maximal k -rational torus, which contains a k -split maximal torus of G . Under these assumptions there is a k -rational Borel subgroup of G containing T . Let U be its unipotent radical. There is a hyperspecial vertex $\mathbf{x} = \mathbf{x}_0$ contained in $\mathcal{A}_{T,\check{k}} \cap \mathcal{B}_k$. Let $r \geq 1$, and let θ be a character of $(\check{T}^0)^\sigma$, which factors through the character (again denoted θ) of \check{T}_r^σ . For each $s \geq r$,

$$S_{\mathbf{x},T,U,s}/\mathbb{U}_s = (\mathbb{G}_s/\mathbb{U}_s)^\sigma = \mathbb{G}_s^\sigma/\mathbb{U}_s^\sigma$$

is a discrete point set. For a surjection of groups $H \twoheadrightarrow K$, let Inf_K^H denote the inflation functor from virtual K -representations to virtual H -representations given by pullback. Since $S_{\mathbf{x},T,U,s}$ and $S_{\mathbf{x},T,U,r}/\mathbb{U}_r$ have the same cohomology groups up to an even degree shift, we then have

$$\begin{aligned} R_{\mathbf{x},T,U,s}^\theta &= \text{Ind}_{\check{B}_s^\sigma}^{\check{G}_s^\sigma} \text{Inf}_{\check{B}_r^\sigma}^{\check{B}_s^\sigma} \theta, \\ \text{Inf}_{\check{G}_r^\sigma}^{\check{G}_s^\sigma} R_{\mathbf{x},T,U,r}^\theta &= \text{Inf}_{\check{G}_r^\sigma}^{\check{G}_s^\sigma} \text{Ind}_{\check{B}_r^\sigma}^{\check{G}_r^\sigma} \theta = \text{Ind}_{\check{B}_s^\sigma \check{G}_s^{r,\sigma}}^{\check{G}_s^\sigma} \text{Inf}_{\check{B}_r^\sigma}^{\check{B}_s^\sigma \check{G}_s^{r,\sigma}} \theta, \end{aligned}$$

where the last formula follows from a general commutativity fact for inflation and induction ($\text{Ind}_{HN}^G \text{Inf}_{HN/N}^{HN} \chi = \text{Inf}_{G/N}^G \text{Ind}_{HN/N}^{G/N} \chi$ for an abstract group G , a subgroup $H \subseteq G$, a normal subgroup $N \subseteq G$, and a representation χ of HN/N). Thus $R_{\mathbf{x},T,U,s}^\theta$ is bigger than $\text{Inf}_{\check{G}_r^\sigma}^{\check{G}_s^\sigma} R_{\mathbf{x},T,U,r}^\theta$.

5. TRACES OF VERY REGULAR ELEMENTS

Let the notation be as in the beginning of Section 3.

Definition 5.1. We say that $s \in \check{P}_{\mathbf{x}}$ is *unramified very regular* with respect to \mathbf{x} if the following conditions hold:

- (i) s is a regular semisimple element of $G_{\check{k}}$,
- (ii) the connected centralizer $Z^\circ(s)$ of s is a \check{k} -split maximal torus of $G_{\check{k}}$ whose apartment contains \mathbf{x} , and
- (iii) $\alpha(s) \not\equiv 1$ modulo \mathfrak{p} for all roots α of $Z^\circ(s)$ in $G_{\check{k}}$.

For $r \geq 2$, we say that $s \in \mathbb{G}_r$ is *unramified very regular*, if s is the image of an unramified very regular element of $\check{P}_{\mathbf{x}}$.

Note that condition (ii) implies condition (i). Note that in condition (iii) the character $\alpha: Z^\circ(s) \rightarrow \mathbb{G}_{m,\check{k}}$ induces a homomorphism of maximal bounded subgroups: $\alpha: \check{Z}^\circ(s) \rightarrow \mathcal{O}^\times$, and hence the condition makes sense.

Remark 5.2. When G is an inner form of GL_n and T is the maximal nonsplit unramified torus in G , Definition 5.1 says that $x \in (\check{T}^0)^\sigma = \mathcal{O}_L^\times$ (here $\check{k} \supseteq L \supseteq k$ is the degree- n -subextension) is unramified very regular if and only if the image of x in $(\mathcal{O}_L/U_L^1) \cong \mathbb{F}_{q^n}^\times$ has trivial $\mathrm{Gal}(\mathbb{F}_{q^n}/\mathbb{F}_q)$ -stabilizer. This is *not* equivalent to (though is implied by) the condition that the image of x in $\mathbb{F}_{q^n}^\times$ is a generator although this last condition is sometimes also associated to the same terminology [Hen92, BW13, CI18]. \diamond

Note that if $s \in \check{P}_{\mathbf{x}}$ is unramified very regular, then we may consider the $W_{\mathbf{x}}(T)$ -homogeneous space $W_{\mathbf{x}}(T, Z^\circ(s))$ (see Section 2.8).

Theorem 5.3. Let $\theta: \check{T}_r^\sigma \rightarrow \overline{\mathbb{Q}}_\ell^\times$ be any character and let $s \in \check{P}_{\mathbf{x}}^\sigma$ be unramified very regular with respect to \mathbf{x} . Then

$$\mathrm{Tr}(g, R_{T,U,r}^\theta) = \sum_{w \in W_{\mathbf{x}}(T, Z^\circ(g))^\sigma} (\theta \circ \mathrm{Ad}(w^{-1}))(g).$$

Corollary 5.4. Let $T' \subset G$ be a k -rational \check{k} -split maximal torus whose apartment contains \mathbf{x} . If T and T' are not conjugate by an element of $\check{P}_{\mathbf{x}}^\sigma$, then for any $s \in T'(k)$ unramified very regular with respect to \mathbf{x} ,

$$\mathrm{Tr}(s, R_{T,U,r}^\theta) = 0.$$

Proof. We need to show that for two such tori, $W_{\mathbf{x}}(T, T')^\sigma = \emptyset$. Suppose there is an element $w \in W_{\mathbf{x}}(T, T')^\sigma$. Then its preimage in $N_{\mathbb{G}_r}(\mathbb{T}_r, \mathbb{T}'_r)$ form a \mathbb{F}_q -rational \mathbb{T}_r -torsor, which by Lang's theorem has a rational point. Doing this for all r and using that the inverse limit of a family of non-empty compact sets is non-empty, we can find an element $n \in \check{P}_{\mathbf{x}}^\sigma$, which conjugates $T(\mathcal{O})$ into $T'(\mathcal{O})$. The centralizer of $T(\mathcal{O})$ in $G(\check{k})$ is $T(\check{k})$ (and similarly for T'), so n also conjugates $T(\check{k})$ into $T'(\check{k})$, and so it conjugates T into T' , which contradicts the assumption. \square

Let B denote the Borel subgroup of G whose unipotent radical is the fixed subgroup U , and let \mathbb{B}_r be the corresponding subgroup of \mathbb{G}_r . The following result shows that \mathbb{B}_r behaves in certain aspects like a Borel subgroup of \mathbb{G}_r (although it is not a Borel subgroup if $r \geq 2$). Similar results in the case that $P_{\mathbf{x}}$ is reductive are shown in [Sta12].

Proposition 5.5. *Let $s \in \check{G}_r$ be an unramified very regular element. If $x \in \check{G}_r$ is such that $s \in x\check{B}_r x^{-1}$, then there exists a unique $w \in W_{\mathbf{x}}(T, Z^0(s))$ such that for any lift $\dot{w} \in \check{G}_r$, we have $x \in \dot{w}\check{B}_r$.*

Proof. The maximal \check{k} -split tori T and $Z^0(s)$ are conjugate by an element $y \in \check{P}_{\mathbf{x}}$, as \mathbf{x} is contained in the intersection of their apartments. Conjugating by y we thus may reduce to the special case that $Z^0(s) = T$.

We first prove the assertion in the case $r = 1$. The image of s in the reductive group \check{G}_1 is regular semisimple and $\mathbb{B}_1 \subseteq \mathbb{G}_1$ is a Borel subgroup. By [DL76, Proposition 4.4(ii)], we see that there is an element $\dot{w} \in \check{G}_1$ normalizing T , and satisfying $x\mathbb{B}_1 x^{-1} = \dot{w}^{-1}\mathbb{B}_1\dot{w}$. By the normalizer theorem, $\dot{w}^{-1}x \in \check{B}_1$, and we are done.

We now prove the assertion for $r \geq 2$. By the above, we see that there exists a unique $w \in W_{\mathbf{x}}(T)$ such that $x \in \dot{w}\check{B}_r\check{G}_r^1$. We proceed by induction; to this end, it suffices to prove that if $x \in \dot{w}\check{B}_r\check{G}_r^{r-1}$, then $x \in \dot{w}\check{B}_r$.

Since \check{G}_r^{r-1} is normal in \check{G}_r , we may write $x = \dot{w}hb$ for some $h \in \check{G}_r^{r-1}$ and $b \in \check{B}_r$. By [MP96, Theorem 4.2], \check{G}_r^{r-1} has an Iwahori decomposition, so we may write $h = h_-h_+$ with $h_- \in \check{U}_r^{-,r-1}$ and $h_+ \in \check{B}_r^{r-1}$. Replacing b by h_+b and h by h_- , we now have $h \in \check{U}_r^{-,r-1}$. Since $x^{-1}sx \in \check{B}_r$ by assumption, we have $h^{-1}\text{Ad}(w^{-1})(s)h \in \check{B}_r$. Writing t for the very regular element $\text{Ad}(w^{-1})(s) \in \check{T}_r$, we deduce $h^{-1}(tht^{-1})t \in \check{B}_r$, and thus $h^{-1}(tht^{-1}) \in \check{B}_r$. Since $h \in \check{U}_r^{-,r-1}$ by construction, $h^{-1}(tht^{-1}) \in \check{B}_r$ only if $h = tht^{-1}$, which holds only when $h = 1$ by Lemma 5.6. \square

Lemma 5.6. *Let $r \geq 2$ and let $t \in \check{T}_r \subset \check{G}_r$ be unramified very regular. If $tht^{-1} = h$ for some $h \in \check{U}_r$, then $h = 1$.*

Proof. Fixing an order on the roots $\Phi^+ = \Phi(T, U)$, we may write h uniquely as $\prod_{\alpha \in \Phi^+} \psi_{\alpha}(h_{\alpha})$, where ψ_{α} is an isomorphism of $\mathbb{U}_{\alpha, r}$ with a framing object coming from the Chevalley system. Then $\prod_{\alpha \in \Phi^+} \psi_{\alpha}(h_{\alpha}) = h = \zeta^{-1}h\zeta = \prod_{\alpha \in \Phi^+} \psi_{\alpha}(\alpha(\zeta^{-1})h_{\alpha})$, and hence (by uniqueness of the presentation as a product) $h_{\alpha} = \alpha(\zeta^{-1})h_{\alpha}$. We have naturally $h_{\alpha} \in \mathfrak{p}^{r_{1, \alpha}}/\mathfrak{p}^{r_{2, \alpha}}$ for appropriate $r_{1, \alpha} \leq r_{2, \alpha} \in \mathbb{Z}$. As ζ^{-1} is very regular, $\alpha(\zeta^{-1}) \not\equiv 1 \pmod{\mathfrak{p}}$, and hence the above equality forces $h_{\alpha} = 0$ for all α . Thus $h = 1$. \square

By Proposition 5.5,

$$S_{T,U}^{(g, \mathbb{T})} := \{x \in \check{G}_r : x^{-1}\sigma(x) \in \check{U}_r \text{ and } gx \in x\check{T}_r^{\sigma}\} = \bigsqcup_{w \in W_{\mathbf{x}}(T, Z^0(g))} S_{T,U}^{(g, \mathbb{T})}(w), \quad (5.1)$$

where

$$S_{T,U}^{(g, \mathbb{T})}(w) := \{x \in \dot{w}\check{B}_r : x^{-1}\sigma(x) \in \check{U}_r \text{ and } gx \in x\check{T}_r^{\sigma}\},$$

for some (any) lift $\dot{w} \in \mathbb{G}_r$ of w . For any k -rational \check{k} -split maximal torus $T' \subset G$ whose apartment contains \mathbf{x} , the preimage of any $w \in W_{\mathbf{x}}(T, T')^{\sigma}$ in \mathbb{G}_r is an \mathbb{F}_q -rational \mathbb{T}_r -torsor, so by Lang's theorem, it contains a \mathbb{F}_q -rational point \dot{w} . For any $w \in W(T, T')^{\sigma}$ we fix such a \dot{w} .

Proposition 5.7. *Let $g \in \check{G}_r^{\sigma}$ be a very regular element. Then*

$$S_{T,U}^{(g, \mathbb{T})}(w) = \begin{cases} \dot{w}\check{T}_r^{\sigma} & \text{if } w \in W_{\mathbf{x}}(T, Z^0(g))^{\sigma}, \\ \emptyset & \text{otherwise.} \end{cases}$$

Proof. Let $w \in W_{\mathbf{x}}(T, Z^0(g))$ and let \dot{w} be any lift of w to \check{G}_r . Assume that $S_{T,U}^{(g, \mathbb{T})}(w) \neq \emptyset$ and let $x \in S_{T,U}^{(g, \mathbb{T})}(w)$. Then $\dot{w}^{-1}x \in \check{B}_r$. After modifying the lift \dot{w} , we may write $x = \dot{w}v$ with

$v \in \check{U}_r$. As $(\check{w}v)^{-1}\sigma(\check{w}v) = x^{-1}\sigma(x) \in \check{U}_r$, we deduce that $\check{w}^{-1}\sigma(\check{w}) \in \check{U}_r$. Projecting to \mathbb{G}_1 , we see that $\bar{w}^{-1}\sigma(\bar{w}) \in \check{U}_1$. The left hand side is a semisimple element in \mathbb{G}_1 , and so we deduce that $w \in W_{\mathbf{x}}(T)^\sigma$.

We may now choose \check{w} to be the σ -stable lift \dot{w} . Then we can write $x = \dot{w}av$ where $a \in \check{T}_r$ and $v \in \check{U}_r$. Then $x^{-1}\sigma(x) = v^{-1}a^{-1}\dot{w}^{-1}\sigma(\dot{w})\sigma(a)\sigma(v) = v^{-1}a^{-1}\sigma(a)\sigma(v) \in \check{U}_r$ and hence we must have $a \in \check{T}_r^\sigma$. Furthermore, by assumption we have $x^{-1}gx = v^{-1}a^{-1}\dot{w}^{-1}g\dot{w}av = v_1^{-1}\dot{w}^{-1}g\dot{w}v_1 \in \check{T}_r^\sigma$, where $v_1 := av_1a^{-1} \in \check{U}_r$. The element $t := \dot{w}^{-1}g\dot{w} \in \check{T}_r^\sigma$ is very regular and now $v^{-1}(tv_1^{-1}) \in \check{T}_r^\sigma$. Hence necessarily $v = tv_1^{-1}$, which forces $v = 1$ by Lemma 5.6. \square

Proof of Theorem 5.3. For any \check{k} -split maximal torus $T' \subset G$, we have a short exact sequence

$$1 \rightarrow (\check{T}_r'^1)^\sigma \rightarrow \check{T}_r'^\sigma \rightarrow \check{T}_1'^\sigma \rightarrow 1$$

of finite abelian groups with $(\check{T}_r'^1)^\sigma$ of p -power order and $\check{T}_1'^\sigma$ of order prime to p . (The surjectivity on the right holds as $\check{T}_1'^\sigma \rightarrow H^1(\text{Gal}(\overline{\mathbb{F}}_q/\mathbb{F}_q), \check{T}_r'^1)$ must be the zero morphism, as the latter is a p -group). This sequence is split.

Applying the above to $T' = Z^0(g)$, we may write $g = st_1$ where $t_1 \in (\check{T}_r'^1)^\sigma$ has p -power order and s is in the image of the splitting and hence of order prime to p . It is easy to see that t_1 and s are both powers of g . Note that s is still very regular and $Z^0(s) = Z^0(g)$. Analogously, applying the above to $T' = T$, for any $\tau \in \check{T}_r^\sigma$, we may write $\tau = \zeta\tau_1$ with $\tau_1 \in (\check{T}_r^1)^\sigma$, and ζ in the image of the splitting. Thus $(g, \tau) \in \check{G}_r^\sigma \times \check{T}_r^\sigma$ has the decomposition

$$(g, \tau) = (s, \zeta) \cdot (t_1, \tau_1),$$

where (s, ζ) and (t_1, τ_1) are both powers of (g, τ) such that (s, ζ) has prime-to- p order and (t_1, τ_1) has p -power order. Averaging over \check{T}_r^σ and applying the Deligne–Lusztig trace formula [DL76, Theorem 3.2] (which we may do by Lemma 4.1), we deduce

$$\begin{aligned} \text{Tr}(g, R_{T,U,r}^\theta) &= \frac{1}{\#\check{T}_r^\sigma} \sum_{\tau \in \check{T}_r^\sigma} \theta(\tau)^{-1} \text{Tr} \left((g, \tau)^*; \sum_i (-1)^i H_c^i(S_{T,U}, \overline{\mathbb{Q}}_\ell) \right) \\ &= \frac{1}{\#\check{T}_r^\sigma} \sum_{\tau \in \check{T}_r^\sigma} \theta(\tau)^{-1} \text{Tr} \left((t_1, \tau_1)^*; \sum_i (-1)^i H_c^i(S_{T,U}^{(s,\zeta)}, \overline{\mathbb{Q}}_\ell) \right), \end{aligned} \quad (5.2)$$

where $S_{T,U}^{(s,\zeta)} := \{x \in \mathbb{G}_r : x^{-1}\sigma(x) \in \mathbb{U}_r, sx\zeta = x\}$ is the set of fixed points of $S_{T,U}$ under (s, ζ) .

We obviously have $S_{T,U}^{(s,\zeta)} \subseteq S_{T,U}^{(g,\mathbb{T})}$, and it now follows easily from Proposition 5.7 that

$$S_{T,U}^{(s,\zeta)} = \begin{cases} \dot{w}\mathbb{T}_r^\sigma & \text{if } \zeta = \text{Ad}(w^{-1})(s^{-1}) \text{ for some (unique) } w \in W_{\mathbf{x}}(T, Z^0(g))^\sigma, \\ \emptyset & \text{otherwise.} \end{cases}$$

Now (t_1, τ_1) acts on a point $\dot{w}a \in \dot{w}\mathbb{T}_r^\sigma$ by $(t_1, \tau_1): \dot{w}a \mapsto t_1\dot{w}a\tau_1 = \dot{w}\text{Ad}(w^{-1})(t_1)a\tau_1$, and thus

$$\begin{aligned} \text{Tr} \left((t_1, \tau_1)^*; \sum_i (-1)^i H_c^i(S_{T,U}^{(s,\zeta)}, \overline{\mathbb{Q}}_\ell) \right) &= \text{Tr} \left((t_1, \tau_1)^*; H_c^0(\dot{w}\mathbb{T}_r^\sigma) \right) \\ &= \begin{cases} \#\check{T}_r^\sigma & \text{if } \tau_1 = \text{Ad}(w^{-1})(t_1^{-1}), \\ 0 & \text{otherwise,} \end{cases} \end{aligned}$$

and Theorem 5.3 now follows from (5.2). \square

REFERENCES

- [BH05] C.J. Bushnell and G. Henniart. The essentially tame local Langlands correspondence, II: Totally ramified representations. *Compos. Math.*, 141(4):979–1011, 2005.
- [Bor91] A. Borel. *Linear algebraic groups*, volume 126 of *Graduate Texts in Mathematics*. Springer-Verlag, 1991.
- [BS17] B. Bhatt and P. Scholze. Projectivity of the Witt vector affine Grassmannian. *Invent. Math.*, 209(2):329–423, August 2017.
- [BT72] F. Bruhat and J. Tits. Groupes réductifs sur un corps local. *Inst. Hautes Études Sci. Publ. Math.*, 41:5–251, 1972.
- [BT84] F. Bruhat and J. Tits. Groupes réductifs sur un corps local II. Schémas en groupes. Existence d’une donnée radicielle valuée. *Inst. Hautes Études Sci. Publ. Math.*, 60:197–376, 1984.
- [BW13] M. Boyarchenko and J. Weinstein. Geometric realization of special cases of local Langlands and Jacquet-Langlands correspondences. *Preprint*, 2013. arXiv:1303.5795.
- [BW16] M. Boyarchenko and J. Weinstein. Maximal varieties and the local Langlands correspondence for $GL(n)$. *J. Amer. Math. Soc.*, 29:177–236, 2016.
- [CI18] C. Chan and A. Ivanov. Affine Deligne–Lusztig varieties at infinite level. 2018. arxiv:1811.11204.
- [DL76] P. Deligne and G. Lusztig. Representations of reductive groups over finite fields. *Ann. of Math.*, 103(1):103–161, 1976.
- [Gre61] M. Greenberg. Schemata over local rings. *Ann. Math.*, 73(3):624–648, 1961.
- [Gre63] M. Greenberg. Schemata over local rings. *Ann. Math.*, 78(2):256–266, 1963.
- [Hen92] G. Henniart. Correspondance de Langlands–Kazhdan explicite dans le cas non ramifié. *Math. Nachr.*, 158(7-26), 1992.
- [HR08] T. Haines and M. Rapoport. On parahoric subgroups. *Appendix to [PR08]*, 2008.
- [Lus79] G. Lusztig. Some remarks on the supercuspidal representations of p -adic semisimple groups. In *Automorphic forms, representations and L-functions, Proc. Symp. Pure Math. 33 Part 1 (Corvallis, Ore., 1977)*, pages 171–175, 1979.
- [Lus04] G. Lusztig. Representations of reductive groups over finite rings. *Represent. Theory*, 8:1–14, 2004.
- [MP94] A. Moy and G. Prasad. Unrefined minimal K-types for p -adic groups. *Invent. Math.*, 116:393–408, 1994.
- [MP96] A. Moy and G. Prasad. Jacquet functors and unrefined minimal k-types. *Comment. Math. Helv.*, 71(1):98–121, 1996.
- [PR08] G. Pappas and M. Rapoport. Twisted loop groups and their affine flag varieties. *Adv. in Math.*, 219:118–198, 2008.
- [Sta09] A. Stasinski. Unramified representations of reductive groups over finite rings. *Represent. Theory*, 13:636–656, 2009.
- [Sta12] A. Stasinski. Reductive group schemes, the greenberg functor, and associated algebraic groups. *J. Pure and Appl. Alg.*, 216(4):1092–1101, 2012.
- [Tit79] J. Tits. Reductive groups over local fields. *Proc. Symp. Pure Math.*, 33:29–69, 1979.
- [Yu01] J.-K. Yu. Construction of tame supercuspidal representations. *J. Amer. Math. Soc.*, 14:579–622, 2001.
- [Yu02] J.-K. Yu. Smooth models associated to concave functions in Bruhat-Tits theory. *Preprint*, 2002.
- [Zhu17] X. Zhu. Affine Grassmannians and the geometric Satake in mixed characteristic. *Ann. of Math.*, 185(2):403–492, 2017. arXiv:1407.8519.

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, FINE HALL, WASHINGTON ROAD, PRINCETON, NJ 08544-1000 USA

Email address: `charchan@math.princeton.edu`

MATHEMATISCHES INSTITUT, UNIVERSITÄT BONN, ENDENICHER ALLEE 60, 53115 BONN, GERMANY

Email address: `ivanov@math.uni-bonn.de`