

Polynomial bounds for chromatic number.
IV. A near-polynomial bound for excluding the five-vertex path

Alex Scott¹

Mathematical Institute, University of Oxford, Oxford OX2 6GG, UK

Paul Seymour²

Princeton University, Princeton, NJ 08544

Sophie Spirkl³

University of Waterloo, Waterloo, Ontario N2L3G1, Canada

August 8, 2021; revised June 26, 2024

¹Research supported by EPSRC grant EP/V007327/1.

²Supported by AFOSR grant FA9550-22-1-0234, and by NSF grant DMS-2154169.

³We acknowledge the support of the Natural Sciences and Engineering Research Council of Canada (NSERC) [funding reference number RGPIN-2020-03912]. Cette recherche a été financée par le Conseil de recherches en sciences naturelles et en génie du Canada (CRSNG) [numéro de référence RGPIN-2020-03912].

Abstract

A graph G is H -free if it has no induced subgraph isomorphic to H . We prove that a P_5 -free graph with clique number $\omega \geq 3$ has chromatic number at most $\omega^{\log_2(\omega)}$. The best previous result was an exponential upper bound $(5/27)3^\omega$, due to Esperet, Lemoine, Maffray, and Morel. A polynomial bound would imply that the celebrated Erdős-Hajnal conjecture holds for P_5 , which is the smallest open case. Thus, there is great interest in whether there is a polynomial bound for P_5 -free graphs, and our result is an attempt to approach that.

1 Introduction

If G, H are graphs, we say G is H -free if no induced subgraph of G is isomorphic to H ; and for a graph G , we denote the number of vertices, the chromatic number, the size of the largest clique, and the size of the largest stable set by $|G|, \chi(G), \omega(G), \alpha(G)$ respectively.

The k -vertex path is denoted by P_k , and P_4 -free graphs are well-understood; every P_4 -free graph G with more than one vertex is either disconnected or disconnected in the complement [24], which implies that $\chi(G) = \omega(G)$. Here we study how $\chi(G)$ depends on $\omega(G)$ for P_5 -free graphs G .

The Gyárfás-Sumner conjecture [10, 25] says:

1.1 Conjecture: *For every forest H there is a function f such that $\chi(G) \leq f(\omega(G))$ for every H -free graph G .*

This is open in general, but has been proved [10] when H is a path, and for several other simple types of tree ([3, 11, 12, 13, 14, 17, 19]; see [18] for a survey). The result is also known if all induced subdivisions of a tree are excluded [17].

A class of graphs is *hereditary* if the class is closed under taking induced subgraphs and under isomorphism, and a hereditary class is said to be χ -bounded if there is a function f such that $\chi(G) \leq f(\omega(G))$ for every graph G in the class (thus, the Gyárfás-Sumner conjecture says that, for every forest H , the class of H -free graphs is χ -bounded). Louis Esperet [8] made the following conjecture:

1.2 (False) Conjecture: *Let \mathcal{G} be a χ -bounded class. Then there is a polynomial function f such that $\chi(G) \leq f(\omega(G))$ for every $G \in \mathcal{G}$.*

Esperet's conjecture was recently shown to be false by Brianiński, Davies and Walczak [2]. However, this raises the further question: which χ -bounded classes are polynomially χ -bounded? In particular, the two conjectures 1.1 and 1.2 would together imply the following, which is still open:

1.3 Conjecture: *For every forest H , there exists $c > 0$ such that $\chi(G) \leq \omega(G)^c$ for every H -free graph G .*

This is a beautiful conjecture. In most cases where the Gyárfás-Sumner conjecture has been proved, the current bounds are very far from polynomial, and 1.3 has been only been proved for a much smaller collection of forests (see [15, 20, 22, 23, 21, 5, 16]). In [23] we proved it for any P_5 -free tree H , but it has not been settled for any tree H that contains P_5 . In this paper we focus on the case $H = P_5$.

The best previously-known bound on the chromatic number of P_5 -free graphs in terms of their clique number, due to Esperet, Lemoine, Maffray, and Morel [9], was exponential:

1.4 *If G is P_5 -free and $\omega(G) \geq 3$ then $\chi(G) \leq (5/27)3^{\omega(G)}$.*

Here we make a significant improvement, showing a “near-polynomial” bound:

1.5 *If G is P_5 -free and $\omega(G) \geq 3$ then $\chi(G) \leq \omega(G)^{\log_2(\omega(G))}$.*

(The cycle of length five shows that we need to assume $\omega(G) \geq 3$. Sumner [25] showed that $\chi(G) \leq 3$ when $\omega(G) = 2$.) Conjecture 1.3 when $H = P_5$ is of great interest, because of a famous conjecture due to Erdős and Hajnal [6, 7], that:

1.6 Conjecture: For every graph H there exists $c > 0$ such that $\alpha(G)\omega(G) \geq |G|^c$ for every H -free graph G .

This is open in general, despite a great deal of effort; and in view of [4], the smallest graph H for which 1.6 is undecided is the graph P_5 . Every forest H satisfying 1.3 also satisfies the Erdős-Hajnal conjecture, and so showing that $H = P_5$ satisfies 1.3 would be a significant result. (See [1] for some other recent progress on this question.)

We use standard notation throughout. When $X \subseteq V(G)$, $G[X]$ denotes the subgraph induced on X . We write $\chi(X)$ for $\chi(G[X])$ when there is no ambiguity.

2 The main proof

We denote the set of nonnegative real numbers by \mathbb{R}_+ , and the set of nonnegative integers by \mathbb{Z}_+ . Let $f : \mathbb{Z}_+ \rightarrow \mathbb{R}_+$ be a function. We say

- f is *non-decreasing* if $f(y) \geq f(x)$ for all integers $x, y \geq 0$ with $y > x \geq 0$;
- f is a *binding* function for a graph G if it is non-decreasing and $\chi(H) \leq f(\omega(H))$ for every induced subgraph H of G ; and
- f is a *near-binding* function for G if f is non-decreasing and $\chi(H) \leq f(\omega(H))$ for every induced subgraph H of G different from G .

In this section we show that if a function f satisfies a certain inequality, then it is a binding function for all P_5 -free graphs. Then at the end we will give a function that satisfies the inequality, and deduce 1.5.

A *cutset* in a graph G is a set X such that $G \setminus X$ is disconnected. A vertex $v \in V(G)$ is *mixed* on a set $A \subseteq V(G)$ or a subgraph A of a graph G if v is not in A and has a neighbour and a non-neighbour in A . It is *complete* to A if it is adjacent to every vertex of A . We begin with the following:

2.1 Let G be P_5 -free, and let f be a near-binding function for G . Let G be connected, and let X be a cutset of G . Then

$$\chi(G \setminus X) \leq f(\omega(G) - 1) + \omega(G)f(\lfloor \omega(G)/2 \rfloor).$$

Proof. We may assume (by replacing X by a subset if necessary) that X is a minimal cutset of G ; and so $G \setminus X$ has at least two components, and every vertex in X has a neighbour in $V(B)$, for every component B of $G \setminus X$. Let B be one such component; we will prove that $\chi(B) \leq f(\omega(G) - 1) + \omega(G)f(\lfloor \omega(G)/2 \rfloor)$, from which the result follows.

Choose $v \in X$ (this is possible since G is connected), and let N be the set of vertices in B adjacent to v . Let the components of $B \setminus N$ be $R_1, \dots, R_k, S_1, \dots, S_\ell$, where R_1, \dots, R_k each have chromatic number more than $f(\lfloor \omega(G)/2 \rfloor)$, and S_1, \dots, S_ℓ each have chromatic number at most $f(\lfloor \omega(G)/2 \rfloor)$. Let S be the union of the graphs S_1, \dots, S_ℓ ; thus, $\chi(S) \leq f(\lfloor \omega(G)/2 \rfloor)$. For $1 \leq i \leq k$, let Y_i be the set of vertices in N with a neighbour in $V(R_i)$, and let $Y = Y_1 \cup \dots \cup Y_k$.

- (1) For $1 \leq i \leq k$, every vertex in Y_i is complete to R_i .

Let $y \in Y_i$. Thus, y has a neighbour in $V(R_i)$; suppose that y is mixed on R_i . Since R_i is connected, there is an edge ab of R_i such that y is adjacent to a and not to b . Now v has a neighbour in each component of $G \setminus X$, and since there are at least two such components, there is a vertex $u \in V(G) \setminus (X \cup V(B))$ adjacent to v . But then $u-v-y-a-b$ is an induced copy of P_5 , a contradiction. This proves (1).

$$(2) \chi(Y) \leq (\omega(G) - 1)f(\lfloor \omega(G)/2 \rfloor).$$

Let $1 \leq i \leq k$. Since $f(\lfloor \omega(G)/2 \rfloor) < \chi(R_i) \leq f(\omega(R_i))$, and f is non-decreasing, it follows that $\omega(R_i) > \omega(G)/2$. By (1), $\omega(G[Y_i]) + \omega(R_i) \leq \omega(G)$, and so $\omega(G[Y_i]) < \omega(G)/2$. Consequently $\chi(Y_i) \leq f(\lfloor \omega(G)/2 \rfloor)$, for $1 \leq i \leq k$. Choose $I \subseteq \{1, \dots, k\}$ minimal such that $\bigcup_{i \in I} Y_i = Y$. From the minimality of I , for each $i \in I$ there exists $y_i \in Y_i$ such that for each $j \in I \setminus \{i\}$ we have that $y_i \notin Y_j$; and so the vertices y_i ($i \in I$) are all distinct. For each $i \in I$ choose $r_i \in V(R_i)$. For all distinct $i, j \in I$, if y_i, y_j are nonadjacent, then $r_i-y_i-v-y_j-r_j$ is isomorphic to P_5 , a contradiction. Hence the vertices y_i ($i \in I$) are all pairwise adjacent, and adjacent to v ; and so $|I| \leq \omega(G) - 1$. Thus, $\chi(Y) = \chi(\bigcup_{i \in I} Y_i) \leq (\omega(G) - 1)f(\lfloor \omega(G)/2 \rfloor)$. This proves (2).

All the vertices in $N \setminus Y$ are adjacent to v , and so $\omega(G[N \setminus Y]) \leq \omega(G) - 1$. Moreover, for $1 \leq i \leq k$, each vertex of R_i is adjacent to each vertex in Y_i , and $Y_i \neq \emptyset$ since B is connected, and so $\omega(R_i) \leq \omega(G) - 1$. Since there are no edges between any two of the graphs $G[N \setminus Y], R_1, \dots, R_k$, their union (Z say) has clique number at most $\omega(G) - 1$ and so has chromatic number at most $f(\omega(G) - 1)$. But $V(B)$ is the union of $Y, V(S)$ and $V(Z)$; and so

$$\chi(B) \leq f(\omega(G) - 1) + (\omega(G) - 1)f(\lfloor \omega(G)/2 \rfloor) + f(\lfloor \omega(G)/2 \rfloor).$$

This proves 2.1. ■

2.2 Let $\Omega \geq 1$, and let $f : \mathbb{Z}_+ \rightarrow \mathbb{R}_+$ be non-decreasing, satisfying the following:

- f is a binding function for every P_5 -free graph H with $\omega(H) \leq \Omega$; and
- $f(w - 1) + (w + 2)f(\lfloor w/2 \rfloor) \leq f(w)$ for each integer $w > \Omega$.

Then f is a binding function for every P_5 -free graph G .

Proof. We prove by induction on $|G|$ that if G is P_5 -free then f is a binding function for G . Thus, we may assume that G is P_5 -free and f is near-binding for G . If G is not connected, or $\omega(G) \leq \Omega$, it follows that f is binding for G , so we assume that G is connected and $\omega(G) > \Omega$. Let us write $w = \omega(G)$ and $m = \lfloor w/2 \rfloor$. If $\chi(G) \leq f(w)$ then f is a binding function for G , so we assume, for a contradiction, that:

$$(1) \chi(G) > f(w - 1) + (w + 2)f(m).$$

We deduce that:

$$(2) \text{ Every cutset } X \text{ of } G \text{ satisfies } \chi(X) > 2f(m).$$

If some cutset X satisfies $\chi(X) \leq 2f(m)$, then since $\chi(G \setminus X) \leq f(w-1) + wf(m)$ by 2.1, it follows that $\chi(G) \leq f(w-1) + (w+2)f(m)$, contrary to (1). This proves (2).

(3) *If P, Q are cliques of G , both of cardinality at least $w/2$, then $G[P \cup Q]$ is connected.*

Suppose not; then there is a minimal subset $X \subseteq V(G) \setminus (P \cup Q)$ such that P, Q are subsets of different components (A, B say) of $G \setminus X$. From the minimality of X , every vertex $x \in X$ has a neighbour in $V(A)$ and a neighbour in $V(B)$. If x is mixed on A and mixed on B , then since A is connected, there is an edge a_1a_2 of A such that x is adjacent to a_1 and not to a_2 ; and similarly there is an edge b_1b_2 of B with x adjacent to b_1 and not to b_2 . But then $a_2-a_1-x-b_1-b_2$ is an induced copy of P_5 , a contradiction; so every $x \in X$ is complete to at least one of A, B . The set of vertices in X complete to A is also complete to P , and hence has clique number at most m , and hence has chromatic number at most $f(m)$; and the same for B . Thus, $\chi(X) \leq 2f(m)$, contrary to (2). This proves (3).

If $v \in V(G)$, we denote its set of neighbours by $N(v)$, or $N_G(v)$. Let $a \in V(G)$, and let B be a component of $G \setminus (N(a) \cup \{a\})$; we will show that $\chi(B) \leq (w-m+2)f(m)$.

A subset Y of $V(B)$ is a *joint* of B if there is a component C of $B \setminus Y$ such that $\chi(C) > f(m)$ and Y is complete to C . If \emptyset is not a joint of B then $\chi(B) < f(m)$ and the claim holds, so we may assume that \emptyset is a joint of B ; let Y be a joint of B chosen with Y maximal, and let C be a component of $B \setminus Y$ such that $\chi(C) > f(m)$ and Y is complete to C .

(4) *If $v \in N(a)$ has a neighbour in $V(C)$, then $\chi(V(C) \setminus N(v)) \leq f(m)$.*

Let $N_C(v)$ be the set of neighbours of v in $V(C)$, and $M = V(C) \setminus N_C(v)$; and suppose that $\chi(M) > f(m)$. Let C' be a component of $G[M]$ with $\chi(C') > f(m)$, and let Z be the set of vertices in $N_C(v)$ that have a neighbour in $V(C')$. Thus, $Z \neq \emptyset$, since $N_C(v), V(C') \neq \emptyset$ and C is connected. If some $z \in Z$ is mixed on C' , let p_1p_2 be an edge of C' such that z is adjacent to p_1 and not to p_2 ; then $a-v-z-p_1-p_2$ is an induced copy of P_5 , a contradiction. So every vertex in Z is complete to $V(C')$; but also every vertex in Y is complete to $V(C)$ and hence to $V(C')$, and so $Y \cup Z$ is a joint of B , contrary to the maximality of Y . This proves (4).

(5) $\chi(Y) \leq f(m)$ and $\chi(C) \leq (w-m+1)f(m)$.

Let X be the set of vertices in $N(a)$ that have a neighbour in $V(C)$. Since C is a component of $B \setminus Y$ and hence a component of $G \setminus (X \cup Y)$, and a belongs to a different component of $G \setminus (X \cup Y)$, it follows that $X \cup Y$ is a cutset of G . By (2), $\chi(X \cup Y) > 2f(m)$. Since $\omega(C) \geq m+1$ (because $\chi(C) > f(m)$, and f is near-binding for G) and every vertex in Y is complete to $V(C)$, it follows that $\omega(G[Y]) \leq w-m-1 \leq m$, and so has chromatic number at most $f(m)$ as claimed; and so $\chi(X) > f(m)$. Consequently there is a clique $P \subseteq X$ with cardinality $w-m$. The subgraph induced on the set of vertices of C complete to P has clique number at most m , and so has chromatic number at most $f(m)$; and for each $v \in P$, the set of vertices of C nonadjacent to v has chromatic number at most $f(m)$ by (4). Thus, $\chi(C) \leq (|P|+1)f(m) = (w-m+1)f(m)$. This proves (5).

$$(6) \chi(B) \leq (w - m + 2)f(m).$$

By (3), every clique contained in $V(B) \setminus (V(C) \cup Y)$ has cardinality less than $w/2$ (because it is anticomplete to the largest clique of C) and so

$$\chi(B \setminus (V(C) \cup Y)) \leq f(m);$$

and hence $\chi(B \setminus Y) \leq (w - m + 1)f(m)$ by (5), since there are no edges between C and $V(B) \setminus (V(C) \cup Y)$. But $\chi(Y) \leq f(m)$ by (5), and so $\chi(B) \leq (w - m + 2)f(m)$. This proves (6).

By (6), $G \setminus N(a)$ has chromatic number at most $(w - m + 2)f(m)$. But $G[N(a)]$ has clique number at most $w - 1$ and so chromatic number at most $f(w - 1)$; and so $\chi(G) \leq f(w - 1) + (w - m + 2)f(m)$, contrary to (1). This proves 2.2. \blacksquare

Now we deduce 1.5, which we restate:

2.3 *If G is P_5 -free and $\omega(G) \geq 3$ then $\chi(G) \leq \omega(G)^{\log_2(\omega(G))}$.*

Proof. Define $f(0) = 0$, $f(1) = 1$, $f(2) = 3$, and $f(x) = x^{\log_2(x)}$ for every real number $x \geq 3$. Let G be P_5 -free. If $\omega(G) \leq 2$ then $\chi(G) \leq 3 = f(2)$, by a result of Sumner [25]; if $\omega(G) = 3$ then $\chi(G) \leq 5 \leq f(3)$, by an application of the result 1.4 of Esperet, Lemoine, Maffray, and Morel [9]; and if $\omega(G) = 4$ then $\chi(G) \leq 15 \leq f(4)$, by another application of 1.4. Consequently every P_5 -free graph G with clique number at most four has chromatic number at most $f(\omega(G))$.

We claim that

$$f(x - 1) + (x + 2)f(\lfloor x/2 \rfloor) \leq f(x)$$

for each integer $x > 4$. If that is true, then by 2.2 with $\Omega = 4$, we deduce that $\chi(G) \leq f(\omega(G))$ for every P_5 -free graph G , and so 1.5 holds. Thus, it remains to show that

$$f(x - 1) + (x + 2)f(\lfloor x/2 \rfloor) \leq f(x)$$

for each integer $x > 4$. This can be verified by direct calculation when $x = 5$, so we may assume that $x \geq 6$.

The derivative of $f(x)/x^4$ is

$$(2\log_2(x) - 4)x^{\log_2(x)-5},$$

and so is nonnegative for $x \geq 4$. Consequently

$$\frac{f(x - 1)}{(x - 1)^4} \leq \frac{f(x)}{x^4}$$

for $x \geq 5$. Since $x^2(x^2 - 2x - 4) \geq (x - 1)^4$ when $x \geq 5$, it follows that

$$\frac{f(x - 1)}{x^2 - 2x - 4} \leq \frac{f(x)}{x^2},$$

that is,

$$f(x - 1) + \frac{2x + 4}{x^2}f(x) \leq f(x),$$

when $x \geq 5$. But when $x \geq 6$ (so that $f(x/2)$ is defined and the first equality below holds), we have

$$f(\lfloor x/2 \rfloor) \leq f(x/2) = (x/2)^{\log_2(x/2)} = (x/2)^{\log_2(x)-1} = (2/x)(x/2)^{\log_2(x)} = (2/x^2)f(x),$$

and so

$$f(x-1) + (x+2)f(\lfloor x/2 \rfloor) \leq f(x)$$

when $x \geq 6$. This proves 2.3. ■

References

- [1] P. Blanco and M. Bucić, “Towards the Erdős-Hajnal conjecture for P_5 -free graphs”, manuscript, September 2022.
- [2] M. Briański, J. Davies and B. Walczak, “Separating polynomial χ -boundedness from χ -boundedness”, [arXiv:2201.08814](https://arxiv.org/abs/2201.08814).
- [3] M. Chudnovsky, A. Scott and P. Seymour, “Induced subgraphs of graphs with large chromatic number. XII. Distant stars”, *J. Graph Theory* **92** (2019), 237–254, [arXiv:1711.08612](https://arxiv.org/abs/1711.08612).
- [4] M. Chudnovsky, A. Scott, P. Seymour and S. Spirkl, “Erdős-Hajnal for graphs with no five-hole”, submitted for publication, [arXiv:2102.04994](https://arxiv.org/abs/2102.04994).
- [5] M. Chudnovsky, A. Scott, P. Seymour and S. Spirkl, “Polynomial bounds for chromatic number. VI. Adding a four-vertex path”, submitted for publication, [arXiv:2202.10412](https://arxiv.org/abs/2202.10412).
- [6] P. Erdős and A. Hajnal, “On spanned subgraphs of graphs”, *Graphentheorie und Ihre Anwendungen* (Oberhof, 1977).
- [7] P. Erdős and A. Hajnal, “Ramsey-type theorems”, *Discrete Applied Math.* **25** (1989), 37–52.
- [8] L. Esperet, *Graph Colorings, Flows and Perfect Matchings*, Habilitation thesis, Université Grenoble Alpes (2017), 24, <https://tel.archives-ouvertes.fr/tel-01850463/document>.
- [9] L. Esperet, L. Lemoine, F. Maffray, and G. Morel, “The chromatic number of $\{P_5, K_4\}$ -free graphs”, *Discrete Math.* **313** (2013), 743–754.
- [10] A. Gyárfás, “On Ramsey covering-numbers”, in *Infinite and Finite Sets, Vol. II* (Colloq., Keszthely, 1973), *Coll. Math. Soc. János Bolyai* **10**, 801–816.
- [11] A. Gyárfás, “Problems from the world surrounding perfect graphs”, *Proceedings of the International Conference on Combinatorial Analysis and its Applications*, (Pokrzywna, 1985), *Zastos. Mat.* **19** (1987), 413–441.
- [12] A. Gyárfás, E. Szemerédi and Zs. Tuza, “Induced subtrees in graphs of large chromatic number”, *Discrete Math.* **30** (1980), 235–344.
- [13] H. A. Kierstead and S. G. Penrice, “Radius two trees specify χ -bounded classes”, *J. Graph Theory* **18** (1994), 119–129.

- [14] H. A. Kierstead and Y. Zhu, “Radius three trees in graphs with large chromatic number”, *SIAM J. Disc. Math.* **17** (2004), 571–581.
- [15] X. Liu, J. Schroeder, Z. Wang and X. Yu, “Polynomial χ -binding functions for t -broom-free graphs”, [arXiv:2106.08871](#).
- [16] I. Schiermeyer and B. Randerath, “Polynomial χ -binding functions and forbidden induced subgraphs: a survey”, *Graphs and Combinatorics* **35** (2019), 1–31.
- [17] A. Scott, “Induced trees in graphs of large chromatic number”, *J. Graph Theory* **24** (1997), 297–311.
- [18] A. Scott and P. Seymour, “A survey of χ -boundedness”, *J. Graph Theory* **95** (2020), 473–504, [arXiv:1812.07500](#).
- [19] A. Scott and P. Seymour, “Induced subgraphs of graphs with large chromatic number. XIII. New brooms”, *European J. Combinatorics* **84** (2020), 103024, [arXiv:1807.03768](#).
- [20] A. Scott, P. Seymour and S. Spirkl, “Polynomial bounds on chromatic number. I. Excluding a biclique and an induced tree”, *J. Graph Theory*, to appear, [arXiv:2104.07927](#).
- [21] A. Scott and P. Seymour, “Polynomial bounds for chromatic number. V. Excluding a tree of radius two and a complete multipartite graph”, submitted for publication, [arXiv:arXiv:2202.05557](#).
- [22] A. Scott, P. Seymour and S. Spirkl, “Polynomial bounds on chromatic number. II. Excluding a star-forest”, *J. Graph Theory*, **101** (2022), 318–322, [arXiv:2107.11780](#).
- [23] A. Scott, P. Seymour and S. Spirkl, “Polynomial bounds on chromatic number. III. Excluding a double star”, *J. Graph Theory*, **101** (2022), 323–340, [arXiv:2108.07066](#).
- [24] D. Seinsche, “On a property of the class of n -colorable graphs”, *J. Combin. Theory, Ser. B*, **16** (1974), 191–193.
- [25] D. P. Sumner, “Subtrees of a graph and chromatic number”, in *The Theory and Applications of Graphs*, (G. Chartrand, ed.), John Wiley & Sons, New York (1981), 557–576.