

An incompressible 2D didactic model with singularity and explicit solutions of the 2D Boussinesq equations

Dongho Chae, Peter Constantin and Jiahong Wu

Abstract. We give an example of a well posed, finite energy, 2D incompressible active scalar equation with the same scaling as the surface quasi-geostrophic equation and prove that it can produce finite time singularities. In spite of its simplicity, this seems to be the first such example. Further, we construct explicit solutions of the 2D Boussinesq equations whose gradients grow exponentially in time for all time. In addition, we introduce a variant of the 2D Boussinesq equations which is perhaps a more faithful companion of the 3D axisymmetric Euler equations than the usual 2D Boussinesq equations.

Mathematics Subject Classification (2010). Primary 35Q35; Secondary 76D03.

Keywords. Inviscid model, singularity, explicit solutions, 2D Boussinesq equations.

1. Introduction

The purpose of this paper is threefold: first, to provide an example of an incompressible 2D active scalar, similar to the inviscid surface quasi-geostrophic (SQG) equation, that possesses a family of solutions which develop finite-time singularities; second, to construct a class of explicit solutions to the inviscid 2D Boussinesq equations that grow exponentially in time; and third, to propose for study a modified 2D Boussinesq system, that appears to provide a closer comparison to the 3D axisymmetric Euler equations than the standard 2D Boussinesq equations. All the calculations are elementary, and the results serve as didactic examples.

The inviscid SQG equation is

$$\begin{cases} \partial_t \theta + u \cdot \nabla \theta = 0, & x \in \mathbb{R}^2, t > 0, \\ u = \nabla^\perp \psi \equiv (-\partial_{x_2}, \partial_{x_1}) \psi, & \Lambda \psi = \theta, \end{cases} \quad (1.1)$$

where θ and ψ are scalar functions of x and t , u denotes the 2D velocity field and $\Lambda = (-\Delta)^{\frac{1}{2}}$ denotes the Zygmund operator, which can be defined through the Fourier transform

$$\widehat{\Lambda f}(\xi) = |\xi| \widehat{f}(\xi).$$

The inviscid SQG equation is useful in modeling atmospheric phenomena such as frontogenesis, the formation of strong fronts between masses of hot and cold air ([4, 6, 7, 11]). In addition, the inviscid SQG equation is a significant example of a 2D active scalar and some of its distinctive features have made it an important testbed for turbulence theories ([1]). Mathematically, the inviscid SQG equation is difficult to analyze, and the issue of whether its solutions can develop singularities in a finite time remains open. Other active scalars with the same scaling $u \sim \theta$ include the porous medium (or Muskat) equation where $u = (0, \theta) + \nabla p$. The blow up problem from smooth initial data is open there as well (see [2] and references therein). Here we give a first example of a 2D incompressible active scalar equation with a velocity field having the same level of regularity and scaling as the active scalar, $u \sim \theta$,

$$\begin{cases} \partial_t \theta + u \cdot \nabla \theta = 0, & x \in \mathbb{R}^2, t > 0, \\ u = \nabla^\perp \psi, & -\partial_{x_2} \psi = \theta. \end{cases} \quad (1.2)$$

The only difference between (1.1) and (1.2) is the equation relating ψ and θ . It is not very difficult to see that (1.2) is locally well-posed in a sufficiently regular functional setting. What is more striking about this model is that (1.2) admits a family of solutions which develop finite-time singularities even though they are initially smooth and have finite energy.

Very recently Luo and Hou performed careful numerical simulations of the 3D axisymmetric incompressible Euler equations which suggested the appearance of a finite-time singularity [9]. We briefly describe the setup and their main result. The spatial domain is the cylinder

$$\left\{ (x_1, x_2, z) : r \equiv \sqrt{x_1^2 + x_2^2} \leq 1, 0 \leq z \leq L \right\}$$

with periodic boundary conditions in the z -direction and no-penetration condition on the solid boundary $r = 1$. The angular components of the velocity, vorticity and stream functions are odd with respect to z across $z = 0$. The velocity on the unit circle $z = 0$, $r = 1$ vanishes, and thus all points on this circle are stagnation points. According to [9], the vorticity of a numerical solution at the stagnation points blows up in finite time. This finite-time singularity does not appear to be well understood theoretically. Motivated by the numerical simulations of Luo and Hou, Kiselev and Sverak recently proved the double exponential growth (in time) of the vorticity gradient of 2D Euler solutions in the unit disk [8]. The odd symmetry, the stagnation point and the boundary appear to be important in their work. Other pursuits for lower bounds for the vorticity gradient of the 2D Euler equation can be found in [5, 12]. Our goal here is to provide explicit solutions to the 2D incompressible Boussinesq equations which exhibit exponential growth in time.

The 2D Boussinesq equations are

$$\begin{cases} \partial_t \omega + u \cdot \nabla \omega = \partial_{x_1} \theta, & x \in \mathbb{R}^2, t > 0, \\ \partial_t \theta + u \cdot \nabla \theta = 0, \\ u = \nabla^\perp \psi, \quad \Delta \psi = \omega. \end{cases} \quad (1.3)$$

As pointed out in [10], (1.3) is closely related to the 3D axisymmetric Euler equations. We construct two families of solutions of (1.3). The first family is given in a fixed domain with a stagnation point at the corner of the domain. The solutions in the family have an odd symmetry with respect to $x_2 = 0$, i.e. they are odd as functions of x_2 , in the direction of gravity, and have $\nabla \theta$ growing exponentially in time. The second family of solutions of (1.3) consists of smooth global solutions in a domain with a moving boundary. Both $\nabla \omega$ and $\nabla \theta$ grow exponentially in time. It may be possible to further exploit and extend these constructions to obtain solutions with gradients with double exponential growth.

We also propose for study the following modified 2D Boussinesq system

$$\begin{cases} \partial_t \omega + u \cdot \nabla \omega = -\partial_{x_2}(\theta^2), & x \in \mathbb{R}^2, t > 0, \\ \partial_t \theta + u \cdot \nabla \theta = 0, \\ u = \nabla^\perp \psi, \quad \Delta \psi = \omega. \end{cases} \quad (1.4)$$

(1.4) differs from (1.3) on the right-hand side of the vorticity equation, and has a different parity symmetry. Note that the gravity now points in the x_1 direction, which might be confusing but it is done to mimic the set-up of Luo and Hou, but the main difference with usual Boussinesq is that both θ and ω are allowed to be odd in x_2 , i.e. in the direction perpendicular to gravity, not parallel to it. As explained in Section 4, (1.4) appears to be a more exact match with the 3D axisymmetric Euler equations as set up in the numerical simulations of Luo and Hou [9]. We construct a class of solutions of (1.4) whose gradients exhibit exponential growth.

The rest of the paper is divided into three sections. The second section details the active scalar model similar to the SQG equation and describes its solutions which develop finite-time singularities. The third section constructs two families of solutions to the 2D Boussinesq equations while the last section presents a variant of the 2D Boussinesq equations and some of its explicit solutions with exponential gradient growth.

2. A 2D model with finite-time singularity

This section presents an active scalar equation transported by an divergence-free velocity field that admits solutions with finite-time singularities.

Consider the initial-value problem (1.2), namely

$$\begin{cases} \partial_t \theta + u \cdot \nabla \theta = 0, & x \in \mathbb{R}^2, t > 0, \\ u = \nabla^\perp \psi, & -\partial_{x_2} \psi = \theta, \\ \psi(x, 0) = \psi_0(x). \end{cases} \quad (2.1)$$

It is not difficult to see that (2.1) is locally well-posed if $\psi_0 \in H^s(\mathbb{R}^2)$ with $s > 3$ and $\theta_0 = -\partial_{x_2} \psi_0 \in L^1(\mathbb{R}^2)$. Now, we assume that the initial stream function ψ_0 , the initial active scalar θ_0 and the initial-velocity u_0 obey the following symmetry, for any $(x_1, x_2) \in \mathbb{R}^2$,

$$\psi_0(x_1, -x_2) = -\psi_0(x_1, x_2), \quad \theta_0(x_1, -x_2) = \theta_0(x_1, x_2),$$

$$u_{01}(x_1, -x_2) = u_{01}(x_1, x_2), \quad u_{02}(x_1, -x_2) = -u_{02}(x_1, x_2).$$

It is easy to check that the corresponding solution of (2.1) preserves this symmetry. In particular, $u_2(x_1, 0, t) = 0$ and the equation for θ on the x_1 -axis becomes

$$\partial_t \theta(x_1, 0, t) + u_1(x_1, 0, t) \partial_{x_1} \theta(x_1, 0, t) = 0.$$

or

$$\partial_t \theta(x_1, 0, t) + \theta(x_1, 0, t) \partial_{x_1} \theta(x_1, 0, t) = 0, \quad (2.2)$$

which is the inviscid Burgers equation. It is well-known that Burgers' equation can develop discontinuities. In fact, if the initial data satisfies

$$\partial_{x_1} \theta_0(x_1, 0) < 0 \quad \text{for some } x_1 \in \mathbb{R},$$

then the corresponding solution $\theta(x_1, 0, t)$ of (2.2) becomes singular in a finite time, namely $\partial_{x_1} \theta$ becomes infinity in a finite time. One could also restrict to a periodic domain and obtain a finite time blow up. For example,

$$\psi_0(x_1, x_2) = -\cos(x_1) \sin(x_2)$$

will generate a solution that becomes singular in a finite time.

3. Explicit solutions of the 2D Boussinesq equations

In this section we construct two families of solutions to the 2D Boussinesq equations given by (1.3). The first family of solutions is defined in a wedge and the gradient of θ grows exponentially in time. The second family of solutions is close to the first family, but it is defined in a smooth domain with a moving boundary, and the gradients of both ω and θ grow exponentially in time.

3.1. Explicit solutions in a wedge

The spatial domain is bounded by two half-lines in the positive half-plane:

$$x_2 = 2x_1, \quad x_2 = -2x_1, \quad x_1 \geq 0.$$

The stream function, the velocity field and the vorticity are given by

$$\begin{aligned}\psi(x) &= \begin{cases} \frac{x_2^2}{2} - x_1x_2, & x_2 \geq 0, \\ -\frac{x_2^2}{2} - x_1x_2, & x_2 < 0; \end{cases} \\ u_1(x) &= -\partial_{x_2}\psi = \begin{cases} -x_2 + x_1, & x_2 \geq 0, \\ x_2 + x_1, & x_2 < 0; \end{cases} \\ u_2(x) &= \partial_{x_1}\psi = -x_2; \\ \omega(x) &= \Delta\psi = \begin{cases} 1, & x_2 \geq 0, \\ -1, & x_2 < 0; \end{cases}\end{aligned}$$

It is clear that $\psi = 0$ on the boundary of the domain and, consequently, u satisfies the no-penetration boundary condition. Given any initial data $\theta_0(x)$ depending only on x_2 and odd with respect to x_1 -axis, namely

$$\theta_0(x) = \theta_0(x_2), \quad \theta_0(-x_2) = -\theta_0(x_2), \quad x_2 \in \mathbb{R},$$

the corresponding solution $\theta(x, t)$ preserves these properties and satisfies

$$\begin{cases} \partial_t\theta(x_2, t) - x_2\partial_{x_2}\theta(x_2, t) = 0, \\ \theta(x, 0) = \theta_0(x_2). \end{cases} \quad (3.1)$$

Integrating on characteristics, θ is given by

$$\theta(x_2, t) = \theta_0(e^t x_2).$$

with arbitrary odd θ_0 . It is easy to see that the gradient of any solution to (3.1) grows exponentially in time. In fact, $\partial_{x_2}\theta$ satisfies

$$\partial_t(\partial_{x_2}\theta) - \partial_{x_2}\theta - x_2\partial_{x_2}(\partial_{x_2}\theta) = 0,$$

which is solved by

$$\partial_{x_2}\theta = e^t (\theta'_0)(e^t x_2).$$

A special example is given by

$$\theta(x, t) = \sin(x_2 e^t), \quad \partial_{x_2}\theta(x, t) = e^t \cos(x_2 e^t).$$

Many other explicit solutions can be obtained by taking different initial data.

3.2. Explicit smooth solutions in a moving domain

The spatial domain in the previous subsection has a sharp corner at the origin and the vorticity has jumps when crossing the x_1 -axis. In this subsection we construct a smooth solution.

For $\sigma(x_2, t)$ to be explicitly determined later, we seek solutions with stream function, velocity field and vorticity given by

$$\psi(x) = \begin{cases} \frac{1}{2}\sigma(x_2, t)x_2^2 - x_1x_2, & x_2 \geq 0, \\ -\frac{1}{2}\sigma(x_2, t)x_2^2 - x_1x_2, & x_2 < 0; \end{cases}$$

$$\begin{aligned}
u_1(x) = -\partial_{x_2}\psi &= \begin{cases} -\sigma(x_2, t)x_2 - \frac{1}{2}x_2^2\partial_{x_2}\sigma + x_1, & x_2 \geq 0, \\ \sigma(x_2, t)x_2 + \frac{1}{2}x_2^2\partial_{x_2}\sigma + x_1, & x_2 < 0; \end{cases} \\
u_2(x) = \partial_{x_1}\psi &= -x_2; \\
\omega(x) = \Delta\psi &= \begin{cases} \frac{1}{2}\partial_{x_2}^2(\sigma(x_2, t)x_2^2), & x_2 \geq 0, \\ -\frac{1}{2}\partial_{x_2}^2(\sigma(x_2, t)x_2^2), & x_2 < 0. \end{cases}
\end{aligned}$$

Consequently, the boundary of the domain is given by $\psi = 0$, namely

$$2x_1 = \sigma(x_2, t)x_2, \quad 2x_1 = -\sigma(x_2, t)x_2.$$

Since the vorticity depends only on x_2 , the vorticity equation is reduced to

$$\partial_t\omega - x_2\partial_{x_2}\omega = 0, \quad \omega(x_2, 0) = \omega_0(x_2),$$

which is solved by

$$\omega(x, t) = \omega_0(e^t x_2).$$

Therefore, $\sigma(x_2, t)$ must satisfy

$$\partial_{x_2}^2(\sigma(x_2, t)x_2^2) = 2\omega_0(e^t x_2), \text{ for } x_2 > 0 \text{ and } -\partial_{x_2}^2(\sigma(x_2, t)x_2^2) = 2\omega_0(e^t x_2), \text{ for } x_2 < 0.$$

We also seek a solution θ depending on x_2 and t only, namely $\theta = \theta(x_2, t)$, then

$$\partial_t\theta - x_2\partial_{x_2}\theta = 0, \quad \theta(x_2, 0) = \theta_0(x_2),$$

which is solved by

$$\theta(x, t) = \theta_0(e^t x_2).$$

Some special examples of the solutions include

$$\psi(x, t) = \frac{1}{6}x_2^3 e^t - x_1 x_2, \quad \omega = x_2 e^t, \quad \theta = x_2 e^t$$

in the domain bounded by $x_1 = \frac{1}{6}x_2^2 e^t$.

4. A variant of the 2D Boussinesq equations

In this section we suggest a new 2D Boussinesq system for study. This new system modifies the standard 2D Boussinesq equations and appears to be a more direct parallel to the 3D axisymmetric Euler equations. A class of explicit solutions are constructed to exemplify the behavior of it solutions.

The modified 2D Boussinesq equations are given in (1.4). We explain why this modified system is identical to the 3D axisymmetric Euler equations, away from the axis of symmetry. We first recall the 3D axisymmetric Euler equations in cylindrical coordinates (see, e.g., [3, 10]). Let \mathbf{e}_r , \mathbf{e}_θ and \mathbf{e}_z be the orthonormal unit vectors defining the cylindrical coordinate system,

$$\mathbf{e}_r = \left(\frac{x_1}{r}, \frac{x_2}{r}, 0 \right), \quad \mathbf{e}_\theta = \left(-\frac{x_2}{r}, \frac{x_1}{r}, 0 \right), \quad \mathbf{e}_z = (0, 0, 1),$$

where $r = \sqrt{x_1^2 + x_2^2}$. A vector field v is axisymmetric if

$$v = v^r(r, z, t)\mathbf{e}_r + v^\theta(r, z, t)\mathbf{e}_\theta + v^z(r, z, t)\mathbf{e}_z.$$

The 3D axisymmetric Euler equations can be written as

$$\begin{cases} \frac{\tilde{D}}{Dt}(ru^\theta) = 0, \\ \frac{\tilde{D}}{Dt}\left(\frac{\omega^\theta}{r}\right) = -\frac{1}{r^4}\partial_z(ru^\theta)^2, \\ (\partial_r^2 + \frac{1}{r}\partial_r + \partial_z^2 - \frac{1}{r^2})\psi^\theta = \omega^\theta, \end{cases} \quad (4.1)$$

where u^θ , ω^θ and ψ^θ are the angular components of the velocity, vorticity and stream function, respectively, and

$$\frac{\tilde{D}}{Dt} = \partial_t + u^r\partial_r + u^z\partial_z.$$

(4.1) is a complete system since u^r and u^z can be recovered from ψ^θ by

$$u^r = -\partial_z\psi^\theta, \quad u^z = \frac{1}{r}\partial_r(r\psi^\theta).$$

Therefore the divergence-free condition

$$\partial_r(ru^r) + \partial_z(ru^z) = 0$$

is automatically satisfied. We remark that the notation in the book of Majda and Bertozzi [10, p.62-66] is slightly different from that in Luo and Hou [9]. In particular, ψ in [10] corresponds to $r\psi^\theta$ in [9] and ω^θ in [10] corresponds to $-\omega^\theta$ in [9].

[9] numerically solved (4.1) in the cylinder

$$D(1, L) = \{(r, z) : 0 \leq r \leq 1, 0 \leq z \leq L\}$$

with the initial data

$$u_0^\theta(r, z) = 100r e^{-30(1-r^2)^4} \sin\left(\frac{2\pi}{L}z\right), \quad \omega_0^\theta(r, z) = \psi_0^\theta(r, z) = 0$$

subject to the periodic boundary condition in z ,

$$u^\theta(r, 0, t) = u^\theta(r, L, t), \quad \omega^\theta(r, 0, t) = \omega^\theta(r, L, t), \quad \psi^\theta(r, 0, t) = \psi^\theta(r, L, t)$$

and the no-flow boundary condition on the solid boundary $r = 1$,

$$\psi^\theta(1, z, t) = 0.$$

Since the initial data $(u_0^\theta, \omega_0^\theta, \psi_0^\theta)$ is odd with respect to $z = 0$, the solution $(u^\theta, \omega^\theta, \psi^\theta)$ is also odd with respect to $z = 0$. Consequently,

$$u^r(1, z, t) = 0, \quad u^z(r, 0, t) = 0.$$

In particular, at $z = 0$ and $r = 1$,

$$u = (u^r, u^\theta, u^z) = 0$$

and all points on this circle are stagnation points.

The modified 2D Boussinesq system that we propose for study is given by (1.4), namely

$$\begin{cases} \partial_t \omega + u \cdot \nabla \omega = -\partial_{x_2}(\rho^2), & x \in \mathbb{R}^2, t > 0, \\ \partial_t \rho + u \cdot \nabla \rho = 0, \\ u = \nabla^\perp \psi, \quad \Delta \psi = \omega. \end{cases} \quad (4.2)$$

We changed the name of the scalar θ to ρ in order to avoid confusing it with the notation for the angular variable. We have the following correspondence between (4.1) and (4.2):

$$r \leftrightarrow x_1, \quad z \leftrightarrow x_2, \quad ru^\theta \leftrightarrow \rho, \quad \frac{\omega^\theta}{r} \leftrightarrow \omega, \quad (u^r, u^z) \leftrightarrow u, \quad \psi^\theta \leftrightarrow \psi.$$

Thus, away from the symmetry axis, the behavior of the solutions to these two systems are expected to be identical. In addition, both ω and ρ can have the same parity symmetry with respect to $x_2 = 0$ (they are both odd). Note that the direction x_2 is not the direction of gravity, but it is perpendicular on it.

We seek a family of solutions to (4.2) independent of x_1 ,

$$\omega = \omega(x_2, t), \quad u_2 = u_2(x_2, t), \quad \rho = \rho(x_2, t).$$

Then, from $\partial_1 u_2 - \partial_2 u_1 = \omega$ we have $-\partial_2 u_1 = \omega(x_2, t)$, which yields

$$u_1 = -\int_0^{x_2} \omega(s, t) ds + \phi(x_1, t)$$

for some function $\phi(\cdot)$. By the divergence free condition we have

$$\partial_1 u_1 = \phi'(x_1, t) = -\partial_2 u_2 = -u_2'(x_2, t),$$

from which we obtain

$$u_2(x_2, t) = -\phi'(x_1, t)x_2 + \psi(x_1, t), \quad \forall x_1, x_2 \in \mathbb{R}$$

for a function $\psi(\cdot)$. This provides us with $\phi'(x_1, t) = C_1(t)$, $\psi(x_1, t) = C_2(t)$, and

$$u_2(x_2, t) = C_1(t)x_2 + C_2(t).$$

The divergence free condition again provides us with

$$u_1(x, t) = -C_1(t)x_1 + f(x_2, t)$$

for some $f(\cdot, \cdot)$. To make a comparison with the setup in [9], we further assume the odd symmetry with respect to $x_2 = 0$. To simplify the calculation, we also assume that u_2 is time independent. Then we have further reduction; $C_1 = \text{constant}$, $C_2 = 0$, and thus.

$$u_2 = -x_2, \quad u_1 = x_1 + f(x_2, t).$$

The equations for ω and ρ are now reduced to

$$\begin{cases} \partial_t \omega - x_2 \partial_{x_2} \omega = -\partial_{x_2}(\rho^2), \\ \partial_t \rho - x_2 \partial_{x_2} \rho = 0. \end{cases}$$

From this system we find that the solution ρ is given by

$$\rho(x_2, t) = \rho_0(e^t x_2),$$

and ω by

$$\omega(x_2, t) = \omega_0(e^t x_2) - 2(e^t - 1)\rho_0(e^t x_2)\rho_0'(e^t x_2)$$

with ρ_0, ω_0 arbitrary functions of one variable. If we choose, in particular, $\rho_0(x_2) = x_2$ (fixed), and $\omega_0(x_2) = x_2^{101}$, $\sinh(x_2)$, $\sinh(\sinh x_2)$, etc, we find that arbitrary growth (algebraic, exponential, double exponential, etc) in ω_0 in space engenders arbitrary growth in time for $|\partial_2 \omega(x_2, t)|$ (exponential, double exponential, triple exponential, etc). Other special example is as follows:

$$\psi = \begin{cases} (1 - \frac{2}{3}e^t(e^t - 1)x_2) \frac{x_2^2}{2} - x_1 x_2, & x_2 \geq 0, \\ -(1 + \frac{2}{3}e^t(e^t - 1)x_2) \frac{x_2^2}{2} - x_1 x_2, & x_2 < 0; \end{cases}$$

$$u_1 = -\partial_{x_2} \psi = \begin{cases} -x_2 + e^t(e^t - 1)x_2^2 + x_1, & x_2 \geq 0, \\ x_2 + e^t(e^t - 1)x_2^2 + x_1, & x_2 < 0; \end{cases}$$

$$u_2 = \partial_{x_1} \psi = -x_2$$

and

$$\omega = \Delta \psi = \begin{cases} 1 - 2x_2 e^t(e^t - 1), & x_2 \geq 0, \\ -1 - 2x_2 e^t(e^t - 1), & x_2 < 0; \end{cases}$$

$$\rho = x_2 e^t.$$

One may also construct examples with fast oscillations such as

$$\omega = \begin{cases} 1 - (e^t - 1) \sin(2x_2 e^t), & x_2 \geq 0, \\ -1 - (e^t - 1) \sin(2x_2 e^t), & x_2 < 0; \end{cases}$$

$$\rho = \sin(2x_2 e^t).$$

Acknowledgements

Chae was partially supported by NRF Grant no. 2006-0093854 and NRF no.2009-0083521. Constantin was partially supported by NSF grants DMS-1209394 and DMS-1265132. Wu was partially supported by NSF grant DMS-1209153. Wu thanks Prof. C. Li, Prof. S. Preston and Dr. A. Sarria for discussions. PC and JW thank the hospitality of the Department of Mathematics of Chung-Ang University, where this work was performed.

References

- [1] W. Blumen, Uniform potential vorticity flow, Part I. Theory of wave interactions and two-dimensional turbulence, *J. Atmos. Sci.* **35** (1978), 774-783.
- [2] A. Castro, D. Córdoba, C. Fefferman and F. Gancedo, Breakdown of smoothness for the Muskat problem, *Arch. Ration. Mech. Anal.* **208** (2013), 805-909.
- [3] D. Chae, On the blow-up problem for the axisymmetric 3D Euler equations, *Nonlinearity* **21** (2008), 2053-2060.
- [4] P. Constantin, A. Majda, and E. Tabak, Formation of strong fronts in the 2-D quasi-geostrophic thermal active scalar, *Nonlinearity* **7** (1994), 1495-1533.
- [5] S. Denisov, Double-exponential growth of the vorticity gradient for the two-dimensional Euler equation, *Proc. Amer. Math. Soc.*, to appear.
- [6] A.E. Gill, *Atmosphere-Ocean Dynamics*, Academic Press, New York, 1982.
- [7] I. Held, R. Pierrehumbert, S. Garner, and K. Swanson, Surface quasi-geostrophic dynamics, *J. Fluid Mech.* **282** (1995), 1-20.
- [8] A. Kiselev and V. Sverak, Small scale creation for solutions of the incompressible two dimensional Euler equations, arXiv: 1310.4799v2 [math.AP] 24 Oct 2013.
- [9] G. Luo and T. Hou, Potentially singular solutions of the 3D incompressible Euler equations, arXiv: 1310.0497v1 [physics.flu-dyn] 1 Oct 2013.
- [10] A.J. Majda and A.L. Bertozzi, *Vorticity and Incompressible Flow*, Cambridge University Press, 2001.
- [11] J. Pedlosky, *Geophysical Fluid Dynamics*, Springer, New York, 1987.
- [12] A. Zlatos, Exponential growth of the vorticity gradient for the Euler equation on the torus, arXiv: 1310.6128v1 [math.AP] 23 Oct 2013.

Dongho Chae

Department of Mathematics, College of Natural Science, Chung-Ang University,
Seoul 156-756, South Korea
e-mail: dchae@cau.ac.kr

Peter Constantin

Program in Applied and Computational Mathematics, Princeton University, Fine
Hall, Washington Road, Princeton, NJ 08544-1000, USA.
e-mail: const@math.princeton.edu

Jiahong Wu

Department of Mathematics, Oklahoma State University, 401 Mathematical Sci-
ences, Stillwater, OK 74078, USA; and Department of Mathematics, Chung-Ang
University, Seoul 156-756, Republic of Korea
e-mail: jiahong@math.okstate.edu