

TORSION IN p -ADIC ÉTALE COHOMOLOGY: REMARKS AND CONJECTURES

GUIDO BOSCO

ABSTRACT. Let C be a complete algebraically closed extension of \mathbb{Q}_p , and let \mathfrak{X} be a smooth formal scheme over \mathcal{O}_C . By the work of Bhatt–Morrow–Scholze, it is known that when \mathfrak{X} is proper, the length of the torsion in the integral p -adic étale cohomology of the generic fiber \mathfrak{X}_C is bounded above by the length of the torsion in the crystalline cohomology of its special fiber. In this note, we focus on the non-proper case and observe that when \mathfrak{X} is affine, the torsion in the integral p -adic étale cohomology of \mathfrak{X}_C can even be expressed as a functor of the special fiber, unlike in the proper case. As a consequence, we show that, surprisingly, if \mathfrak{X} is affine, the integral p -adic étale cohomology groups of \mathfrak{X}_C have finite torsion subgroups. We discuss further applications and propose conjectures predicting the torsion in the integral p -adic étale cohomology of a broader class of rigid-analytic varieties over C .

CONTENTS

1.	Introduction	1
2.	Rigidity of torsion	4
3.	Preliminaries on log de Rham–Witt cohomology	6
4.	Finiteness of torsion and applications	8
	References	13

1. INTRODUCTION

Let p be a prime number. We denote by C an algebraically closed complete non-archimedean extension of \mathbb{Q}_p and we write \mathcal{O}_C for its ring of integers.

In recent years, there has been increasing interest in the study of the integral/rational p -adic (pro-)étale cohomology of rigid-analytic varieties over C that are not necessarily proper, such as local Shimura varieties. See e.g. the works [CDN20a], [CDN20b], [CDN21], [Bos21], [Bos23]. One difficulty in the latter works is that, in general, the rational p -adic étale cohomology groups of non-proper rigid-analytic varieties over C are not finite-dimensional \mathbb{Q}_p -vector spaces. Consequently, it becomes important to understand whether such cohomology groups have a reasonable topological structure, or, at the integral level, to understand the torsion in the integral p -adic étale cohomology of such spaces.

Let us begin with the building blocks of rigid-analytic varieties, namely affinoid spaces. While writing [Bos23], and in particular thinking about [Bos23, Conjecture 1.15] (cf. [CDN22, Remarque 0.3]), we arrived from several directions at the following conjecture.

Date: May 15, 2025.

Conjecture 1.1. *Let X be a smooth affinoid rigid space over C .*

- (i) *For all $i \in \mathbb{Z}$, the torsion subgroup of the pro-étale cohomology group $H_{\text{proét}}^i(X, \mathbb{Z}_p)$ is finite.*
- (ii) *For all $n \in \mathbb{Z}_{\geq 0}$, the p^n -torsion subgroup of the Picard group $\text{Pic}(X)$ is finite.*

We note that Conjecture 1.1(i) implies in particular that, for X a smooth affinoid rigid space over C , the cohomology group $H_{\text{proét}}^i(X, \mathbb{Q}_p)$ has a natural structure of \mathbb{Q}_p -Banach space (see Proposition 4.9), therefore it is not pathological as one might think.

The first goal of this note is to prove Conjecture 1.1 in the good reduction case. More precisely:

Theorem 1.2 (Theorem 4.1, Corollary 4.7). *Conjecture 1.1 holds true in the case X has an affine smooth p -adic formal model over \mathcal{O}_C .*

We prove Theorem 1.2 by reducing it to a problem in characteristic p . For the first part of the statement, i.e. in order to prove Conjecture 1.1(i) in the good reduction case, we will rely on the following result expressing the p^n -torsion in the integral p -adic étale cohomology in terms of the special fiber.

We denote by k the residue field of C .

Theorem 1.3 (Theorem 2.1, Theorem 4.14). *Let \mathfrak{X} be a smooth p -adic formal scheme over \mathcal{O}_C that is affine, or, more generally, can be written as a countable increasing union of affine opens. Then, for all $i \in \mathbb{Z}$ and $n \in \mathbb{Z}_{\geq 0}$ we have a natural isomorphism*

$$H_{\text{proét}}^{i+1}(\mathfrak{X}_C, \mathbb{Z}_p)[p^n] \cong H^1(\mathfrak{X}_k, W\Omega_{\log}^i)[p^n].$$

Here, we denote by $W\Omega_{\log}^i$ the logarithmic de Rham–Witt sheaf (see Notation 2.4).

Remark 1.4 (Comparison to the proper case). We note that Theorem 1.3 cannot hold for any smooth p -adic formal scheme \mathfrak{X} over \mathcal{O}_C . In fact, in the case \mathfrak{X} is proper, Bhatt–Morrow–Scholze proved that (cf. [BMS18, Theorem 1.1, (ii)]), for all $i, n \in \mathbb{Z}_{\geq 0}$, we have the inequality

$$\text{length}_{\mathbb{Z}_p}(H_{\text{proét}}^i(\mathfrak{X}_C, \mathbb{Z}_p)[p^n]) \leq \text{length}_{W(k)}(H_{\text{crys}}^i(\mathfrak{X}_k/W(k))[p^n]),$$

but, the torsion in $H_{\text{proét}}^i(\mathfrak{X}_C, \mathbb{Z}_p)$ is not a functor of the special fiber, [BMS18, Remark 2.4]. Therefore, Theorem 1.3 is somewhat orthogonal to the cited results in the proper case.

The proof of Theorem 1.3 in the affine case relies on the work [BMS19], in particular on the comparison between p -adic étale and syntomic cohomology, and crucially uses a rigidity result for the syntomic cohomology proved by Antieau–Mathew–Morrow–Nikolaus, [AMMN22]. Using Theorem 1.3 in the affine case, we prove Theorem 1.2 by employing results on the structure of logarithmic de Rham–Witt cohomology due to Illusie–Raynaud, [IR83]. On the other hand, the extension of Theorem 1.3 beyond the affine case relies on the same finiteness results shown in the proof of Theorem 1.2.

We expect that one can prove Conjecture 1.1 in the semistable reduction case using a logarithmic variant of the results in this paper.

Remark 1.5 (On Drinfeld’s upper half-spaces). In light of the above, it is natural to conjecture, in particular, an extension of Theorem 1.3 to the semistable reduction case; see Conjecture 4.16 for a precise statement. More generally, one could ask for a version of Theorem 1.3 for \mathfrak{X} a semistable p -adic formal scheme over \mathcal{O}_C whose generic fiber \mathfrak{X}_C can be written as a countable increasing union

of open affinoids. The appeal of a positive answer to the latter question lies in the fact that it could be applied, for example, to Stein rigid spaces over C admitting a semistable formal model over \mathcal{O}_C , such as the d -dimensional Drinfeld's upper half-space \mathbb{H}_C^d over C , for $d \in \mathbb{Z}_{\geq 1}$. In this example, this would lead to a conceptual proof of a result by Colmez–Dospinescu–Nizioł, [CDN21], which states that

$$H_{\text{proét}}^i(\mathbb{H}_C^d, \mathbb{Z}_p) \text{ is torsion-free for all } i \geq 0. \quad (1.1)$$

In fact, denoting by \mathfrak{H} the standard semistable formal model over \mathcal{O}_C of \mathbb{H}_C^d , [GK05, §6.1], [CDN20b, §5.1],¹ by a result of Grösse–Klönne (see [CDN20b, Corollary 6.25(2)]), we have

$$H^j(\mathfrak{H}_k/k^0, W\Omega_{\log}^i) = 0 \text{ for all } j > 0 \text{ and } i \geq 0$$

(see Notation 4.15 for the log structures we are considering here). In [CDN21], the result (1.1) is deduced from an explicit computation of $H_{\text{proét}}^i(\mathbb{H}_C^d, \mathbb{Z}_p)$, [CDN21, Theorem 1.1]; on the other hand, as explained in the introduction of *op. cit.*, using (1.1), this computation can be immediately deduced from its rational variant, previously treated in [CDN20b, Theorem 6.55].

We conclude recalling that, according to a folklore conjecture, any smooth affinoid rigid space over C should have a semistable p -adic formal model over \mathcal{O}_C . However, it would be interesting to find a proof of Conjecture 1.1 working purely in terms of rigid-analytic varieties. We hope that, in the future, tools from integral p -adic Hodge theory of rigid-analytic varieties will become available to help address these questions.

Acknowledgments. The main ideas behind this note emerged during conversations with Matthew Morrow in the fall of 2021. However, I was only able to complete the proof of Theorem 1.2 recently, after coming across Illusie–Raynaud's result [IR83, II, Corollaire 3.11]. I warmly thank Matthew for his help and encouragement. I am grateful to Alexander Petrov for many helpful comments and for suggesting Lemma 4.3, which led to a simplification of my previous argument. I also thank Gabriel Dospinescu, Akhil Mathew, and Peter Scholze for comments or corrections on a previous version of this note and for subsequent discussions. Additionally, I would like to thank Marco D'Addezio, Ian Gleason, and Wiesława Nizioł for helpful conversations. Last but not least, I am grateful to the anonymous referee for their invaluable comments, which improved both the content and the exposition of this note.

Notation and conventions 1.6. Let p be a prime number. Unless otherwise stated, we denote by C an algebraically closed complete non-archimedean extension of \mathbb{Q}_p , and we write \mathcal{O}_C for its ring of integers.

All formal schemes are p -adic and locally of finite type over the base. Given a formal scheme \mathfrak{X} over \mathcal{O}_C , we denote by \mathfrak{X}_C its generic fiber regarded as an adic space. In general, all rigid-analytic varieties will be regarded as adic spaces.

We fix an uncountable cardinal κ as in [Sch21, Lemma 4.1]. Given an adic space X over \mathbb{Q}_p , we denote by $X_{\text{proét}}$ its κ -small pro-étale site. We adopt similar notation and conventions for the κ -small pro-étale site of a (formal) scheme.

For an abelian group A and $m \in \mathbb{Z}_{\geq 1}$, we denote by $A[m] \subseteq A$ the kernel of the multiplication by m on A .

¹More precisely, \mathfrak{H} is the base change to \mathcal{O}_C of the semistable formal model over \mathbb{Z}_p of the d -dimensional Drinfeld's upper half-space over \mathbb{Q}_p considered in *loc. cit.*

2. RIGIDITY OF TORSION

Our starting point is the following result.

Theorem 2.1. *Let \mathfrak{X} be an affine smooth formal scheme over \mathcal{O}_C . Let k denote the residue field of C . Then, for all $i \in \mathbb{Z}$ and $n \in \mathbb{Z}_{\geq 0}$ we have a natural isomorphism*

$$H_{\text{proét}}^{i+1}(\mathfrak{X}_C, \mathbb{Z}_p)[p^n] \cong H^1(\mathfrak{X}_k, W\Omega_{\log}^i)[p^n].$$

Here, we denote by $W\Omega_{\log}^i$ the logarithmic de Rham–Witt sheaf (see Notation 2.4).

We will prove Theorem 2.1 via comparison with syntomic cohomology, using crucially a rigidity result for the latter, due to Antieau–Mathew–Morrow–Nikolaus, [AMMN22, Theorem 5.2].

2.0.1. p -adic Tate twists. We begin by recalling some crucial results, due to Bhatt–Morrow–Scholze [BMS19], on the quasisyntomic sheaves $\mathbb{Z}_p(i)^{\text{syn}}$.

Notation 2.2. Let QSyn denote the quasisyntomic site, [BMS19, §4], and let $\text{QRSPerfd} \subset \text{QSyn}$ denote its basis consisting of the quasiregular semiperfectoid rings, [BMS19, Proposition 4.21].

For $i \in \mathbb{Z}_{\geq 0}$, we denote by $\mathbb{Z}_p(i)^{\text{syn}}$ the sheaf on QSyn introduced in [BMS19, §7.4], defined for $R \in \text{QRSPerfd}$ by

$$\mathbb{Z}_p(i)^{\text{syn}}(R) := \text{fib}(\text{Fil}_{\mathcal{N}}^i \widehat{\Delta}_R\{i\} \xrightarrow{\varphi_i-1} \widehat{\Delta}_R\{i\})$$

where, in terms of [BS22] and [BL22], we denote by $\widehat{\Delta}_R\{i\}$ the Breuil–Kisin twisted Nygaard completed absolute prismatic cohomology of R , we write $\text{Fil}_{\mathcal{N}}^*$ for the Nygaard filtration ([BMS19, Theorem 1.12(3)], [BS22, Theorem 13.1]), and we denote by φ_i the i -th divided Frobenius map.

For $n \in \mathbb{Z}_{\geq 1}$, we define the sheaf on QSyn

$$\mathbb{Z}/p^n(i)^{\text{syn}} := \mathbb{Z}_p(i)^{\text{syn}}/p^n.$$

In mixed characteristic, the sheaves defined above can be compared to p -adic nearby cycles, as we now recall. In the following, given a smooth formal scheme \mathfrak{X} over \mathcal{O}_C , we regard $\mathbb{Z}_p(i)^{\text{syn}}$ as a sheaf on the pro-étale site $\mathfrak{X}_{\text{proét}}$ using [BMS19, Remark 10.4].

Theorem 2.3 ([BMS19, Theorem 10.1]). *Let \mathfrak{X} be a smooth formal scheme over \mathcal{O}_C with generic fiber X . Denote by $\psi : X_{\text{proét}} \rightarrow \mathfrak{X}_{\text{proét}}$ the natural morphism of sites. Then, there is a natural isomorphism of sheaves of complexes on the pro-étale site $\mathfrak{X}_{\text{proét}}$*

$$\mathbb{Z}_p(i)^{\text{syn}} \simeq \tau^{\leq i} R\psi_* \mathbb{Z}_p(i) \quad (2.1)$$

and, for all $n \in \mathbb{Z}_{\geq 1}$, there is a natural isomorphism

$$\mathbb{Z}/p^n(i)^{\text{syn}} \simeq \tau^{\leq i} R\psi_* \mathbb{Z}/p^n(i). \quad (2.2)$$

Moreover, in equal characteristic p , the sheaves $\mathbb{Z}_p(i)^{\text{syn}}$ can be compared to the logarithmic de Rham–Witt sheaves. To state this result precisely, we introduce the following notation.

Notation 2.4. Let k be a perfect field of characteristic p , and let X be a smooth scheme over k .

We denote by

$$W\Omega^\bullet = \lim_n W_n\Omega^\bullet \quad \text{on } X_{\text{proét}}$$

the *de Rham–Witt complex* of Bloch–Deligne–Illusie of X , [Ill79, §I.1], regarded as a complex of pro-étale sheaves on X . It comes equipped with a Frobenius operator F and a Verschiebung operator V satisfying $FV = VF = p$. We denote by

$$W\Omega_{\log}^{\bullet} = \lim_n W_n\Omega_{\log}^{\bullet} \quad \text{on } X_{\text{proét}}$$

the *logarithmic de Rham–Witt complex* of X , [Ill79, §I.5.7]. We recall that each term $W\Omega_{\log}^i$ fits in the following exact sequence of pro-étale sheaves on X

$$0 \rightarrow W\Omega_{\log}^i \rightarrow W\Omega^i \xrightarrow{F-1} W\Omega^i \rightarrow 0,$$

[Ill79, I, Théorème 5.7.2].

Remark 2.5. We recall from [BMS19, Proposition 8.4] that we have isomorphisms

$$W\Omega^i \cong R\lim_n W_n\Omega^i, \quad W\Omega_{\log}^i \cong R\lim_n W_n\Omega_{\log}^i$$

on $X_{\text{proét}}$.

We will use several times the following result of Illusie.

Lemma 2.6 ([Ill79, I, Corollaire 5.7.5]). *Let k be a perfect field of characteristic p , and let X be a smooth scheme over k . For all $n \in \mathbb{Z}_{\geq 1}$ the natural map*

$$W\Omega_{\log}^i/p^n \rightarrow W_n\Omega_{\log}^i$$

is an isomorphism of pro-étale sheaves on X .

In the following, we regard $\mathbb{Z}_p(i)^{\text{syn}}$ as a sheaf on the pro-étale site $X_{\text{proét}}$.

Theorem 2.7 ([BMS19, Corollary 8.21]). *Let k be a perfect field of characteristic p , and let X be a smooth scheme over k . Then, there is a natural isomorphism of sheaves of complexes on the pro-étale site $X_{\text{proét}}$*

$$\mathbb{Z}_p(i)^{\text{syn}} \simeq W\Omega_{\log}^i[-i] \tag{2.3}$$

and, for all $n \in \mathbb{Z}_{\geq 1}$, there is a natural isomorphism

$$\mathbb{Z}/p^n(i)^{\text{syn}} \simeq W_n\Omega_{\log}^i[-i]. \tag{2.4}$$

Proof. The isomorphism (2.3) is [BMS19, Corollary 8.21]. Reducing (2.3) modulo p^n , we obtain the isomorphism (2.4), thanks to Lemma 2.6. \square

2.0.2. Rigidity of syntomic cohomology. Let $i \in \mathbb{Z}_{\geq 0}$. By [AMMN22, Theorem 5.1, (2)], one can extend the definition of the *integral syntomic cohomology*

$$\mathbb{Z}_p(i)^{\text{syn}}(R) = R\Gamma_{\text{syn}}(R, \mathbb{Z}_p(i))$$

to all p -adically complete rings R , via left Kan extension from finitely generated p -adically complete polynomial \mathbb{Z}_p -algebras.

We can then state the following rigidity result.

Theorem 2.8 ([AMMN22, Theorem 5.2]). *Let (R, I) be a henselian pair where R and R/I are p -adically complete rings. Then, the homotopy fiber*

$$\text{fib}(\mathbb{Z}_p(i)^{\text{syn}}(R) \rightarrow \mathbb{Z}_p(i)^{\text{syn}}(R/I))$$

is concentrated in cohomological degrees $\leq i$.

As an immediate consequence, we have the following. We will switch from algebraic to geometric notation to denote syntomic cohomology.

Corollary 2.9. *Let $\mathfrak{X} = \mathrm{Spf}(R)$ be an affine formal scheme over \mathcal{O}_C . Denote by k the residue field of C . Then, the homotopy fiber*

$$\mathrm{fib}(\mathbb{Z}_p(i)^{\mathrm{syn}}(\mathfrak{X}) \rightarrow \mathbb{Z}_p(i)^{\mathrm{syn}}(\mathfrak{X}_k))$$

is concentrated in cohomological degrees $\leq i$.

Proof. Denote by \mathfrak{m}_C the maximal ideal of \mathcal{O}_C , so that $k = \mathcal{O}_C/\mathfrak{m}_C$ is the residue field of C . We observe that the pair $(R, \mathfrak{m}_C R)$ is henselian: by [Sta24, Tag 0DYD], we can reduce to checking that (R, pR) and $(R/p, \mathfrak{m}_C R/p)$ are henselian pairs; for the first one, by [Sta24, Tag 0ALJ], it suffices to note that R is p -adically complete; for the second pair, we observe that the ideal $\mathfrak{m}_C R/p$ of R/p is locally nilpotent, and then we apply [Sta24, Tag 0ALI]. Thus, the statement follows by applying Theorem 2.8 to the henselian pair $(R, \mathfrak{m}_C R)$. \square

We are ready to prove the main result of this section.

Proof of Theorem 2.1. Considering the short exact sequences

$$0 \rightarrow H_{\mathrm{pro\acute{e}t}}^i(\mathfrak{X}_C, \mathbb{Z}_p(i))/p^n \rightarrow H_{\mathrm{pro\acute{e}t}}^i(\mathfrak{X}_C, \mathbb{Z}/p^n(i)) \rightarrow H_{\mathrm{pro\acute{e}t}}^{i+1}(\mathfrak{X}_C, \mathbb{Z}_p(i))[p^n] \rightarrow 0$$

and

$$0 \rightarrow H^i(\mathbb{Z}_p(i)^{\mathrm{syn}}(\mathfrak{X}))/p^n \rightarrow H^i(\mathbb{Z}/p^n(i)^{\mathrm{syn}}(\mathfrak{X})) \rightarrow H^{i+1}(\mathbb{Z}_p(i)^{\mathrm{syn}}(\mathfrak{X}))[p^n] \rightarrow 0,$$

using Theorem 2.3 (together with the compatibility of the isomorphisms (2.1) and (2.2) of *loc. cit.* under the reduction modulo p^n), we deduce that we have a natural isomorphism

$$H_{\mathrm{pro\acute{e}t}}^{i+1}(\mathfrak{X}_C, \mathbb{Z}_p(i))[p^n] \cong H^{i+1}(\mathbb{Z}_p(i)^{\mathrm{syn}}(\mathfrak{X}))[p^n]. \quad (2.5)$$

On the other hand, by Corollary 2.9, we have a natural isomorphism

$$H^{i+1}(\mathbb{Z}_p(i)^{\mathrm{syn}}(\mathfrak{X})) \cong H^{i+1}(\mathbb{Z}_p(i)^{\mathrm{syn}}(\mathfrak{X}_k)). \quad (2.6)$$

Therefore, combining (2.5), (2.6), and using Theorem 2.7, we deduce that

$$H_{\mathrm{pro\acute{e}t}}^{i+1}(\mathfrak{X}_C, \mathbb{Z}_p(i))[p^n] \cong H^1(\mathfrak{X}_k, W\Omega_{\log}^i)[p^n]. \quad (2.7)$$

We obtain the statement observing that, as C is algebraically closed, we can ignore the twist (i) on the left hand side of (2.7). \square

3. PRELIMINARIES ON LOG DE RHAM–WITT COHOMOLOGY

Notation 3.1. In this section, we denote by k an algebraically closed field of characteristic p .

Our goal here is to collect a number of known results on the logarithmic de Rham–Witt cohomology of smooth proper schemes over k that will be useful in the next section, when combined with Theorem 2.1. The main result that we will need is Corollary 3.6(ii).

We start with the following results of Illusie–Raynaud on the structural properties of the logarithmic de Rham–Witt cohomology. Similar results were also obtained by Milne, [Mil86, Lemma 1.8].

Lemma 3.2 ([IR83, IV, Corollaire 3.5(a)]). *Let X be a smooth proper scheme over k . For all $i, j \in \mathbb{Z}$, we have a short exact sequence*

$$0 \rightarrow H^j(X, W\Omega_{\log}^i) \rightarrow H^j(X, W\Omega^i) \xrightarrow{F-1} H^j(X, W\Omega^i) \rightarrow 0.$$

Remark 3.3. Given a smooth proper scheme X over k , in [IR83, §IV.3] the cohomology group $H^j(X, W\Omega_{\log}^i)$ denotes $\lim_n H^j(X, W_n\Omega_{\log}^i)$. However, it is also shown in the proof of [IR83, IV, Corollaire 3.5(a)], that the inverse system $\{H^{j-1}(X, W_n\Omega_{\log}^i)\}_n$ is Mittag-Leffler, therefore, in our notation $H^j(X, W\Omega_{\log}^i) := H^j(R\Gamma(X, W\Omega_{\log}^i))$, we have $H^j(X, W\Omega_{\log}^i) \simeq \lim_n H^j(X, W_n\Omega_{\log}^i)$.

Lemma 3.4 ([IR83]). *Let X be a smooth proper scheme over k . Let $i, j \in \mathbb{Z}$.*

(i) *We have a natural (in X) short exact sequence*

$$0 \rightarrow M^{ij}(X) \rightarrow H^j(X, W\Omega_{\log}^i) \rightarrow N^{ij}(X) \rightarrow 0$$

where $N^{ij}(X)$ is a finitely generated \mathbb{Z}_p -module and $M^{ij}(X)$ is killed by a power of p .

(ii) *The following facts are equivalent:*

(a) $M^{ij}(X) = 0$,

(b) $d : H^j(X, W\Omega^{i-1}) \rightarrow H^j(X, W\Omega^i)$ is the zero map,

(c) $H^j(X, W\Omega^i)^{V=0}$ is finite-dimensional over k , where V denotes the Verschiebung operator.

(iii) *The conditions in (ii) are satisfied for $j = 0, 1$.*

Proof. Parts (i) and (ii) follow from [IR83, IV, Théorème 3.3(b), Corollaire 3.5(b) & II, Corollaire 3.8]. For part (iii), we claim that, for $j = 0, 1$, we have that $H^j(X, W\Omega^i)^{V=0}$ is finite-dimensional over k . In fact, for $j = 0$, by [III79, II, Corollaire 2.17], we even have that $H^0(X, W\Omega^i)$ is a finite free module over $W(k)$. For $j = 1$, this is the content of (the proof of) [IR83, II, Corollaire 3.11]. \square

Remark 3.5. We note that some of the previous results were generalized to the log-smooth case by Lorenzon, [Lor02].

Let us now collect some consequences. For part (i) of the result below, see also [Pet24, Lemma A.2].

Corollary 3.6 ([III79], [IR83]). *Let X be a smooth proper scheme over k . Let $i \in \mathbb{Z}$.*

(i) *For all $j \in \mathbb{Z}$, we have an isomorphism of \mathbb{Z}_p -modules*

$$H^j(X, W\Omega_{\log}^i) \cong \mathbb{Z}_p^{\oplus r} \oplus T$$

for some $r \in \mathbb{Z}_{\geq 0}$, with T killed by a power of p .

(ii) *For $j = 0, 1$, the cohomology group $H^j(X, W\Omega_{\log}^i)$ is a finitely generated \mathbb{Z}_p -module, and for $j = 0$ it is also a free \mathbb{Z}_p -module.*

Proof. Part (i) readily follows from Lemma 3.4(i) and the classification of finitely generated \mathbb{Z}_p -modules. The first assertion of part (ii) follows combining part (i) and (iii) of Lemma 3.4. The last assertion follows recalling that $H^0(X, W\Omega_{\log}^i) = H^0(X, W\Omega^i)^{F=1}$ and, by [III79, II, Corollaire 2.17], $H^0(X, W\Omega^i)$ is a finite free module over $W(k)$. \square

Remark 3.7. Keeping the assumptions of Corollary 3.6, by [III79, II, Proposition 5.9], we also have that the cohomology group $H^2(X, W\Omega_{\log}^1)$ is a finitely generated \mathbb{Z}_p -module. On the other hand, for X a supersingular K3 surface we have that, $H^3(X, W\Omega_{\log}^1) \cong k$, [III79, §II.7.2], thus the torsion subgroup T in Corollary 3.6(i) can be infinite in general.

In the next section, we will only need the finiteness result of Corollary 3.6(ii). However, we chose to present an overview of the general structural properties of log de Rham–Witt cohomology to recall how Corollary 3.6(ii) can be deduced from such general results, and to illustrate that such a finiteness result can fail dramatically in general, as recalled in Remark 3.7 above.

4. FINITENESS OF TORSION AND APPLICATIONS

4.0.1. Finiteness. Our first goal here is to prove the following surprising finiteness result, Theorem 4.1, which thanks to Theorem 2.1 can be reduced to a problem in characteristic p .

Theorem 4.1. *Let \mathfrak{X} be an affine smooth formal scheme over \mathcal{O}_C . Then, for all $i \in \mathbb{Z}$, the cohomology group $H_{\text{proét}}^i(\mathfrak{X}_C, \mathbb{Z}_p)$ has finite torsion subgroup.*

Proof. The statement follows combining Theorem 2.1 and Proposition 4.2(ii) below. \square

The following is the crucial technical result that we use for the proof of Theorem 4.1. We will prove it by reduction to the proper case, via refined alterations and purity of the logarithmic de Rham–Witt sheaves.

Proposition 4.2. *Let k be an algebraically closed field of characteristic p . Let Y be a smooth variety over k .² Let $i \in \mathbb{Z}$.*

- (i) (Illusie–Raynaud) *For all $n \in \mathbb{Z}_{\geq 1}$, the cohomology group $H^0(Y, W_n \Omega_{\log}^i)$ is a finite abelian group.*
- (ii) *The cohomology group $H^1(Y, W \Omega_{\log}^i)$ has finite torsion subgroup.*

Proof. Part (i) in the case Y is smooth proper over k follows from Corollary 3.6(ii), using the short exact sequence

$$0 \rightarrow H^0(Y, W \Omega_{\log}^i) / p^n \rightarrow H^0(Y, W_n \Omega_{\log}^i) \rightarrow H^1(Y, W \Omega_{\log}^i)[p^n] \rightarrow 0, \quad (4.1)$$

which follows from Lemma 2.6.

Next, we want to show that we can reduce to the proper case. Let Y be a smooth variety over k . Using a refinement of de Jong’s alterations, [BS, Theorem 1.2], there exist an étale morphism $X \rightarrow Y$ over k , and a dense open immersion $X \hookrightarrow \overline{X}$ into a smooth projective variety \overline{X} over k whose complement is a strict normal crossing divisor $D \subset \overline{X}$. Since $W_n \Omega_{\log}^i$ satisfies étale descent, we have that $H^0(Y, W_n \Omega_{\log}^i)$ injects into $H^0(X, W_n \Omega_{\log}^i)$, hence it suffices to prove the statement with X in place of Y . For this, we consider the localization exact sequence

$$H_D^0(\overline{X}, W_n \Omega_{\log}^i) \rightarrow H^0(\overline{X}, W_n \Omega_{\log}^i) \rightarrow H^0(X, W_n \Omega_{\log}^i) \rightarrow H_D^1(\overline{X}, W_n \Omega_{\log}^i)$$

where H_D^\bullet denotes the cohomology with support in the closed subscheme $D \subset \overline{X}$. Then, using the purity of the logarithmic de Rham–Witt sheaves, [Gro85, II, Théorème 3.5.8, (3.5.19)], [Shi07, Theorem 3.1], we deduce the exact sequence

$$0 \rightarrow H^0(\overline{X}, W_n \Omega_{\log}^i) \rightarrow H^0(X, W_n \Omega_{\log}^i) \rightarrow \bigoplus_{r=1}^m H^0(D_r, W_n \Omega_{\log}^{i-1}) \quad (4.2)$$

where we wrote $D = \bigcup_{r=1}^m D_r$ as union of its irreducible components D_r . As \overline{X} and D_r , for $r = 1, \dots, m$, are smooth proper varieties over k , part (i) follows from the previous case and the exact sequence (4.2).

²Here, a *variety over k* is a separated, integral, scheme of finite type over $\text{Spec } k$.

For part (ii), let $M := H^1(Y, W\Omega_{\log}^i)$. By part (i) and the exact sequence (4.1) for $n = 1$, we have that $M[p]$ is a finite abelian group. Observing that M is derived p -adically complete,³ we deduce the statement from Lemma 4.3 below. \square

We used the following simple lemma.

Lemma 4.3. *Let M be a derived p -adically complete \mathbb{Z}_p -module. Suppose that the p -torsion submodule $M[p]$ is finite, then the torsion submodule $M_{\text{tors}} = \text{colim}_n M[p^n]$ is finite.*

Proof. We denote by $T_p(M) := \lim_n M[p^n]$ the p -adic Tate module of M , where the transition maps $M[p^n] \rightarrow M[p^{n-1}]$ are given by multiplication by p . Since M is derived p -adically complete (and concentrated in degree 0), by [Sta24, Tag 0BKG] we have that

$$T_p(M) = H^{-1}(M^\wedge) = H^{-1}(M) = 0$$

where $(-)^\wedge$ denotes the derived p -adic completion. As $M[p]$ is finite, we deduce that there exists a sufficiently big integer m such that the map $M[p^{m+1}] \rightarrow M[p]$, given by multiplication by p^m , is the zero map. In other words, $M_{\text{tors}} = M[p^m]$. Now, using the exact sequence

$$0 \rightarrow M[p^{m-1}] \rightarrow M[p^m] \rightarrow M[p]$$

where the right map is given by multiplication by p^{m-1} , arguing by induction we deduce that $M[p^m]$ is finite. Putting everything together, we conclude that M_{tors} is finite, as desired. \square

Remark 4.4. In the following, we keep the notation of Proposition 4.2.

- (1) After a first draft of this note was distributed, Alexander Petrov informed us that Proposition 4.2(i) can also be deduced from [Ill79, Corollaire 2.5.6, Remarque 2.5.7 (Page 534)]⁴. In fact, the latter states the following: denote $\nu(i)$ the étale sheaf on Y defined by

$$\nu(i) := \ker(Z\Omega^i \xrightarrow{C-1} \Omega^i)$$

where we write $Z\Omega^i \subset \Omega^i$ for the subsheaf of cocycles (i.e. the kernel of the differential $d : \Omega^i \rightarrow \Omega^{i+1}$), and we denote by C the composite $Z\Omega^i \rightarrow H^i(\Omega^\bullet) \cong \Omega^i$, where the latter is the Cartier isomorphism; then, we have that

$$H^0(Y, \nu(i)) \text{ is finite-dimensional over } \mathbb{F}_p. \quad (4.3)$$

This combined with the comparison between the sheaves $\nu(i)$ and Ω_{\log}^i , [Ill79, Théorème 2.4.2 (Page 528)] proves Proposition 4.2(i) for $n = 1$. Then, Proposition 4.2(i) for a general $n \in \mathbb{Z}_{\geq 1}$ can be deduced from the case $n = 1$. For this, we can argue by induction on n , using the exact sequence of pro-étale sheaves on Y , which follows from Lemma 2.6:

$$0 \rightarrow W_{n-1}\Omega_{\log}^i \rightarrow W_n\Omega_{\log}^i \rightarrow \Omega_{\log}^i \rightarrow 0$$

where the right map is given by reduction modulo p .

³In fact, $R\Gamma(Y, W\Omega_{\log}^i) = R\lim_n R\Gamma(Y, W_n\Omega_{\log}^i)$ is derived p -adically complete, as it is limit of p^n -torsion complexes. Hence, we can use [BS15, Proposition 3.4.4].

⁴As stated by Illusie before [Ill79, Proposition 2.5.2 (Page 532)], these results are based on some unpublished notes of Raynaud. That is the reason why we attributed Proposition 4.2(i) to Illusie and Raynaud.

- (2) Using the recent stacky approach to syntomic cohomology by Drinfeld and Bhatt–Lurie, one can show, using the same arguments as in [Bha22, Proposition 4.4.7], that we have a natural isomorphism

$$\mathbb{F}_p(i)^{\text{syn}}(Y) \simeq R\Gamma(Y, \nu(i))[-i]$$

where we kept the notation of point (1). The latter isomorphism combined with (4.3) implies that $H^i(\mathbb{F}_p(i)^{\text{syn}}(Y))$ is finite-dimensional over \mathbb{F}_p . In particular, using this one can give a proof of Theorem 4.1 bypassing the use of log de Rham–Witt cohomology. We remark that it would also be interesting to give a proof of Corollary 2.9 purely in terms of the stacky approach to syntomic cohomology.

We observe the following easy consequence of Theorem 4.1.

Corollary 4.5. *Let \mathfrak{X} be an affine smooth formal scheme over \mathcal{O}_C . Then, for all $i \in \mathbb{Z}$, we have a natural isomorphism*

$$H_{\text{proét}}^i(\mathfrak{X}_C, \mathbb{Z}_p) \xrightarrow{\sim} \lim_n H_{\text{proét}}^i(\mathfrak{X}_C, \mathbb{Z}/p^n).$$

Proof. Considering the short exact sequence

$$0 \rightarrow H_{\text{proét}}^{i-1}(\mathfrak{X}_C, \mathbb{Z}_p)/p^n \rightarrow H_{\text{proét}}^{i-1}(\mathfrak{X}_C, \mathbb{Z}/p^n) \rightarrow H_{\text{proét}}^i(\mathfrak{X}_C, \mathbb{Z}_p)[p^n] \rightarrow 0,$$

by Theorem 4.1 and the Mittag-Leffler criterion, [Gro61, Proposition 13.2.2], we have that

$$R^1 \lim_n H_{\text{proét}}^{i-1}(\mathfrak{X}_C, \mathbb{Z}/p^n) = 0.$$

This implies the statement. \square

4.0.2. p -torsion in the Picard group. Let us now collect another interesting consequence of Proposition 4.2(i). A proof of the following result was sketched in [GJRW96, Lemma 2.8], assuming resolution of singularities in characteristic p .

Proposition 4.6. *Let k be an algebraically closed field of characteristic p , and let Y be a smooth variety over k . Then, for all $n \in \mathbb{Z}_{\geq 0}$,*

$$\text{Pic}(Y)[p^n] \text{ is finite.}$$

Proof. By [BMS19, Proposition 7.17], we have a natural isomorphism

$$\mathbb{Z}_p(1)^{\text{syn}}(Y) \simeq R\Gamma_{\text{ét}}(Y, \mathbb{G}_m)^\wedge[-1]$$

where $(-)^\wedge$ denotes the derived p -adic completion. Combining this with Theorem 2.4, we deduce that we have a short exact sequence

$$0 \rightarrow H^0(Y, \mathbb{G}_m)/p^n \rightarrow H^0(Y, W_n\Omega^1) \rightarrow \text{Pic}(Y)[p^n] \rightarrow 0.$$

In particular, $H^0(Y, W_n\Omega^1)$ surjects onto $\text{Pic}(Y)[p^n]$, and then the statement follows from Proposition 4.2(i). \square

Corollary 4.7. *Let \mathfrak{X} be an affine smooth formal scheme over \mathcal{O}_C . Then, for all $n \in \mathbb{Z}_{\geq 0}$,*

$$\text{Pic}(\mathfrak{X}_C)[p^n] \text{ is finite.}$$

Proof. Let k denote the residue field of C . By [Lüt16, Lemma 6.2.4], [Heu21, Lemma 3.6], we have natural isomorphisms of Picard groups

$$\text{Pic}(\mathfrak{X}_C) \cong \text{Pic}(\mathfrak{X}) \cong \text{Pic}(\mathfrak{X}_k).$$

Then, the statement follows from Proposition 4.6. \square

4.0.3. Rational p -adic étale cohomology of affinoids. As a corollary of Theorem 4.1, we will deduce that, for X an affinoid rigid space over C having an affine smooth formal model over \mathcal{O}_C , the rational p -adic pro-étale cohomology groups of X are \mathbb{Q}_p -Banach spaces.

Notation 4.8. We recall from 1.6 that we fixed an uncountable cardinal κ as in [Sch21, Lemma 4.1]. We denote by CondAb the category of κ -condensed abelian groups (the prefix “ κ ” will be tacit), [CS19, Definition 2.1]. We recall that CondAb identifies with the category of sheaves of abelian groups on $\text{Spa}(C)_{\text{proét}}$ (see e.g. [Bos21, Remark 2.9]).

For an adic space $f : X \rightarrow \text{Spa}(C)$, and a sheaf of abelian groups \mathcal{F} on $X_{\text{proét}}$, we will regard the pro-étale cohomology complex $R\Gamma_{\text{proét}}(X, \mathcal{F}) = Rf_{\text{proét},*}\mathcal{F}$ in the derived category $D(\text{CondAb})$ (see also [Bos21, Definition 2.7]).

Proposition 4.9. *Let X be a quasi-compact and quasi-separated rigid-analytic variety over C . Let $i \in \mathbb{Z}$. If $H_{\text{proét}}^i(X, \mathbb{Z}_p)$ has bounded p^∞ -torsion, then the condensed \mathbb{Q}_p -vector space $H_{\text{proét}}^i(X, \mathbb{Q}_p)$ is a Banach space.*

For the proof, we will use the following general lemma. We refer the reader to [BS15, §3.4] for the definition of derived completion in a general replete topos, as well as its basic properties, which will be used freely in the following. In particular, we denote $(-)_p^\wedge : D(\text{CondAb}) \rightarrow D(\text{CondAb})$ the derived p -adic completion.

Lemma 4.10. *Let M be a condensed abelian group, and denote by $M(*)$ its underlying abelian group. Suppose that the following conditions are satisfied:*

- (i) *the natural maps of condensed abelian groups $M \rightarrow M_p^\wedge \leftarrow (M(*))^{\text{disc}}_p^\wedge$ are isomorphisms, where we regard $M(*)$ as a discrete condensed abelian group;*
- (ii) *$M(*)$ has bounded p^∞ -torsion.*

Then, the condensed \mathbb{Q}_p -vector space $M[1/p]$ is a Banach space.

Proof. By condition (i) there exists an exact sequence of condensed abelian groups as follows:

$$\left(\bigoplus_I \mathbb{Z}\right)_p^\wedge \xrightarrow{f} \left(\bigoplus_J \mathbb{Z}\right)_p^\wedge \rightarrow M \rightarrow 0$$

for some sets I, J . We write N for the target of the map f and L for the image of f , so that we have a short exact sequence of condensed abelian groups

$$0 \rightarrow L \rightarrow N \rightarrow M \rightarrow 0. \quad (4.4)$$

Using, in addition, condition (ii) we deduce that the exact sequence (4.4) is given by taking the limit over $n \in \mathbb{Z}_{\geq 0}$ of the exact sequence of inverse systems

$$\{M(*)[p^n]\} \rightarrow \{L(*)/p^n\} \rightarrow \{N(*)/p^n\} \rightarrow \{M(*)/p^n\} \rightarrow 0$$

where all the objects in sight are regarded as discrete condensed abelian groups. We deduce in particular that, inverting p in (4.4), $L[1/p]$ is a closed subspace of the \mathbb{Q}_p -Banach space $N[1/p]$, and then $M[1/p]$ is a \mathbb{Q}_p -Banach space too, as desired. \square

Remark 4.11. Let us denote by $D_{p\text{-comp}}(\text{Ab}) \subset D(\text{Ab})$ the category of derived p -complete complexes of abelian groups. Consider the functor

$$D_{p\text{-comp}}(\text{Ab}) \rightarrow D(\text{CondAb}) : K \mapsto (K^{\text{disc}})_p^\wedge \quad (4.5)$$

where K^{disc} denotes the complex of condensed abelian group associated with K endowed with the discrete topology, and we write $(-)_p^\wedge$ for the derived p -adic completion. We observe that the functor (4.5) induces a t -exact equivalence (for the natural t -structures) between $D_{p\text{-comp}}(\text{Ab})$ and the full subcategory of $D(\text{CondAb})$ spanned by the complexes M such that the natural maps $M \rightarrow M_p^\wedge \leftarrow (M(*))^{\text{disc}}_p^\wedge$ are isomorphisms.

Proof of Proposition 4.9. We want to apply Lemma 4.10 with $M := H_{\text{proét}}^i(X, \mathbb{Z}_p)$. It suffices to show that M satisfies the condition (i) of the latter lemma; then, the statement follows observing that $H_{\text{proét}}^i(X, \mathbb{Q}_p) = H_{\text{proét}}^i(X, \mathbb{Z}_p)[1/p]$, as X is quasi-compact and quasi-separated. In view of Remark 4.11, and fact that if a complex of abelian groups is derived p -complete then each of its cohomology groups is so, [BS15, Proposition 3.4.4], we can reduce to showing that $R\Gamma_{\text{proét}}(X, \mathbb{Z}_p)$ lies in the essential image of the functor (4.5). Thus, recalling that $R\Gamma_{\text{proét}}(X, \mathbb{Z}_p) = R\lim_n R\Gamma_{\text{proét}}(X, \mathbb{Z}/p^n)$, it suffices to show that $R\Gamma_{\text{proét}}(X, \mathbb{Z}/p^n)$ is a complex of discrete condensed abelian groups, for all $n \in \mathbb{Z}_{\geq 1}$. This follows from [Sch21, Proposition 16.6]. \square

Combining Theorem 4.1 and Proposition 4.9, we obtain the following result.

Theorem 4.12. *Let \mathfrak{X} be an affine smooth formal scheme over \mathcal{O}_C . Then, for all $i \in \mathbb{Z}$, the condensed \mathbb{Q}_p -vector space $H_{\text{proét}}^i(\mathfrak{X}_C, \mathbb{Q}_p)$ is a Banach space.*

4.0.4. Beyond the affine case. Our next goal is to extend Theorem 2.1 to a wider class of formal schemes.

Notation 4.13. In this section, we denote by k the residue field of C .

In the proof of the following theorem, it will be crucial once again the finiteness result stated in Proposition 4.2(i).

Theorem 4.14. *Let \mathfrak{X} be a smooth formal scheme over \mathcal{O}_C that admits an open affine covering $\{\mathfrak{U}_r\}_{r \in \mathbb{Z}_{\geq 0}}$ with $\mathfrak{U}_r \subseteq \mathfrak{U}_{r+1}$. Then, for all $i \in \mathbb{Z}$ and $n \in \mathbb{Z}_{\geq 0}$ we have a natural isomorphism*

$$H_{\text{proét}}^{i+1}(\mathfrak{X}_C, \mathbb{Z}_p)[p^n] \cong H^1(\mathfrak{X}_k, W\Omega_{\log}^i)[p^n].$$

Proof. It suffices to show that Corollary 2.9 extends to \mathfrak{X} a smooth formal scheme over \mathcal{O}_C as in the statement, and then we can use the same argument of the proof of Theorem 2.1. Namely, we want to show that the homotopy fiber

$$\text{fib}(\mathbb{Z}_p(i)^{\text{syn}}(\mathfrak{X}) \rightarrow \mathbb{Z}_p(i)^{\text{syn}}(\mathfrak{X}_k))$$

is concentrated in cohomological degrees $\leq i$. First, we note that, by the derived Nakayama lemma, the latter assertion is equivalent to its version modulo p , i.e. replacing $\mathbb{Z}_p(i)^{\text{syn}}$ by $\mathbb{F}_p(i)^{\text{syn}}$. In other words, we want to prove that, for all $j \geq i + 1$, the natural map

$$H^j(\mathbb{F}_p(i)^{\text{syn}}(\mathfrak{X})) \rightarrow H^j(\mathbb{F}_p(i)^{\text{syn}}(\mathfrak{X}_k))$$

is an isomorphism. Consider the short exact sequence

$$0 \rightarrow R^1 \lim_r H^{j-1}(\mathbb{F}_p(i)^{\text{syn}}(\mathfrak{U}_r)) \rightarrow H^j(\mathbb{F}_p(i)^{\text{syn}}(\mathfrak{X})) \rightarrow \lim_r H^j(\mathbb{F}_p(i)^{\text{syn}}(\mathfrak{U}_r)) \rightarrow 0$$

and the analogous short exact sequence with \mathfrak{X} replaced by \mathfrak{X}_k and \mathfrak{U}_r replaced by its special fiber $U_r := (\mathfrak{U}_r)_k$. By Corollary 2.9 (applied to each affine \mathfrak{U}_r), we deduce that it is sufficient to show that

$$R^1 \lim_r H^i(\mathbb{F}_p(i)^{\text{syn}}(\mathfrak{U}_r)) = 0 \tag{4.6}$$

and

$$R^1 \lim_r H^i(\mathbb{F}_p(i)^{\text{syn}}(U_r)) = 0. \quad (4.7)$$

First of all, we note that (4.6) holds true if and only if (4.7) holds true. In fact, (4.6) is equivalent to $H^{i+1}(\mathbb{F}_p(i)^{\text{syn}}(\mathfrak{U}_r)) \xrightarrow{\sim} \lim_r H^{i+1}(\mathbb{F}_p(i)^{\text{syn}}(\mathfrak{U}_r))$, which, in turn, again by Corollary 2.9, is equivalent to (4.7). Next, we observe that, by Theorem 2.7, the vanishing (4.7) is equivalent to

$$R^1 \lim_r H^0(U_r, \Omega_{\log}^i) = 0. \quad (4.8)$$

Therefore, we are reduced to check the vanishing (4.8): this follows from Proposition 4.2(i) and the Mittag-Leffler criterion, [Gro61, Proposition 13.2.2]. \square

At this point, it is natural to conjecture an extension of Theorem 4.14 to the semistable reduction case. In order to state the conjecture precisely, we need to introduce some notation.

Notation 4.15. We say that a formal scheme over \mathcal{O}_C is *semistable* if it has étale locally semistable coordinates in the sense of [CK19, §1.5]. Given a semistable formal scheme \mathfrak{X} over \mathcal{O}_C we endow \mathfrak{X} with the canonical log structure, i.e. the one given by the sheafification of the subpresheaf $\mathcal{O}_{\mathfrak{X}, \text{ét}} \cap (\mathcal{O}_{\mathfrak{X}, \text{ét}}[1/p])^\times \hookrightarrow \mathcal{O}_{\mathfrak{X}, \text{ét}}$, and the special fiber \mathfrak{X}_k with the pullback log structure.

We write k^0 for k with the log structure associated with $\mathbb{Z}_{\geq 0} \rightarrow k$, $1 \mapsto 0$, $0 \mapsto 1$.

Conjecture 4.16. *Let \mathfrak{X} be a semistable formal scheme over \mathcal{O}_C that admits an open affine covering $\{\mathfrak{U}_r\}_{r \in \mathbb{Z}_{\geq 0}}$ with $\mathfrak{U}_r \subseteq \mathfrak{U}_{r+1}$. Then, for all $i \in \mathbb{Z}$ and $n \in \mathbb{Z}_{\geq 0}$ we have a natural isomorphism*

$$H_{\text{proét}}^{i+1}(\mathfrak{X}_C, \mathbb{Z}_p)[p^n] \cong H^1(\mathfrak{X}_k/k^0, W\Omega_{\log}^i)[p^n].$$

Here, we use the logarithmic de Rham-Witt cohomology for log schemes as defined in [Lor02].

REFERENCES

- [AMMN22] Benjamin Antieau, Akhil Mathew, Matthew Morrow, and Thomas Nikolaus, *On the Beilinson fiber square*, Duke Math. J. **171** (2022), no. 18, 3707–3806.
- [Bha22] Bhargav Bhatt, *Prismatic F -Gauges*, <https://www.math.ias.edu/~bhatt/teaching/mat549f22/lectures.pdf>, 2022.
- [BL22] Bhargav Bhatt and Jacob Lurie, *Absolute prismatic cohomology*, <https://arxiv.org/abs/2201.06120>, 2022, Preprint.
- [BMS18] Bhargav Bhatt, Matthew Morrow, and Peter Scholze, *Integral p -adic Hodge theory*, Publ. Math. Inst. Hautes Études Sci. **128** (2018), 219–397.
- [BMS19] ———, *Topological Hochschild homology and integral p -adic Hodge theory*, Publ. Math. Inst. Hautes Études Sci. **129** (2019), 199–310.
- [Bos21] Guido Bosco, *On the p -adic pro-étale cohomology of Drinfeld symmetric spaces*, <https://arxiv.org/abs/2110.10683>, 2021, Preprint.
- [Bos23] ———, *Rational p -adic Hodge theory for rigid-analytic varieties*, <https://arxiv.org/abs/2306.06100>, 2023, Preprint.
- [BS] Bhargav Bhatt and Andrew Snowden, *Refined alterations*, <http://www-personal.umich.edu/~bhattb/math/alterationsepsilon.pdf>.
- [BS15] Bhargav Bhatt and Peter Scholze, *The pro-étale topology for schemes*, Astérisque (2015), no. 369, 99–201.
- [BS22] ———, *Prisms and prismatic cohomology*, Ann. of Math. (2) **196** (2022), no. 3, 1135–1275.
- [CDN20a] Pierre Colmez, Gabriel Dospinescu, and Wiesława Nizioł, *Cohomologie p -adique de la tour de Drinfeld: le cas de la dimension 1*, J. Amer. Math. Soc. **33** (2020), no. 2, 311–362.
- [CDN20b] ———, *Cohomology of p -adic Stein spaces*, Invent. Math. **219** (2020), no. 3, 873–985.
- [CDN21] ———, *Integral p -adic étale cohomology of Drinfeld symmetric spaces*, Duke Math. J. **170** (2021), no. 3, 575–613.

- [CDN22] ———, *Cohomologie des courbes analytiques p -adiques*, Camb. J. Math. **10** (2022), no. 3, 511–655.
- [CK19] Kestutis Cesnavicius and Teruhisa Koshikawa, *The A_{inf} -cohomology in the semistable case*, Compos. Math. **155** (2019), no. 11, 2039–2128.
- [CS19] Dustin Clausen and Peter Scholze, *Lectures on Condensed Mathematics*, <https://www.math.uni-bonn.de/people/scholze/Condensed.pdf>, 2019.
- [GJRW96] Robert Guralnick, David B. Jaffe, Wayne Raskind, and Roger Wiegand, *On the Picard group: torsion and the kernel induced by a faithfully flat map*, J. Algebra **183** (1996), no. 2, 420–455.
- [GK05] Elmar Grosse-Klönne, *Frobenius and monodromy operators in rigid analysis, and Drinfel'd's symmetric space*, J. Algebraic Geom. **14** (2005), no. 3, 391–437.
- [Gro61] A. Grothendieck, *Éléments de géométrie algébrique. III. Étude cohomologique des faisceaux cohérents. I*, Inst. Hautes Études Sci. Publ. Math. (1961), no. 11, 167.
- [Gro85] Michel Gros, *Classes de Chern et classes de cycles en cohomologie de Hodge-Witt logarithmique*, Mém. Soc. Math. France (N.S.) (1985), no. 21, 87.
- [Heu21] Ben Heuer, *Line bundles on perfectoid covers: case of good reduction*, <https://arxiv.org/abs/2105.05230>, 2021, Preprint.
- [Ill79] Luc Illusie, *Complexe de de Rham-Witt et cohomologie cristalline*, Ann. Sci. École Norm. Sup. (4) **12** (1979), no. 4, 501–661.
- [IR83] Luc Illusie and Michel Raynaud, *Les suites spectrales associées au complexe de de Rham-Witt*, Inst. Hautes Études Sci. Publ. Math. (1983), no. 57, 73–212.
- [Lor02] Pierre Lorenzon, *Logarithmic Hodge-Witt forms and Hyodo-Kato cohomology*, J. Algebra **249** (2002), no. 2, 247–265.
- [Lüt16] Werner Lütkebohmert, *Rigid geometry of curves and their Jacobians*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], vol. 61, Springer, Cham, 2016.
- [Mil86] J. S. Milne, *Values of zeta functions of varieties over finite fields*, Amer. J. Math. **108** (1986), no. 2, 297–360.
- [Pet24] Alexander Petrov, *Appendix to "Boundedness of the p -primary torsion of the Brauer group of products of varieties"*, by Alexei N. Skorobogatov, <https://arxiv.org/pdf/2404.19150>, 2024, Preprint.
- [Sch13] Peter Scholze, *p -adic Hodge theory for rigid-analytic varieties*, Forum Math. Pi **1** (2013), e1, 77.
- [Sch21] ———, *Étale cohomology of diamonds*, <https://arxiv.org/abs/1709.07343>, 2021, Preprint.
- [Shi07] Atsushi Shiho, *On logarithmic Hodge-Witt cohomology of regular schemes*, J. Math. Sci. Univ. Tokyo **14** (2007), no. 4, 567–635.
- [Sta24] The Stacks Project Authors, *Stacks Project*, <https://stacks.math.columbia.edu>, 2024.

MAX-PLANCK-INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, 53111 BONN, GERMANY
E-mail address: bosco@mpim-bonn.mpg.de