



## COMBINATORIAL APPLICATIONS OF THE HODGE–RIEMANN RELATIONS

June Huh

### Abstract

Why do natural and interesting sequences often turn out to be log-concave? We give one of many possible explanations, from the viewpoint of “standard conjectures”. We illustrate with several examples from combinatorics.

### 1 Log-concave and unimodal sequences

Logarithmic concavity is a property of a sequence of real numbers, occurring throughout algebra, geometry, and combinatorics. A sequence of real numbers  $a_0, \dots, a_d$  is *log-concave* if

$$a_i^2 \geq a_{i-1}a_{i+1} \text{ for all } i.$$

When all the entries are positive, the log-concavity implies unimodality, a property easier to visualize: the sequence is *unimodal* if there is an index  $i$  such that

$$a_0 \leq \dots \leq a_{i-1} \leq a_i \geq a_{i+1} \geq \dots \geq a_d.$$

A rich variety of log-concave and unimodal sequences arising in combinatorics can be found in the surveys [Brenti \[1994\]](#) and [Stanley \[1989, 2000\]](#). For an extensive discussion of log-concavity and its applications in probability and statistics, see [Dharmadhikari and Joag-Dev \[1988\]](#), [Marshall, Olkin, and Arnold \[2011\]](#), and [Saumard and Wellner \[2014\]](#).

Why do natural and interesting sequences often turn out to be log-concave? Below we give one of many possible explanations, from the viewpoint of *standard conjectures*. To illustrate, we discuss three combinatorial sequences appearing in [Stanley \[2000, Problem 25\]](#), in Sections [2.4](#), [2.5](#), and [2.8](#). Another heuristic, based on the physical principle that the entropy of a system should be concave as a function of the energy, can be found in [Okounkov \[2003\]](#).

Let  $X$  be a mathematical object of “dimension”  $d$ . Often it is possible to construct from  $X$  in a natural way a graded vector space over the real numbers

$$A(X) = \bigoplus_{q=0}^d A^q(X),$$

together with a symmetric bilinear map  $P : A(X) \times A(X) \rightarrow \mathbb{R}$  and a graded linear map  $L : A^\bullet(X) \rightarrow A^{\bullet+1}(X)$  that is symmetric with respect to  $P$ . The linear operator  $L$  usually comes in as a member of a family  $K(X)$ , a convex cone in the space of linear operators on  $A(X)$ .<sup>1</sup> For example,  $A(X)$  may be the cohomology of real  $(q, q)$ -forms on a compact Kähler manifold (Gromov [1990]), the ring of algebraic cycles modulo homological equivalence on a smooth projective variety (Grothendieck [1969]), McMullen's algebra generated by the Minkowski summands of a simple convex polytope (McMullen [1993]), the combinatorial intersection cohomology of a convex polytope (Karu [2004]), the reduced Soergel bimodule of a Coxeter group element (Elias and Williamson [2014]), or the Chow ring of a matroid (Section 2.6).

Often, but not always,  $A(X)$  has the structure of a graded algebra,  $P$  is determined by the multiplicative structure of  $A(X)$  up to a constant multiple, and  $L$  is the multiplication by an element in  $A^1(X)$ . In any case, we expect the following properties to hold for the triple  $(A(X), P(X), K(X))$  for every nonnegative integer  $q \leq \frac{d}{2}$ :

(PD) The bilinear pairing

$$A^q(X) \times A^{d-q}(X) \longrightarrow \mathbb{R}, \quad (\eta, \xi) \longmapsto P(\eta, \xi)$$

is nondegenerate (the Poincaré duality for  $X$ ).

(HL) For any  $L_1, \dots, L_{d-2q} \in K(X)$ , the linear map

$$A^q(X) \longrightarrow A^{d-q}(X), \quad \eta \longmapsto \left( \prod_{i=1}^{d-2q} L_i \right) \eta$$

is bijective (the hard Lefschetz theorem for  $X$ ).

(HR) For any  $L_0, L_1, \dots, L_{d-2q} \in K(X)$ , the bilinear form

$$A^q(X) \times A^q(X) \longrightarrow \mathbb{R}, \quad (\eta_1, \eta_2) \longmapsto (-1)^q P(\eta_1, \left( \prod_{i=1}^{d-2q} L_i \right) \eta_2)$$

is positive definite on the kernel of the linear map

$$A^q(X) \longrightarrow A^{d-q+1}(X), \quad \eta \longmapsto \left( \prod_{i=0}^{d-2q} L_i \right) \eta$$

(the Hodge-Riemann relation for  $X$ ).

All three properties are known to hold for the objects listed above except one, which is the subject of Grothendieck's standard conjectures on algebraic cycles. The known proofs of the hard Lefschetz theorems and the Hodge-Riemann relations for different objects have certain structural similarities, but there is no known way of deducing one of them from the others.

<sup>1</sup>“P” is for Poincaré, “L” is for Lefschetz, and “K” is for Kähler.

Hard Lefschetz theorems for various  $X$ 's have found numerous applications to problems of combinatorial nature. An early survey of these applications can be found in Stanley [1984]. We highlight the following three:

- (1) Erdős-Moser conjecture (Erdős [1965]), proved by Stanley [1980b]: Let  $E$  be a subset of  $\mathbb{R}$  and let  $f(E, k)$  be the number of subsets of  $E$  whose elements sum to  $k$ . If the cardinality of  $E$  is  $2n + 1$ , then

$$f(E, k) \leq f([-n, n] \cap \mathbb{Z}, 0).$$

- (2) McMullen's  $g$ -conjecture (McMullen [1971]), proved by Billera and Lee [1980] and Stanley [1980a]: The  $f$ -vector of a  $d$ -dimensional convex polytope  $P$  is the sequence  $f_0(P), \dots, f_d(P)$ , where

$$f_i(P) = \text{the number of } (i - 1)\text{-dimensional faces of } P.$$

The  $h$ -vector of  $P$  is the sequence  $h_0(P), \dots, h_d(P)$  defined by the identity

$$\sum_{i=0}^d h_i(P)x^i = \sum_{i=0}^d f_i(P)x^i(1-x)^{d-i}.$$

The  $g$ -conjecture gives a complete numerical characterization of the  $h$ -vectors of simplicial polytopes. In particular, for any  $d$ -dimensional simplicial polytope  $P$ ,

$$h_i(P) = h_{d-i}(P) \text{ and } h_i(P) \leq h_{i+1}(P) \text{ for all } i < d/2.$$

- (3) Dowling-Wilson conjecture (Dowling and Wilson [1974, 1975]), proved by Huh and Wang [2017]: Let  $E$  be a finite subset of a vector space, and let  $w_i(E)$  be the number of  $i$ -dimensional subspaces spanned by subsets of  $E$ . If  $E$  spans a  $d$ -dimensional subspace, then

$$w_i(E) \leq w_{d-i}(E) \text{ and } w_i(E) \leq w_{i+1}(E) \text{ for all } i < d/2.$$

All known proofs of the above statements use some version of HL.

When the Poincaré duality for  $X$  is known, the Hodge-Riemann relation for  $X$  is stronger than the hard Lefschetz theorem for  $X$  in the sense that, for every  $q$ ,

$$\text{HR in degrees at most } q \implies \text{HL in degrees at most } q.$$

In the remainder of this survey, we give an overview of applications of the Hodge-Riemann relations to concrete problems. We remark that most known applications only use the following immediate consequence of HR in degrees  $q \leq 1$ : For any  $L_1, \dots, L_{d-2} \in K(X)$ , any matrix representing the symmetric bilinear form

$$A^1(X) \times A^1(X) \longrightarrow \mathbb{R}, \quad (\eta_1, \eta_2) \longmapsto P(\eta_1, \left( \prod_{i=0}^{d-2} L_i \right) \eta_2)$$

has exactly one positive eigenvalue. One notable exception is the implication

Grothendieck standard conjectures on algebraic cycles  $\implies$

Weil conjectures on zeta functions over finite fields,

which was one of the main motivations for formulating the standard conjectures (Colmez and Serre [2001], Kleiman [1968], and Kleiman [1994]). It will be interesting to find applications of HR for  $q > 1$  in other contexts too.

## 2 Applications of the Hodge-Riemann relations

**2.1 Mixed discriminants and permanents.** The notion of mixed discriminant arises when one combines the determinant with the matrix sum. To define the mixed discriminant, let  $\mathbf{A} = (A_1, \dots, A_d)$  be a collection of real symmetric  $d \times d$  matrices, and consider the function

$$\det_{\mathbf{A}} : \mathbb{R}^d \longrightarrow \mathbb{R}, \quad (t_1, \dots, t_d) \longmapsto \det(t_1 A_1 + \dots + t_d A_d),$$

which is a homogeneous polynomial of degree  $d$ . The number

$$D(A_1, \dots, A_d) = \frac{\partial^d}{\partial t_1 \dots \partial t_d} \det_{\mathbf{A}}(t_1, \dots, t_d)$$

is called the *mixed discriminant* of  $\mathbf{A}$ . The mixed discriminant is symmetric in  $\mathbf{A}$ , and it is nonnegative whenever all the matrices in  $\mathbf{A}$  are positive semidefinite.<sup>2</sup>

Let  $\mathbf{P} = (P_1, \dots, P_{d-2})$  be any collection of  $d \times d$  positive semidefinite matrices. Define a symmetric bilinear form  $\text{HR}(\mathbf{P})$  on the space of real symmetric  $d \times d$  matrices by

$$\text{HR}(\mathbf{P}) : \text{Sym}_d \times \text{Sym}_d \longrightarrow \mathbb{R}, \quad (\eta_1, \eta_2) \longmapsto D(\eta_1, \eta_2, P_1, \dots, P_{d-2}).$$

Aleksandrov [1938] proved the following statement and used it in his proof of the Aleksandrov-Fenchel inequality for mixed volumes of convex bodies. To avoid trivialities, we suppose that  $\text{HR}(\mathbf{P})$  is not identically zero.

**Theorem 1.** Any matrix representing  $\text{HR}(\mathbf{P})$  has exactly one positive eigenvalue.

It follows from Cauchy's eigenvalue interlacing theorem that, for any positive semidefinite  $d \times d$  matrices  $A_1, \dots, A_d$ ,

$$\det \begin{pmatrix} D(A_1, A_1, A_3, \dots, A_d) & D(A_1, A_2, A_3, \dots, A_d) \\ D(A_1, A_2, A_3, \dots, A_d) & D(A_2, A_2, A_3, \dots, A_d) \end{pmatrix} \leq 0.$$

**Theorem 1** is, in fact, a Hodge-Riemann relation in degree 1. The object  $X$  is the  $d$ -dimensional complex vector space  $\mathbb{C}^d$ , the algebra  $A(X)$  is the ring of real differential

<sup>2</sup>The latter fact can be viewed as a Hodge-Riemann relation in degree 0.

forms with constant coefficients on  $\mathbb{C}^d$ , and the cone  $K(X)$  is the spectrahedral cone of all  $d \times d$  positive definite matrices. Elementary proofs of the Hodge-Riemann relation for this  $X$  in any degree can be found in Gromov [1990] and Timorin [1998].

In the important special case when all the matrices are diagonal, the mixed discriminant is a permanent. Precisely, if  $A = (a_{ij})$  is an  $d \times d$  matrix and if  $A_j$  is the diagonal matrix whose  $j$ -th diagonal element is  $a_{jj}$ , then

$$d! D(A_1, \dots, A_d) = \text{per}(A) := \sum_{\sigma} \prod_{i=1}^d a_{i\sigma(i)},$$

where  $\sigma$  runs through all permutations of  $\{1, \dots, d\}$ . Therefore, for any column vectors  $a_1, \dots, a_d$  in  $\mathbb{R}^n$  with nonnegative entries,

$$\text{per}(a_1, a_2, a_3, \dots, a_d)^2 \geq \text{per}(a_1, a_1, a_3, \dots, a_d) \text{per}(a_2, a_2, a_3, \dots, a_d).$$

The above special case of the Hodge-Riemann relations for  $\mathbb{C}^d$  was the main ingredient in Egorychev’s and Falikman’s proofs of van der Waerden’s conjecture that the permanent of any doubly stochastic  $d \times d$  matrix is at least  $d!/d^d$ . See Knuth [1981] and van Lint [1982] for more on van der Waerden’s permanent conjecture.

**2.2 Mixed volumes of convex bodies.** The notion of mixed volume arises when one combines the volume with the Minkowski sum. For any collection of convex bodies  $\mathbf{P} = (P_1, \dots, P_d)$  in  $\mathbb{R}^d$ , consider the function

$$\text{vol}_{\mathbf{P}} : \mathbb{R}_{\geq 0}^d \longrightarrow \mathbb{R}_{\geq 0}, \quad (t_1, \dots, t_d) \longmapsto \text{vol}(t_1 P_1 + \dots + t_d P_d).$$

Minkowski noticed that  $\text{vol}_{\mathbf{P}}$  is a homogeneous polynomial of degree  $d$ , and called the number

$$V(P_1, \dots, P_d) = \frac{\partial^d}{\partial t_1 \dots \partial t_d} \text{vol}_{\mathbf{P}}(t_1, \dots, t_d)$$

the *mixed volume* of  $\mathbf{P}$ . The mixed volume is symmetric in  $\mathbf{P}$ , and it is nonnegative for any  $\mathbf{P}$ .<sup>3</sup>

Now let  $\eta_1, \dots, \eta_n$  be another collection of convex bodies in  $\mathbb{R}^d$ , and define an  $n \times n$  matrix  $AF = (AF_{ij})$  by

$$AF_{ij} = V(\eta_i, \eta_j, P_1, \dots, P_{d-2}).$$

If  $AF \neq 0$ , then the mixed volume analog of Theorem 1 holds.

**Theorem 2.** The matrix  $AF$  has exactly one positive eigenvalue.

It follows that the mixed volume satisfies the *Aleksandrov-Fenchel inequality*

$$\det \begin{pmatrix} V(P_1, P_1, P_3, \dots, P_d) & V(P_1, P_2, P_3, \dots, P_d) \\ V(P_1, P_2, P_3, \dots, P_d) & V(P_2, P_2, P_3, \dots, P_d) \end{pmatrix} \leq 0.$$

<sup>3</sup>The latter fact can be viewed as a Hodge-Riemann relation in degree 0.

In particular, the sequence of mixed volumes of two convex bodies is log-concave:

$$V(\underbrace{P_1, \dots, P_1}_i, \underbrace{P_2, \dots, P_2}_{d-i})^2 \geq V(\underbrace{P_1, \dots, P_1}_{i-1}, \underbrace{P_2, \dots, P_2}_{d-i+1}) V(\underbrace{P_1, \dots, P_1}_{i+1}, \underbrace{P_2, \dots, P_2}_{d-i-1}).$$

Aleksandrov reduced [Theorem 2](#) to the case when the Minkowski sum of all the relevant convex bodies, say  $\Delta$ , is a simple convex polytope. Under this hypothesis, [Theorem 2](#) is a Hodge-Riemann relation in degree 1 ([Gromov \[1990\]](#), [McMullen \[1993\]](#), and [Teissier \[1979\]](#)). The object  $X$  is the convex polytope  $\Delta$ , the algebra  $A(X)$  is McMullen's polytope algebra generated by the Minkowski summands of  $\Delta$ , and the cone  $K(X)$  is the cone of convex polytopes that share the normal fan with  $\Delta$ . Elementary proofs of the Hodge-Riemann relation for this  $X$  in any degree can be found in [Fleming and Karu \[2010\]](#), [McMullen \[1993\]](#), and [Timorin \[1999\]](#).

The Alexandrov-Fenchel inequality has been used to understand linear extensions of partially ordered sets. For example, [Chung, Fishburn, and Graham \[1980\]](#) conjectured that, for any finite poset  $Q$ ,

$$\Pr_i(x)^2 \geq \Pr_{i-1}(x)\Pr_{i+1}(x) \text{ for all } i \text{ and all } x \in Q,$$

where  $\Pr_i(x)$  is the fraction of linear extensions of  $Q$  in which  $x$  is the  $i$ -th largest element. [Stanley \[1981\]](#) proved the conjecture by constructing suitable convex polytopes from  $x \in Q$  and using the Alexandrov-Fenchel inequality. For another example, write  $\Pr(x_1 < x_2)$  for the fraction of linear extensions of  $Q$  in which  $x_1$  is smaller than  $x_2$ . [Kahn and Saks \[1984\]](#) employed Stanley's method to deduce the following remarkable fact from the Alexandrov-Fenchel inequality:

If  $Q$  is not a chain, then there are elements  $x_1, x_2 \in Q$  such that

$$3/11 < \Pr(x_1 < x_2) < 8/11.$$

This confirmed a conjecture of [Fredman \[1975/76\]](#) and [Linial \[1984\]](#) that the information theoretic lower bound for the general sorting problem is tight up to a multiplicative constant.

**2.3 The correlation constant of a field.** Let  $G$  be a finite connected graph, let  $i, j$  be distinct edges, and let  $T$  be a random spanning tree of  $G$ . Kirchhoff's effective resistance formula can be used to show that the probability that  $i$  is in  $T$  can only decrease by assuming that  $j$  is in  $T$ :

$$\Pr(i \in T) \geq \Pr(i \in T \mid j \in T).$$

In other words, the number  $b_-$  of spanning trees containing given edges satisfies

$$\frac{b_i}{b} \geq \frac{b_{ij}}{b_j}.$$

Now let  $M$  be a finite spanning subset of a vector space  $V$ , let  $i, j$  be distinct nonzero vectors in  $M$ , and write  $b_-$  for the number of bases in  $M$  containing given vectors. Do

we still have the negative correlation

$$\frac{b_i}{b} \geq \frac{b_{ij}}{b_j}?$$

The previous statement on graphs is the special case when  $M$  is the vertex-edge incidence matrix over the field with two elements.

[Seymour and Welsh \[1975\]](#) gave the first example of  $M$  over a field of characteristic 2 with  $\frac{b_i b_{ij}}{b_i b_j} = \frac{36}{35}$  for some  $i$  and  $j$ . How large can the ratio be?

**Definition 3.** The *correlation constant* of a field  $k$  is the supremum of  $\frac{b_i b_{ij}}{b_i b_j}$  over all pairs of distinct vectors  $i$  and  $j$  in finite vector configurations in vector spaces over  $k$ .

The correlation constant may be an interesting invariant of a field, although it is not immediately clear that the constant is finite. In fact, the finiteness of the correlation constant is one of the consequences of the Hodge-Riemann relations for vector configurations. Let  $n$  be the number of vectors in  $M$ , and let  $\text{HR}(M)$  be the symmetric  $n \times n$  matrix

$$\text{HR}(M)_{ij} = \begin{cases} 0 & \text{if } i = j, \\ b_{ij} & \text{if } i \neq j. \end{cases}$$

To avoid the trivial case  $\text{HR}(M) = 0$ , we suppose that the dimension of  $V$  is at least 2. For example, if  $K_4$  is the set of six column vectors of the matrix

$$\begin{pmatrix} 1 & 1 & 1 & 0 & 0 & 0 \\ -1 & 0 & 0 & 1 & 1 & 0 \\ 0 & -1 & 0 & -1 & 0 & 1 \\ 0 & 0 & -1 & 0 & -1 & -1 \end{pmatrix},$$

then  $\text{HR}(K_4)$  is the  $6 \times 6$  symmetric matrix

$$\begin{pmatrix} 0 & 3 & 3 & 3 & 3 & 4 \\ 3 & 0 & 3 & 3 & 4 & 3 \\ 3 & 3 & 0 & 4 & 3 & 3 \\ 3 & 3 & 4 & 0 & 3 & 3 \\ 3 & 4 & 3 & 3 & 0 & 3 \\ 4 & 3 & 3 & 3 & 3 & 0 \end{pmatrix}.$$

In [Huh and Wang \[2017\]](#), the following statement was deduced from [Theorem 12](#).

**Theorem 4.** The matrix  $\text{HR}(M)$  has exactly one positive eigenvalue.

In fact, the same statement holds more generally for any matroid  $M$  ([Huh and Wang \[ibid., Remark 15\]](#)). To deduce a bound on the correlation constant, consider the restriction of  $\text{HR}(M)$  to the three-dimensional subspace of  $\mathbb{R}^n$  spanned by  $\mathbf{e}_i$ ,  $\mathbf{e}_j$ , and  $(1, \dots, 1)$ . Cauchy’s eigenvalue interlacing theorem shows that the resulting  $3 \times 3$  symmetric matrix also has exactly one positive eigenvalue. Expressing the  $3 \times 3$  determinant, which should be nonnegative, we get the inequality

$$\frac{b_i b_{ij}}{b_i b_j} \leq 2 - 2(\dim V)^{-1}.$$

Thus the correlation constant of any field is at most 2. What is the correlation constant of, say,  $\mathbb{Z}/2\mathbb{Z}$ ? Does the correlation constant depend on the field?

**2.4 The chromatic polynomial of a graph.** Generalizing earlier work of Birkhoff, [Whitney \[1932\]](#) introduced the *chromatic polynomial* of a connected graph  $G$  as the function on  $\mathbb{N}$  defined by

$$\chi_G(q) = \text{the number of proper } q\text{-colorings of } G.$$

In other words,  $\chi_G(q)$  is the number of ways to color the vertices of  $G$  using  $q$  colors so that the endpoints of every edge have different colors. Whitney noticed that the chromatic polynomial is indeed a polynomial. In fact, we can write

$$\chi_G(q)/q = a_0(G)q^d - a_1(G)q^{d-1} + \cdots + (-1)^d a_d(G)$$

for some positive integers  $a_0(G), \dots, a_d(G)$ , where  $d$  is one less than the number of vertices.

*Example 5.* The cycle  $C_4$  with 4 vertices and 4 edges has the chromatic polynomial

$$\chi_{C_4}(q) = 1q^4 - 4q^3 + 6q^2 - 3q.$$


The chromatic polynomial was originally devised as a tool for attacking the Four Color Problem, but soon it attracted attention in its own right. [Read \[1968\]](#) conjectured that the coefficients of the chromatic polynomial form a unimodal sequence for any graph. A few years later, [Hoggar \[1974\]](#) conjectured more generally that the coefficients form a log-concave sequence:

$$a_i(G)^2 \geq a_{i-1}(G)a_{i+1}(G) \text{ for any } i \text{ and } G.$$

Notice that the chromatic polynomial can be computed using the *deletion-contraction relation*: if  $G \setminus e$  is the deletion of an edge  $e$  from  $G$  and  $G/e$  is the contraction of the same edge, then

$$\chi_G(q) = \chi_{G \setminus e}(q) - \chi_{G/e}(q).$$

The first term counts the proper colorings of  $G$ , the second term counts the otherwise-proper colorings of  $G$  where the endpoints of  $e$  are permitted to have the same color, and the third term counts the otherwise-proper colorings of  $G$  where the endpoints of  $e$  are mandated to have the same color. For example, to compute the chromatic polynomial of the cycle  $C_4$  in [Example 5](#), we write




and use that the chromatic polynomials of the two smaller graphs are  $q(q-1)^3$  and  $q(q-1)(q-2)$ , respectively. Note that, in general, the sum of log-concave sequences need not be log-concave.

The log-concavity conjecture for chromatic polynomials was proved in [Huh \[2012\]](#) by showing that the absolute values of the coefficients of  $\chi_G(q)/(q-1)$  are mixed

multiplicities of certain homogeneous ideals constructed from  $G$ . The notion of mixed multiplicities is a commutative algebraic analog of the notion of mixed volumes, and it can be shown that mixed multiplicities of homogeneous ideals satisfy a version of the Aleksandrov-Fenchel inequality. To formulate the underlying Hodge-Riemann relation in purely combinatorial terms was the primary motivation for [Adiprasito, Huh, and Katz \[2015\]](#). The main result of [Adiprasito, Huh, and Katz \[ibid.\]](#) will be reviewed in [Section 2.6](#) below.

**2.5 Counting independent sets.** How many linearly independent collection of  $i$  vectors are there in a given configuration of vectors? Let's write  $M$  for a finite subset of a vector space and  $f_i(M)$  for the number of independent subsets of  $M$  of size  $i$ .

*Example 6.* Let  $F$  be the set of all nonzero vectors in the three-dimensional vector space over the field with two elements. Nontrivial dependencies between elements of  $F$  can be read off from the picture of the Fano plane shown below.



The nonempty independent subsets of  $F$  correspond to the seven points in  $F$ , the twenty-one pairs of points in  $F$ , and the twenty-eight triple of points in  $F$  not in one of the seven lines:

$$f_0(F) = 1, \quad f_1(F) = 7, \quad f_2(F) = 21, \quad f_3(F) = 28.$$

[Welsh \[1971\]](#) conjectured that the sequence  $f_i(M)$  is unimodal for any  $M$ . Shortly after, [Mason \[1972\]](#) conjectured more generally that the sequence is log-concave:

$$f_i(M)^2 \geq f_{i-1}(M)f_{i+1}(M) \text{ for any } i \text{ and } M.$$

In any small specific case, the conjecture can be verified by computing the  $f_i(M)$ 's by the *deletion-contraction relation*: if  $M \setminus v$  is the deletion of a nonzero vector  $v$  from  $M$  and  $M/v$  is the projection of  $M$  in the direction of  $v$ , then

$$f_i(M) = f_i(M \setminus v) + f_{i-1}(M/v).$$

The first term counts the number of independent subsets of size  $i$ , the second term counts the independent subsets of size  $i$  not containing  $v$ , and the third term counts the independent subsets of size  $i$  containing  $v$ . As in the case of graphs, we notice the apparent conflict between the log-concavity conjecture and the additive nature of  $f_i(M)$ .

The log-concavity conjecture for  $f_i(M)$  was proved in [Lenz \[2013\]](#) by combining a geometric construction of [Huh and Katz \[2012\]](#) and a matroid-theoretic construction of [Brylawski \[1977\]](#). Given a spanning subset  $M$  of a  $d$ -dimensional vector space over a

field  $k$ , one can construct a  $d$ -dimensional smooth projective variety  $X(\mathbf{M})$  over  $k$  and globally generated line bundles  $L_1, L_2$  on  $X(\mathbf{M})$  so that

$$f_i(\mathbf{M}) = \int_{X(\mathbf{M})} L_1^{d-i} L_2^i.$$

The Hodge-Riemann relation for smooth projective varieties is known to hold in degrees  $q \leq 1$  (Grothendieck [1958] and Segre [1937]), and this implies the log-concavity of  $f_i(\mathbf{M})$  as in Sections 2.1, 2.2. To express and verify the general Hodge-Riemann relation for  $X(\mathbf{M})$  in purely combinatorial terms was another motivation for Adiprasito, Huh, and Katz [2015].

**2.6 The Hodge-Riemann relations for matroids.** In the 1930s, Hassler Whitney observed that several notions in graph theory and linear algebra fit together in a common framework, that of *matroids* (Whitney [1935]). This observation started a new subject with applications to a wide range of topics like characteristic classes, optimization, and moduli spaces.

**Definition 7.** A *matroid*  $\mathbf{M}$  on a finite set  $E$  is a collection of subsets of  $E$ , called *flats* of  $\mathbf{M}$ , satisfying the following axioms:

- (1) The ground set  $E$  is a flat.
- (2) If  $F_1$  and  $F_2$  are flats, then  $F_1 \cap F_2$  is a flat.
- (3) If  $F$  is a flat, then any element not in  $F$  is contained in exactly one flat covering  $F$ .

Here, a flat is said to *cover* another flat  $F$  if it is minimal among the flats properly containing  $F$ .

For our purposes, we may and will suppose that  $\mathbf{M}$  is *loopless*:

- (4) The empty subset of  $E$  is a flat.

Every maximal chain of flats in  $F$  has the same length, and this common length is called the *rank* of the flat  $F$ . The rank of the flat  $E$  is called the rank of the matroid  $\mathbf{M}$ . Matroids are determined by their *independent sets* (the idea of “*general position*”), and can be alternatively defined in terms of independent sets (Oxley [2011, Chapter 1]).

*Example 8.* Let  $E$  be the set of edges of a finite graph  $G$ . Call a subset  $F$  of  $E$  a flat when there is no edge in  $E \setminus F$  whose endpoints are connected by a path in  $F$ . This defines a *graphic matroid* on  $E$ .

*Example 9.* A *projective space*  $\mathbb{P}$  is a set with distinguished subsets, called *lines*, satisfying:

- (1) Every line contains more than two points.
- (2) If  $x, y$  are distinct points, then there is exactly one line  $xy$  containing  $x$  and  $y$ .

(3) If  $x, y, z, w$  are distinct points, no three collinear, then


the line  $xy$  intersects the line  $zw \implies$  the line  $xz$  intersects the line  $yw$ .

A subspace of  $\mathbb{P}^2$  is a subset  $\mathbb{S}$  of  $\mathbb{P}^2$  such that

$x$  and  $y$  are distinct points in  $\mathbb{S} \implies$  the line  $xy$  is in  $\mathbb{S}$ .

For any finite subset  $E$  of  $\mathbb{P}^2$ , the collection of sets of the form  $E \cap \mathbb{S}$  has the structure of a matroid. Matroids arising from subsets of projective spaces over a field  $k$  are said to be *realizable* over  $k$  (the idea of “coordinates”).

Not surprisingly, the notion of realizability is sensitive to the field  $k$ . A matroid may arise from a vector configuration over one field while no such vector configuration exists over another field.



Among the rank 3 matroids pictured above, where rank 1 flats are represented by points and rank 2 flats containing more than 2 points are represented by lines, the first is realizable over  $k$  if and only if the characteristic of  $k$  is 2, the second is realizable over  $k$  if and only if the characteristic of  $k$  is not 2, and the third is not realizable over any field. Recently, Nelson [2016] showed that almost all matroids are not realizable over any field Nelson [ibid.].

**Definition 10.** We introduce variables  $x_F$ , one for each nonempty proper flat  $F$  of  $M$ , and consider the polynomial ring

$$S(M) = \mathbb{R}[x_F]_{F \neq \emptyset, F \neq E}.$$

The *Chow ring*  $A(M)$  of  $M$  is the quotient of  $S(M)$  by the ideal generated by the linear forms

$$\sum_{i_1 \in F} x_{F_i_1} - \sum_{i_2 \in F} x_{F_i_2},$$

one for each pair of distinct elements  $i_1$  and  $i_2$  of  $E$ , and the quadratic monomials

$$x_{F_1} x_{F_2},$$

one for each pair of incomparable nonempty proper flats  $F_1$  and  $F_2$  of  $M$ . We have

$$A(M) = \bigoplus_q A^q(M),$$

where  $A^q(M)$  is the span of degree  $q$  monomials in  $A(M)$ .

Feichtner and Yuzvinsky introduced the Chow ring of  $M$  (Feichtner and Yuzvinsky [2004]). When  $M$  is realizable over a field  $k$ , it is the Chow ring of the “wonderful” compactification of the complement of a hyperplane arrangement defined over  $k$  studied by De Concini and Procesi [1995].

To formulate the hard Lefschetz theorem and the Hodge-Riemann relations for  $A(M)$ , we define a matroid analog of the Kähler cone in complex geometry.

**Definition 11.** A real-valued function  $c$  on  $2^E$  is said to be *strictly submodular* if

$$c_{\emptyset} = 0, \quad c_E = 0,$$

and, for any two incomparable subsets  $I_1, I_2 \subseteq E$ ,

$$c_{I_1} + c_{I_2} > c_{I_1 \cap I_2} + c_{I_1 \cup I_2}.$$

We note that strictly submodular functions exist. For example,

$$I \mapsto |I||E \setminus I|$$

is a strictly submodular function. A strictly submodular function  $c$  defines an element

$$L_c = \sum_F c_F x_F \in A^1(M).$$

The Kähler cone  $K(M)$  is defined to be the set of all such elements in  $A^1(M)$ .

Now let  $d + 1$  be the rank of  $M$ , and write “deg” for the unique linear isomorphism

$$\text{deg}: A^d(M) \longrightarrow \mathbb{R}$$

which maps  $x_{F_1} \cdots x_{F_d}$  to 1 for every maximal chain  $F_1 \subsetneq \cdots \subsetneq F_d$  of nonempty proper flats (Adiprasito, Huh, and Katz [2015, Proposition 5.10]). We are ready to state the hard Lefschetz theorem and the Hodge-Riemann relation for  $M$  (Adiprasito, Huh, and Katz [ibid., Theorem 8.9]).

**Theorem 12.** Let  $q$  be a nonnegative integer  $\leq \frac{d}{2}$ , and let  $L_0, L_1, \dots, L_{d-2q} \in K(M)$ .

(PD) The product in  $A(M)$  defines a nondegenerate bilinear pairing

$$A^q(M) \times A^{d-q}(M) \longrightarrow \mathbb{R}, \quad (\eta, \xi) \longmapsto \text{deg}(\eta \xi).$$

(HL) The multiplication by  $L_1, \dots, L_{d-2q}$  defines a linear bijection

$$A^q(M) \longrightarrow A^{d-q}(M), \quad \eta \longmapsto \left( \prod_{i=1}^{d-2q} L_i \right) \eta.$$

(HR) The symmetric bilinear form

$$A^q(M) \times A^q(M) \longrightarrow \mathbb{R}, \quad (\eta_1, \eta_2) \longmapsto (-1)^q \text{deg}\left(\left(\prod_{i=1}^{d-2q} L_i\right) \eta_1 \eta_2\right)$$

is positive definite on the kernel of the multiplication map

$$A^q(\mathbf{M}) \longrightarrow A^{d-q+1}(\mathbf{M}), \quad \eta \longmapsto \left( \prod_{i=0}^{d-2q} L_i \right) \eta.$$

We highlight the following consequence of HR in degrees  $\leq 1$ : For any  $\xi_1, \xi_2 \in K(\mathbf{M})$ ,

$$\begin{pmatrix} \deg\left(\left(\prod_{i=1}^{d-2} L_i\right) \xi_1 \xi_1\right) & \deg\left(\left(\prod_{i=1}^{d-2} L_i\right) \xi_1 \xi_2\right) \\ \deg\left(\left(\prod_{i=1}^{d-2} L_i\right) \xi_1 \xi_2\right) & \deg\left(\left(\prod_{i=1}^{d-2} L_i\right) \xi_2 \xi_2\right) \end{pmatrix}$$

has exactly one positive eigenvalue. Taking the determinant, we get an analog of the Alexandrov-Fenchel inequality

$$\deg\left(\left(\prod_{i=1}^{d-2} L_i\right) \xi_1 \xi_2\right)^2 \geq \deg\left(\left(\prod_{i=1}^{d-2} L_i\right) \xi_1 \xi_1\right) \deg\left(\left(\prod_{i=1}^{d-2} L_i\right) \xi_2 \xi_2\right).$$

We apply the inequality to the *characteristic polynomial*  $\chi_{\mathbf{M}}(q)$ , a generalization of the chromatic polynomial  $\chi_G(q)$  to a matroid  $\mathbf{M}$  that is not necessarily graphic (Welsh [1976, Chapter 15]). For this, we consider two distinguished elements of  $A^1(\mathbf{M})$ . For fixed  $j \in E$ , the elements are

$$\alpha = \sum_{j \in F} x_F, \quad \beta = \sum_{j \notin F} x_F.$$

The two elements do not depend on the choice of  $j$ , and they are limits of elements of the form  $\ell_c$  for a strictly submodular function  $c$ . A bijective counting argument in Adiprasito, Huh, and Katz [2015] shows that

$$e_i(\mathbf{M}) = \deg(\alpha^i \beta^{d-i}) \text{ for every } i,$$

where  $e_i(\mathbf{M})$  is the sequence of integers satisfying the identity

$$\chi_{\mathbf{M}}(q)/(q-1) = e_0(\mathbf{M})q^d - e_1(\mathbf{M})q^{d-1} + \dots + (-1)^d e_d(\mathbf{M}).$$

Thus the sequence  $e_i(\mathbf{M})$  is log-concave, which implies the following conjecture of Heron [1972], Rota [1971], and Welsh [1976]:

The coefficients of  $\chi_{\mathbf{M}}(q)$  form a log-concave sequence for any matroid  $\mathbf{M}$ .

The above implies the log-concavity of the sequence  $a_i(G)$  in Section 2.4 and the log-concavity of the sequence  $f_i(\mathbf{M})$  in Section 2.5. See Oxley [2011, Chapter 15] and White [1987, Chapter 8] for overviews and historical accounts.

**2.7 The reliability polynomial of a network.** Let  $G$  be a finite connected graph with  $v$  vertices and  $n$  edges. The *reliability* of  $G$  is the probability that any two vertices remain connected when each edge is independently removed with the same probability  $q$ . Let's write  $o_i(G)$  for the number of  $i$ -edge operational states. For example,  $o_{v-1}(G)$  is the number of spanning trees and  $o_{n-1}(G)$  is the number of non-bridges. Thus the reliability of  $G$  is

$$R_G(q) = \sum_i o_i(G)(1-q)^i q^{n-i}.$$

We define a sequence of integers  $h_0(G), \dots, h_d(G)$  by the identity

$$R_G(q)/(1-q)^{v-1} = h_d(G)q^d + h_{d-1}(G)q^{d-1} + \dots + h_0(G),$$

where  $d$  is one more than the difference  $n - v$ .

*Example 13.* The complete graph on 4 vertices has the reliability polynomial

$$\begin{aligned} R_{K_4}(q) &= 16q^3(1-q)^3 + 15q^2(1-q)^4 + 6q(1-q)^5 + 1(1-q)^6 \\ &= (1-q)^3(6q^3 + 6q^2 + 3q + 1). \end{aligned}$$

The numbers  $h_i$  are closely related to the numbers  $f_i$  of independent sets in [Section 2.5](#). Writing  $M$  for the dual of the graphic matroid of  $G$ , we have

$$\sum_{i=0}^d h_i(G)x^i = \sum_{i=0}^d f_i(M)x^i(1-x)^{d-i} = \sum_{i=0}^d h_i(M)x^i.$$

[Dawson \[1984\]](#) conjectured that the sequence  $h_i(M)$  defined by the second equality is log-concave for any matroid  $M$ :

$$h_i(M)^2 \geq h_{i-1}(M)h_{i+1}(M) \text{ for any } i \text{ and } M.$$

[Colbourn \[1987\]](#) independently conjectured the same in the context of reliability polynomials.

When  $M$  is the dual of a graphic matroid, or more generally when  $M$  is realizable over the complex numbers, the log-concavity conjecture for  $h_i(M)$  was proved in [Huh \[2015\]](#) by applying an algebraic analog of the Alexandrov-Fenchel inequality to the variety of critical points of the master function of a realization of  $M$  studied by [Denham, Garrounian, and Schulze \[2012\]](#). The underlying combinatorial Hodge-Riemann relation is yet to be formulated, and Dawson's conjecture for general matroids remains open. The argument in the complex realizable case is tightly connected to the geometry of characteristic cycles ([Huh \[2013\]](#)), suggesting that the combinatorial Hodge-Riemann relation in this context will be strictly stronger than that of [Section 2.6](#).

**2.8 Unsolved problems.** The log-concavity of a sequence is not only important because of its applications but because it hints the existence of a structure that satisfies PD, HL, and HR. We close by listing some of the most interesting sequences that are conjectured to be log-concave.

- (1) Rota’s unimodality conjecture (Rota [1971]): If  $w_k(M)$  is the number of rank  $k$  flats of a rank  $d$  matroid  $M$ , then the sequence  $w_0(M), \dots, w_d(M)$  is unimodal. Welsh [1976] conjectured more generally that the sequence is log-concave.
- (2) Fox’s trapezoidal conjecture (Fox [1962]): The sequence of absolute values of the coefficients of the Alexander polynomial of an alternating knot strictly increases, possibly plateaus, then strictly decreases. Stoimenow [2005] conjectured more generally that the sequence is log-concave.
- (3) Kazhdan-Lusztig polynomials of matroids (Elias, Proudfoot, and Wakefield [2016]): For any matroid  $M$ , the coefficients of the Kazhdan-Lusztig polynomial of  $M$  form a nonnegative log-concave sequence.

## References

- Karim Adiprasito, June Huh, and Eric Katz (Nov. 2015). “Hodge Theory for Combinatorial Geometries”. arXiv: [1511.02888](#) (cit. on pp. [3087](#), [3088](#), [3090](#), [3091](#)).
- Alexander D Aleksandrov (1938). “Zur Theorie der gemischten Volumina von konvexen Körpern, IV. Die gemischten Diskriminanten und die gemischten Volumina”. *Mat. Sbornik* 3.45, pp. 227–251 (cit. on p. [3082](#)).
- Louis J. Billera and Carl W. Lee (1980). “Sufficiency of McMullen’s conditions for  $f$ -vectors of simplicial polytopes”. *Bull. Amer. Math. Soc. (N.S.)* 2.1, pp. 181–185. MR: [551759](#) (cit. on p. [3081](#)).
- Francesco Brenti (1994). “Log-concave and unimodal sequences in algebra, combinatorics, and geometry: an update”. In: *Jerusalem combinatorics ’93*. Vol. 178. Contemp. Math. Amer. Math. Soc., Providence, RI, pp. 71–89. MR: [1310575](#) (cit. on p. [3079](#)).
- Tom Brylawski (1977). “The broken-circuit complex”. *Trans. Amer. Math. Soc.* 234.2, pp. 417–433. MR: [468931](#) (cit. on p. [3087](#)).
- F. R. K. Chung, P. C. Fishburn, and R. L. Graham (1980). “On unimodality for linear extensions of partial orders”. *SIAM J. Algebraic Discrete Methods* 1.4, pp. 405–410. MR: [593850](#) (cit. on p. [3084](#)).
- Charles J. Colbourn (1987). *The combinatorics of network reliability*. International Series of Monographs on Computer Science. The Clarendon Press, Oxford University Press, New York, pp. xii+160. MR: [902584](#) (cit. on p. [3092](#)).
- Pierre Colmez and Jean-Pierre Serre, eds. (2001). *Correspondance Grothendieck-Serre*. Vol. 2. Documents Mathématiques (Paris) [Mathematical Documents (Paris)]. Société Mathématique de France, Paris, pp. xii+288. MR: [1942134](#) (cit. on p. [3082](#)).
- Jeremy E. Dawson (1984). “A collection of sets related to the Tutte polynomial of a matroid”. In: *Graph theory, Singapore 1983*. Vol. 1073. Lecture Notes in Math. Springer, Berlin, pp. 193–204. MR: [761018](#) (cit. on p. [3092](#)).
- C. De Concini and C. Procesi (1995). “Wonderful models of subspace arrangements”. *Selecta Math. (N.S.)* 1.3, pp. 459–494. MR: [1366622](#) (cit. on p. [3090](#)).

- Graham Denham, Mehdi Garrounian, and Mathias Schulze (2012). “[A geometric deletion-restriction formula](#)”. *Adv. Math.* 230.4–6, pp. 1979–1994. MR: [2927361](#) (cit. on p. [3092](#)).
- Sudhakar Dharmadhikari and Kumar Joag-Dev (1988). *Unimodality, convexity, and applications*. Probability and Mathematical Statistics. Academic Press, Inc., Boston, MA, pp. xiv+278. MR: [954608](#) (cit. on p. [3079](#)).
- Thomas A. Dowling and Richard M. Wilson (1974). “[The slimmest geometric lattices](#)”. *Trans. Amer. Math. Soc.* 196, pp. 203–215. MR: [0345849](#) (cit. on p. [3081](#)).
- (1975). “[Whitney number inequalities for geometric lattices](#)”. *Proc. Amer. Math. Soc.* 47, pp. 504–512. MR: [0354422](#) (cit. on p. [3081](#)).
- Ben Elias, Nicholas Proudfoot, and Max Wakefield (2016). “[The Kazhdan-Lusztig polynomial of a matroid](#)”. *Adv. Math.* 299, pp. 36–70. MR: [3519463](#) (cit. on p. [3093](#)).
- Ben Elias and Geordie Williamson (2014). “[The Hodge theory of Soergel bimodules](#)”. *Ann. of Math. (2)* 180.3, pp. 1089–1136. MR: [3245013](#) (cit. on p. [3080](#)).
- P. Erdős (1965). “Extremal problems in number theory”. In: *Proc. Sympos. Pure Math., Vol. VIII*. Amer. Math. Soc., Providence, R.I., pp. 181–189. MR: [0174539](#) (cit. on p. [3081](#)).
- Eva Maria Feichtner and Sergey Yuzvinsky (2004). “[Chow rings of toric varieties defined by atomic lattices](#)”. *Invent. Math.* 155.3, pp. 515–536. MR: [2038195](#) (cit. on p. [3090](#)).
- Balin Fleming and Kalle Karu (2010). “[Hard Lefschetz theorem for simple polytopes](#)”. *J. Algebraic Combin.* 32.2, pp. 227–239. MR: [2661416](#) (cit. on p. [3084](#)).
- R. H. Fox (1962). “Some problems in knot theory”. In: *Topology of 3-manifolds and related topics (Proc. The Univ. of Georgia Institute, 1961)*. Prentice-Hall, Englewood Cliffs, N.J., pp. 168–176. MR: [0140100](#) (cit. on p. [3093](#)).
- Michael L. Fredman (1975/76). “[How good is the information theory bound in sorting?](#)” *Theoret. Comput. Sci.* 1.4, pp. 355–361. MR: [0416100](#) (cit. on p. [3084](#)).
- M. Gromov (1990). “Convex sets and Kähler manifolds”. In: *Advances in differential geometry and topology*. World Sci. Publ., Teaneck, NJ, pp. 1–38. MR: [1095529](#) (cit. on pp. [3080](#), [3083](#), [3084](#)).
- A. Grothendieck (1958). “[Sur une note de Mattuck-Tate](#)”. *J. Reine Angew. Math.* 200, pp. 208–215. MR: [0136607](#) (cit. on p. [3088](#)).
- (1969). “Standard conjectures on algebraic cycles”. In: *Algebraic Geometry (Internat. Colloq., Tata Inst. Fund. Res., Bombay, 1968)*. Oxford Univ. Press, London, pp. 193–199. MR: [0268189](#) (cit. on p. [3080](#)).
- A. P. Heron (1972). “Matroid polynomials”, pp. 164–202. MR: [0340058](#) (cit. on p. [3091](#)).
- S. G. Hoggar (1974). “Chromatic polynomials and logarithmic concavity”. *J. Combinatorial Theory Ser. B* 16, pp. 248–254. MR: [0342424](#) (cit. on p. [3086](#)).
- June Huh (2012). “[Milnor numbers of projective hypersurfaces and the chromatic polynomial of graphs](#)”. *J. Amer. Math. Soc.* 25.3, pp. 907–927. MR: [2904577](#) (cit. on p. [3086](#)).
- (2013). “[The maximum likelihood degree of a very affine variety](#)”. *Compos. Math.* 149.8, pp. 1245–1266. MR: [3103064](#) (cit. on p. [3092](#)).

- (2015). “*[h](#)-vectors of matroids and logarithmic concavity*”. *Adv. Math.* 270, pp. 49–59. MR: [3286530](#) (cit. on p. [3092](#)).
- June Huh and Eric Katz (2012). “*Log-concavity of characteristic polynomials and the Bergman fan of matroids*”. *Math. Ann.* 354.3, pp. 1103–1116. MR: [2983081](#) (cit. on p. [3087](#)).
- June Huh and Botong Wang (2017). “*Enumeration of points, lines, planes, etc.*”. *Acta Math.* 218.2, pp. 297–317. MR: [3733101](#) (cit. on pp. [3081](#), [3085](#)).
- Jeff Kahn and Michael Saks (1984). “*Balancing poset extensions*”. *Order* 1.2, pp. 113–126. MR: [764319](#) (cit. on p. [3084](#)).
- Kalle Karu (2004). “*Hard Lefschetz theorem for nonrational polytopes*”. *Invent. Math.* 157.2, pp. 419–447. MR: [2076929](#) (cit. on p. [3080](#)).
- S. L. Kleiman (1968). “*Algebraic cycles and the Weil conjectures*”. In: *Dix exposés sur la cohomologie des schémas*. Vol. 3. Adv. Stud. Pure Math. North-Holland, Amsterdam, pp. 359–386. MR: [292838](#) (cit. on p. [3082](#)).
- Steven L. Kleiman (1994). “*The standard conjectures*”. In: *Motives (Seattle, WA, 1991)*. Vol. 55. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, RI, pp. 3–20. MR: [1265519](#) (cit. on p. [3082](#)).
- Donald E. Knuth (1981). “*A permanent inequality*”. *Amer. Math. Monthly* 88.10, pp. 731–740, 798. MR: [668399](#) (cit. on p. [3083](#)).
- Matthias Lenz (2013). “*The  $f$ -vector of a representable-matroid complex is log-concave*”. *Adv. in Appl. Math.* 51.5, pp. 543–545. MR: [3118543](#) (cit. on p. [3087](#)).
- Nathan Linial (1984). “*The information-theoretic bound is good for merging*”. *SIAM J. Comput.* 13.4, pp. 795–801. MR: [764179](#) (cit. on p. [3084](#)).
- J. H. van Lint (1982). “*The van der Waerden conjecture: two proofs in one year*”. *Math. Intelligencer* 4.2, pp. 72–77. MR: [672919](#) (cit. on p. [3083](#)).
- Albert W. Marshall, Ingram Olkin, and Barry C. Arnold (2011). *Inequalities: theory of majorization and its applications*. Second. Springer Series in Statistics. Springer, New York, pp. xxviii+909. MR: [2759813](#) (cit. on p. [3079](#)).
- J. H. Mason (1972). “*Matroids: unimodal conjectures and Motzkin’s theorem*”, pp. 207–220. MR: [0349445](#) (cit. on p. [3087](#)).
- P. McMullen (1971). “*The numbers of faces of simplicial polytopes*”. *Israel J. Math.* 9, pp. 559–570. MR: [0278183](#) (cit. on p. [3081](#)).
- Peter McMullen (1993). “*On simple polytopes*”. *Invent. Math.* 113.2, pp. 419–444. MR: [1228132](#) (cit. on pp. [3080](#), [3084](#)).
- Peter Nelson (May 2016). “*Almost all matroids are non-representable*”. arXiv: [1605.04288](#) (cit. on p. [3089](#)).
- Andrei Okounkov (2003). “*Why would multiplicities be log-concave?*” In: *The orbit method in geometry and physics (Marseille, 2000)*. Vol. 213. Progr. Math. Birkhäuser Boston, Boston, MA, pp. 329–347. MR: [1995384](#) (cit. on p. [3079](#)).
- James Oxley (2011). *Matroid theory*. Second. Vol. 21. Oxford Graduate Texts in Mathematics. Oxford University Press, Oxford, pp. xiv+684. MR: [2849819](#) (cit. on pp. [3088](#), [3091](#)).

- Ronald C. Read (1968). “An introduction to chromatic polynomials”. *J. Combinatorial Theory* 4, pp. 52–71. MR: [0224505](#) (cit. on p. [3086](#)).
- Gian-Carlo Rota (1971). “Combinatorial theory, old and new”, pp. 229–233. MR: [0505646](#) (cit. on pp. [3091](#), [3093](#)).
- Adrien Saumard and Jon A. Wellner (2014). “Log-concavity and strong log-concavity: a review”. *Stat. Surv.* 8, pp. 45–114. MR: [3290441](#) (cit. on p. [3079](#)).
- Beniamino Segre (1937). “Intorno ad un teorema di Hodge sulla teoria della base per le curve di una superficie algebrica”. *Ann. Mat. Pura Appl.* 16.1, pp. 157–163. MR: [1553294](#) (cit. on p. [3088](#)).
- P. D. Seymour and D. J. A. Welsh (1975). “Combinatorial applications of an inequality from statistical mechanics”. *Math. Proc. Cambridge Philos. Soc.* 77, pp. 485–495. MR: [0376378](#) (cit. on p. [3085](#)).
- Richard P. Stanley (1980a). “The number of faces of a simplicial convex polytope”. *Adv. in Math.* 35.3, pp. 236–238. MR: [563925](#) (cit. on p. [3081](#)).
- (1980b). “Weyl groups, the hard Lefschetz theorem, and the Sperner property”. *SIAM J. Algebraic Discrete Methods* 1.2, pp. 168–184. MR: [578321](#) (cit. on p. [3081](#)).
- (1981). “Two combinatorial applications of the Aleksandrov-Fenchel inequalities”. *J. Combin. Theory Ser. A* 31.1, pp. 56–65. MR: [626441](#) (cit. on p. [3084](#)).
- (1984). “Combinatorial applications of the hard Lefschetz theorem”. In: *Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Warsaw, 1983)*. PWN, Warsaw, pp. 447–453. MR: [804700](#) (cit. on p. [3081](#)).
- (1989). “Log-concave and unimodal sequences in algebra, combinatorics, and geometry”. In: *Graph theory and its applications: East and West (Jinan, 1986)*. Vol. 576. Ann. New York Acad. Sci. New York Acad. Sci., New York, pp. 500–535. MR: [1110850](#) (cit. on p. [3079](#)).
- (2000). “Positivity problems and conjectures in algebraic combinatorics”. In: *Mathematics: frontiers and perspectives*. Amer. Math. Soc., Providence, RI, pp. 295–319. MR: [1754784](#) (cit. on p. [3079](#)).
- Alexander Stoimenow (2005). “Newton-like polynomials of links”. *Enseign. Math.* (2) 51.3-4, pp. 211–230. MR: [2214886](#) (cit. on p. [3093](#)).
- Bernard Teissier (1979). “Du théorème de l’index de Hodge aux inégalités isopérimétriques”. *C. R. Acad. Sci. Paris Sér. A-B* 288.4, A287–A289. MR: [524795](#) (cit. on p. [3084](#)).
- V. A. Timorin (1998). “Mixed Hodge-Riemann bilinear relations in a linear context”. *Funktsional. Anal. i Prilozhen.* 32.4, pp. 63–68, 96. MR: [1678857](#) (cit. on p. [3083](#)).
- (1999). “An analogue of the Hodge-Riemann relations for simple convex polyhedra”. *Uspekhi Mat. Nauk* 54.2(326), pp. 113–162. MR: [1711255](#) (cit. on p. [3084](#)).
- D. J. A. Welsh (1971). “Combinatorial problems in matroid theory”. In: *Combinatorial Mathematics and its Applications (Proc. Conf., Oxford, 1969)*. Academic Press, London, pp. 291–306. MR: [0278975](#) (cit. on p. [3087](#)).
- (1976). *Matroid theory*. L. M. S. Monographs, No. 8. Academic Press [Harcourt Brace Jovanovich, Publishers], London-New York, pp. xi+433. MR: [0427112](#) (cit. on pp. [3091](#), [3093](#)).

Neil White, ed. (1987). *Combinatorial geometries*. Vol. 29. Encyclopedia of Mathematics and its Applications. Cambridge University Press, Cambridge, pp. xii+212. MR: [921064](#) (cit. on p. [3091](#)).

Hassler Whitney (1932). “A logical expansion in mathematics”. *Bull. Amer. Math. Soc.* 38.8, pp. 572–579. MR: [1562461](#) (cit. on p. [3086](#)).

– (1935). “On the Abstract Properties of Linear Dependence”. *Amer. J. Math.* 57.3, pp. 509–533. MR: [1507091](#) (cit. on p. [3088](#)).

Received 2017-11-29.

June Huh  
Institute for Advanced Study  
Fuld Hall  
1 Einstein Drive  
Princeton, NJ  
USA  
[huh@princeton.edu](mailto:huh@princeton.edu)  
[junehuh@ias.edu](mailto:junehuh@ias.edu)

