

The Hilbert–Smith conjecture for three-manifolds

John Pardon

9 April 2012; Revised 22 January 2013

Abstract

We show that every locally compact group which acts faithfully on a connected three-manifold is a Lie group. By known reductions, it suffices to show that there is no faithful action of \mathbb{Z}_p (the p -adic integers) on a connected three-manifold. If \mathbb{Z}_p acts faithfully on M^3 , we find an interesting \mathbb{Z}_p -invariant open set $U \subseteq M$ with $H_2(U) = \mathbb{Z}$ and analyze the incompressible surfaces in U representing a generator of $H_2(U)$. It turns out that there must be one such incompressible surface, say F , whose isotopy class is fixed by \mathbb{Z}_p . An analysis of the resulting homomorphism $\mathbb{Z}_p \rightarrow \text{MCG}(F)$ gives the desired contradiction. The approach is local on M .

MSC 2010 Primary: 57S10, 57M60, 20F34, 57S05, 57N10

MSC 2010 Secondary: 54H15, 55M35, 57S17

Keywords: transformation groups, Hilbert–Smith conjecture, Hilbert’s Fifth Problem, three-manifolds, incompressible surfaces

1 Introduction

By a *faithful action* of a topological group G on a topological manifold M , we mean a continuous injection $G \rightarrow \text{Homeo}(M)$ (where $\text{Homeo}(M)$ has the compact-open topology). A well-known problem is to characterize the locally compact topological groups which can act faithfully on a manifold. In particular, there is the following conjecture, which is a natural generalization of Hilbert’s Fifth Problem.

Conjecture 1.1 (Hilbert–Smith). *If a locally compact topological group G acts faithfully on some connected n -manifold M , then G is a Lie group.*

It is well-known (see, for example, Lee [17] or Tao [41]) that as a consequence of the solution of Hilbert’s Fifth Problem (by Gleason [8, 9], Montgomery–Zippin [22, 23], as well as further work by Yamabe [46, 47]), any counterexample G to Conjecture 1.1 must contain an embedded copy of \mathbb{Z}_p (the p -adic integers). Thus it is equivalent to consider the following conjecture.

Conjecture 1.2. *There is no faithful action of \mathbb{Z}_p on any connected n -manifold M .*

Conjecture 1.1 also admits the following reformulation, which we hope will help the reader better understand its flavor.

Conjecture 1.3. *Given a connected n -manifold M with metric d and open set $U \subseteq M$, there exists $\epsilon > 0$ such that the subset:*

$$\{\phi \in \text{Homeo}(M) \mid d(x, \phi(x)) < \epsilon \text{ for all } x \in U\} \quad (1.1)$$

of $\text{Homeo}(M)$ contains no nontrivial compact subgroup.

For any specific manifold M , Conjectures 1.1–1.3 for M are equivalent. They also have the following simple consequence for *almost periodic* homeomorphisms of M (a homeomorphism is said to be almost periodic if and only if the subgroup of $\text{Homeo}(M)$ it generates has compact closure; see Gottschalk [10] for other equivalent definitions).

Conjecture 1.4. *For every almost periodic homeomorphism f of a connected n -manifold M , there exists $r > 0$ such that f^r is in the image of some homomorphism $\mathbb{R} \rightarrow \text{Homeo}(M)$.*

Conjecture 1.1 is known in the cases $n = 1, 2$ (see Montgomery–Zippin [23, pp233,249]). By consideration of $M \times \mathbb{R}$, clearly Conjecture 1.1 in dimension n implies the same in all lower dimensions.

The most popular approach to the conjectures above is via Conjecture 1.2. Yang [48] showed that for any counterexample to Conjecture 1.2, the orbit space M/\mathbb{Z}_p must have cohomological dimension $n + 2$. Conjecture 1.2 has been established for various regularity classes of actions, C^2 actions by Bochner–Montgomery [4], $C^{0,1}$ actions by Repovš–Ščepin [32], $C^{0, \frac{n}{n+2} + \epsilon}$ actions by Maleshich [18], quasiconformal actions by Martin [19], and uniformly quasisymmetric actions on doubling Ahlfors regular compact metric measure manifolds with Hausdorff dimension in $[1, n + 2)$ by Mj [20]. In the negative direction, it is known by work of Walsh [44, p282 Corollary 5.15.1] that there *does* exist a continuous decomposition of any compact PL n -manifold into cantor sets of arbitrarily small diameter if $n \geq 3$ (see also Wilson [45, Theorem 3]). By work of Raymond–Williams [31], there are faithful actions of \mathbb{Z}_p on n -dimensional compact metric spaces which achieve the cohomological dimension jump of Yang [48] for every $n \geq 2$.

In this paper, we establish the aforementioned conjectures for $n = 3$.

Theorem 1.5. *There is no faithful action of \mathbb{Z}_p on any connected three-manifold M .*

1.1 Rough outline of the proof of Theorem 1.5

We suppose the existence of a continuous injection $\mathbb{Z}_p \rightarrow \text{Homeo}(M)$ and derive a contradiction.

Since $p^k \mathbb{Z}_p \cong \mathbb{Z}_p$, we may replace \mathbb{Z}_p with one of its subgroups $p^k \mathbb{Z}_p$ for any large $k \geq 0$. The subgroups $p^k \mathbb{Z}_p \subseteq \mathbb{Z}_p$ form a neighborhood base of the identity in \mathbb{Z}_p ; hence by continuity of the action, as $k \rightarrow \infty$ these subgroups converge to the identity map on M . By picking a suitable Euclidean chart of M and a suitably large $k \geq 0$, we reduce to the case where M is an open subset of \mathbb{R}^3 and the action of \mathbb{Z}_p is very close to the identity.¹

¹There are two motivations for this reduction (which is valid in any dimension). First, recall that a topological group is NSS (“has no small subgroups”) iff there exists an open neighborhood of the identity which contains no nontrivial subgroup. Then a theorem of Yamabe [47, p364 Theorem 3] says that a locally

The next step is to produce a compact connected \mathbb{Z}_p -invariant subset $Z \subseteq M$ satisfying the following two properties:²

1. On a coarse scale, Z looks like a handlebody of genus two.
2. The action of \mathbb{Z}_p on $H^1(Z)$ is nontrivial.

The eventual contradiction will come by combining the first (coarse) property of Z with the second (fine) property of Z . Constructing such a set Z follows a natural strategy: we take the orbit of a closed handlebody of genus two and attach the orbit of a small loop connecting two points on the boundary. However, the construction requires checking certain connectedness properties of a number of different orbit sets, and is currently the least transparent part of the proof.

Now we consider an open set U defined roughly as $N_\epsilon(Z) \setminus Z$ (only roughly, since we need to ensure that U is \mathbb{Z}_p -invariant and $H_2(U) = \mathbb{Z}$). The set of isotopy classes of incompressible surfaces in U representing a generator of $H_2(U)$ forms a lattice, and this lets us find an incompressible surface F in U which is fixed up to isotopy by \mathbb{Z}_p . We think of the surface F as a sort of “approximate boundary” of Z . Even though \mathbb{Z}_p does not act naturally on F itself, we do get a natural homomorphism $\mathbb{Z}_p \rightarrow \text{MCG}(F)$ with finite image. The two properties of Z translate into the following two properties of the action of \mathbb{Z}_p on $H_1(F)$:

1. There is a submodule of $H_1(F)$ fixed by \mathbb{Z}_p on which the intersection form is:

$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \oplus \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \tag{1.2}$$

2. The action of \mathbb{Z}_p on $H_1(F)$ is nontrivial.

This means we have a cyclic subgroup $\mathbb{Z}/p \subseteq \text{MCG}(F)$ such that $H_1(F)^{\mathbb{Z}/p}$ has a submodule on which the intersection form is given by (1.2). The Nielsen classification of cyclic subgroups of the mapping class group shows that this is impossible. This contradiction completes the proof of Theorem 1.5.

We conclude this introduction with a few additional remarks on the proof. First, the proof is a local argument on M , similar in that respect to the proof of Newman’s theorem [24]. Second, our proof works essentially verbatim with any pro-finite group in place of \mathbb{Z}_p (though this is not particularly surprising). Finally, we remark that assuming the action of \mathbb{Z}_p on M is *free* (as is traditional in some approaches to Conjecture 1.2) does not produce any significant simplifications to the argument.

compact topological group is a Lie group iff it is NSS. Thus the relevant property of \mathbb{Z}_p which distinguishes it from a Lie group is the existence of the small subgroups $p^k\mathbb{Z}_p$. Second, recall Newman’s theorem [24] which implies that a compact Lie group acting nontrivially on a manifold must have large orbits. In essence, we are extending Newman’s theorem to the group \mathbb{Z}_p (however the proof will be quite different).

²The motivation to consider such a set is to attempt a *dimension reduction* argument. In other words, we would like to conclude that ∂Z is a closed surface with a faithful action of \mathbb{Z}_p , and therefore contradicts the (known) case of Conjecture 1.2 with $n = 2$. This, of course, is not possible since Z could *a priori* have wild boundary. We will nevertheless be able to construct a closed surface F which serves as an “approximate boundary” of Z .

1.2 Acknowledgements

We thank Ian Agol for suggesting Lemmas 2.14 and 2.19 concerning lattice properties of incompressible surfaces. We also thank Mike Freedman and Steve Kerckhoff for helpful conversations. We thank the referee for their comments and for giving this paper a very close reading.

The author was partially supported by a National Science Foundation Graduate Research Fellowship under grant number DGE-1147470.

2 The lattice of incompressible surfaces

In this section, we study a natural partial order on the set of incompressible surfaces in a particularly nice class of three-manifolds; we call such three-manifolds “quasicylinders” (see Definition 2.4). For a quasicylinder M , we let $\mathcal{S}(M)$ denote the set of isotopy classes of incompressible surfaces in M which generate $H_2(M)$. The main result of this section (suggested by Ian Agol [1]) is that $\mathcal{S}(M)$ is a lattice (see Lemma 2.19) under its natural partial order. Related ideas may be found in papers of Schultens [36] and Przytycki–Schultens [30] on the contractibility of the Kakimizu complex [14].³

Since we will ultimately use the results of this section to study groups of *homeomorphisms*, we need constructions which are functorial with respect to homeomorphisms. To study the properties of these constructions, however, we use methods and results in PL/DIFF three-manifold theory (for example, general position). Thus in this section we will always state explicitly which category (TOP/PL/DIFF) we are working in, since this will change frequently (and there is no straightforward way of working just in a single category). We will, of course, need the key result that every topological three-manifold can be triangulated, as proved by Moise [21] and Bing [3] (see also Hamilton [12] for a modern proof based on the methods of Kirby–Siebenmann [16]; the PL structure is unique, but we do not need this).

By surface, we always mean a closed connected orientable surface. By isotopy (resp. PL isotopy) of surfaces, we always mean ambient isotopy through homeomorphisms (resp. PL homeomorphisms).

Theorem 2.1 ([3, p62 Theorem 8]). *Let M_1, M_2 be two PL three-manifolds where M_2 has a metric d . Let $\phi : M_1 \rightarrow M_2$ be a homeomorphism, and let $f : M_1 \rightarrow \mathbb{R}_{>0}$ be continuous. Then there exists a PL homeomorphism $\phi_1 : M_1 \rightarrow M_2$ with $d(\phi(x), \phi_1(x)) \leq f(x)$.*

Lemma 2.2. *Let M be a PL three-manifold, and let F be a bicollared surface in M . Then there exists an isotopy of M supported in an arbitrarily small neighborhood of F which maps F to a PL surface.*

Proof. Let $\phi : F \times [-1, 1] \rightarrow M$ be a bicollar, which we may assume is arbitrarily close to $F = \phi(F \times \{0\})$. Now pick a PL structure on F and apply Theorem 2.1 to $\phi|_{F \times (0,1)}$ and a

³The Kakimizu complex is a sort of “ \mathbb{Z} -equivariant order complex” of $\mathcal{S}(\widetilde{\mathbb{S}^3 \setminus K})$, where $\widetilde{\mathbb{S}^3 \setminus K}$ denotes the infinite cyclic cover of the knot complement $\mathbb{S}^3 \setminus K$. Even though technically $\widetilde{\mathbb{S}^3 \setminus K}$ is not a quasicylinder under our definition, it is easy to give a modified definition (allowing manifolds with boundary) to which the methods of this section would apply.

function f which decreases sufficiently rapidly near the ends of $(0, 1)$. The resulting ϕ_1 then extends continuously to $F \times [0, 1]$ and agrees with ϕ on $F \times \{0, 1\}$. Now splice $\phi_1|_{F \times [0, 1]}$ and $\phi|_{F \times [-1, 0]}$ together to get a bicollar $\phi_2 : F \times [-1, 1] \rightarrow M$ which is PL on $F \times (0, 1)$. Now using the bicollar ϕ_2 we can easily construct an isotopy of M which sends $F = \phi_2(F \times \{0\})$ to $\phi_2(F \times \{\frac{1}{2}\})$, which is PL. \square

Lemma 2.3. *Let M be an irreducible orientable TOP (resp. PL) three-manifold, and let F_1, F_2 be two bicollared (resp. PL) π_1 -injective surfaces in M . If F_1, F_2 are homotopic, then there is a compactly supported (resp. PL) isotopy of M which sends F_1 to F_2 .*

Proof. Waldhausen [43, p76 Corollary 5.5] proves this in the PL category if M is compact with boundary. It is clear that this implies our PL statement, since the given homotopy will be supported in a compact region of M , which is in turn contained in a compact irreducible submanifold with boundary.

For the TOP category, first pick a PL structure on M , and use Lemma 2.2 to straighten F_1, F_2 by a compactly supported isotopy. Now use the PL version of this lemma. \square

Definition 2.4. A *quasicylinder* is an irreducible orientable three-manifold M with exactly two ends and $H_2(M) \cong \mathbb{Z}$. For example, $\Sigma_g \times \mathbb{R}$ is a quasicylinder for $g \geq 1$.

Lemma 2.5. *Let M be a quasicylinder. For an embedded surface $F \subseteq M$, the following are equivalent:*

1. F is nonzero in $H_2(M)$.
2. F separates the two ends of M .
3. F generates $H_2(M)$.

Proof. (1) \implies (2). A path from one end to the other gives a non-torsion class in $H_1^{\text{lf}}(M)$. Thus its Poincaré dual is a non-torsion class in $H^2(M)$, and thus defines a nonzero map $H_2(M) \rightarrow \mathbb{Z}$. Since F is nonzero in $H_2(M) \cong \mathbb{Z}$, every such path must therefore intersect F .

(2) \implies (3). If F separates the two ends, then there is a path from one end to the other which intersects F exactly once. Thus the Poincaré dual of the class of this path in $H_1^{\text{lf}}(M)$ evaluates to 1 on $F \in H_2(M)$. Thus F represents a primitive element of $H_2(M) \cong \mathbb{Z}$, and thus generates it.

(3) \implies (1). Trivial. \square

Definition 2.6. For a TOP quasicylinder M , let $\mathcal{S}_{\text{TOP}}(M)$ be the set of bicollared π_1 -injective embedded surfaces in M generating $H_2(M)$, modulo homotopy.

Definition 2.7. For a PL quasicylinder M , let $\mathcal{S}_{\text{PL}}(M)$ be the set of PL π_1 -injective embedded surfaces in M generating $H_2(M)$, modulo homotopy.

Remark 2.8. $\mathcal{S}_{\text{PL}}(M)$ is always nonempty, since we can pick a PL embedded surface representing a generator of $H_2(M)$ and then take some maximal compression thereof (as in the proof of Lemma 2.17 below), which will be π_1 -injective by the loop theorem.

Definition 2.9. A *directed quasicylinder* is a quasicylinder along with a labelling of the ends with \pm . For an embedded surface $F \subseteq M$ separating the two ends, let $(M \setminus F)_{\pm}$ denote the two connected components of $M \setminus F$ (the subscripts corresponding to the labelling of the ends). It is easy to see that $(M \setminus F)_{\pm}$ are both (directed) quasicylinders.

Definition 2.10. Let M be a directed quasicylinder. For two embedded surfaces $F_1, F_2 \subseteq M$ separating the two ends of M , we say $F_1 \leq F_2$ iff $F_1 \subseteq (M \setminus F_2)_-$. Define a relation \leq on $\mathcal{S}_{\text{TOP}}(M)$ (resp. $\mathcal{S}_{\text{PL}}(M)$) by declaring that $\mathfrak{F}_1 \leq \mathfrak{F}_2$ if and only if $\mathfrak{F}_1, \mathfrak{F}_2$ have embedded representatives F_1, F_2 with $F_1 \leq F_2$.

Lemma 2.11. *Let M be a PL directed quasicylinder. Then the natural map $\psi : \mathcal{S}_{\text{PL}}(M) \rightarrow \mathcal{S}_{\text{TOP}}(M)$ is a bijection satisfying $\mathfrak{F}_1 \leq \mathfrak{F}_2 \iff \psi(\mathfrak{F}_1) \leq \psi(\mathfrak{F}_2)$.*

Proof. The natural map $(\mathcal{S}_{\text{PL}}(M), \leq) \rightarrow (\mathcal{S}_{\text{TOP}}(M), \leq)$ is clearly injective, and by Lemma 2.2 it is surjective.

If $\mathfrak{F}_1, \mathfrak{F}_2$ have bicollared representatives F_1, F_2 with $F_1 \leq F_2$, then by Lemma 2.2 they can be straightened preserving the relation $F_1 \leq F_2$. The other direction is obvious, since PL surfaces have a bicollar. \square

Having established that $\mathcal{S}_{\text{PL}}(M)$ and $\mathcal{S}_{\text{TOP}}(M)$ are naturally isomorphic, we henceforth use the notation $\mathcal{S}(M)$ for both.

Lemma 2.12. *Let M be a PL quasicylinder. Suppose F is a PL embedded surface in M separating the two ends of M , and suppose γ is a PL arc from one end of M to the other which intersects F transversally in exactly one point. Denote by $\pi_1(M, \gamma)$ one of the groups $\{\pi_1(M, p)\}_{p \in \gamma}$ (they are all naturally isomorphic given the path γ).⁴ Then for any surface $G \subseteq M$ homotopic to F , there is a canonical map $\pi_1(G) \rightarrow \pi_1(M, \gamma)$ (defined up to inner automorphism of the domain).*

Proof. Assume that G intersects γ transversally. Let us call two intersections of γ with G *equivalent* iff there is a path between them on G which, when spliced with the path between them on γ , becomes null-homotopic in M (this is an equivalence relation). Note that during a general position homotopy of G , the mod 2 cardinalities of the equivalence classes of intersections with γ remain the same. Thus since G is homotopic to F and $\#(F \cap \gamma) = 1$, there is a unique equivalence class of intersection $G \cap \gamma$ of odd cardinality. Picking any one of these points as basepoint on G and on γ gives the same map $\pi_1(G) \rightarrow \pi_1(M, \gamma)$ up to inner automorphism of the domain. This map is clearly constant under homotopy of G . \square

Lemma 2.13. *Let M be a PL quasicylinder. Let F be a PL π_1 -injective surface in M separating the two ends of M . Then any homotopy of F to itself induces the trivial element of $\text{MCG}(F)$.*

Proof. Pick a PL arc γ from one end of M to the other which intersects F transversally exactly once. By Lemma 2.12, we get a canonical map $\pi_1(F) \rightarrow \pi_1(M, \gamma)$ which is constant as we move F by homotopy. Since this map is injective, we know $\pi_1(F)$ up to inner automorphism as a subgroup of $\pi_1(M, \gamma)$. \square

⁴A categorical way of phrasing this is as follows. Let γ denote the category whose objects are points $p \in \gamma$, with a single morphism $p \rightarrow p'$ for all $p, p' \in \gamma$. The fundamental group is a functor $\pi_1(M, \cdot) : \gamma \rightarrow \mathbf{Groups}$, where the morphism $p \rightarrow p'$ is sent to the isomorphism $\pi_1(M, p) \rightarrow \pi_1(M, p')$ given by the path from p to p' along γ . Then $\pi_1(M, \gamma)$ is defined as the limit/colimit of this functor.

Lemma 2.14. *Let M be a directed quasicylinder. Then the pair $(\mathcal{S}(M), \leq)$ is a partially-ordered set. That is, for all $\mathfrak{F}_1, \mathfrak{F}_2, \mathfrak{F}_3 \in \mathcal{S}(M)$ we have:*

1. (reflexivity) $\mathfrak{F}_1 \leq \mathfrak{F}_1$.
2. (antisymmetry) $\mathfrak{F}_1 \leq \mathfrak{F}_2 \leq \mathfrak{F}_1 \implies \mathfrak{F}_1 = \mathfrak{F}_2$.
3. (transitivity) $\mathfrak{F}_1 \leq \mathfrak{F}_2 \leq \mathfrak{F}_3 \implies \mathfrak{F}_1 \leq \mathfrak{F}_3$.

Proof. By Lemma 2.11, it suffices to work in the PL category.

Reflexivity is obvious.

For transitivity, suppose $\mathfrak{F}_1 \leq \mathfrak{F}_2 \leq \mathfrak{F}_3$. Then pick representatives F_1, F_2, F'_2, F'_3 such that $F_1 \leq F_2$ and $F'_2 \leq F'_3$. By Lemma 2.3, there is a PL isotopy from F_2 to F'_2 . Applying this isotopy to F_1 produces $F'_1 \leq F'_2 \leq F'_3$.

For antisymmetry, suppose $\mathfrak{F}_1 \leq \mathfrak{F}_2$ and $\mathfrak{F}_2 \leq \mathfrak{F}_1$. Pick PL representatives $F_1 \leq F_2 \leq F'_1 \leq F'_2$ (this is possible by the argument used for transitivity). Now pick a PL arc γ from one end of M to the other which intersects each of F_1, F'_1, F_2, F'_2 exactly once. Since the maps $\pi_1(F_1), \pi_1(F'_1), \pi_1(F_2), \pi_1(F'_2) \rightarrow \pi_1(M, \gamma)$ are injective, we will identify each of the groups on the left with its image in $\pi_1(M, \gamma)$. By Lemma 2.12, we have $\pi_1(F_i) = \pi_1(F'_i)$ under this identification.

By van Kampen's theorem we have:

$$\pi_1(M, \gamma) = \pi_1((M \setminus F_2)_-, \gamma) \underset{\pi_1(F_2)}{*} \pi_1((M \setminus F_2)_+, \gamma) \quad (2.1)$$

Since $F_1 \leq F_2 \leq F'_1$, we have $\pi_1(F_1) \subseteq \pi_1((M \setminus F_2)_-, \gamma)$ and $\pi_1(F'_1) \subseteq \pi_1((M \setminus F_2)_+, \gamma)$. Since $\pi_1(F_1) = \pi_1(F'_1)$, we have $\pi_1(F_1) \subseteq \pi_1((M \setminus F_2)_-, \gamma) \cap \pi_1((M \setminus F_2)_+, \gamma)$. It follows from the properties of the amalgamated product (2.1) that this intersection is just $\pi_1(F_2)$. Thus we have $\pi_1(F_1) \subseteq \pi_1(F_2)$. A symmetric argument shows the reverse inclusion, so we have $\pi_1(F_1) = \pi_1(F_2)$. Now an easy obstruction theory argument shows F_1 and F_2 are homotopic (using the fact that $\pi_2(M) = 0$ by the sphere theorem [29]). Thus $\mathfrak{F}_1 = \mathfrak{F}_2$. \square

Lemma 2.15. *Let M be a directed quasicylinder. Let F be a bicollared embedded π_1 -injective surface. Then the natural map $\psi : \mathcal{S}((M \setminus F)_-) \rightarrow \mathcal{S}(M)$ is a bijection of $\mathcal{S}((M \setminus F)_-)$ with the set $\{\mathfrak{G} \in \mathcal{S}(M) \mid \mathfrak{G} \leq [F]\}$. Furthermore, this bijection satisfies $\mathfrak{G}_1 \leq \mathfrak{G}_2 \iff \psi(\mathfrak{G}_1) \leq \psi(\mathfrak{G}_2)$.*

Proof. By Lemma 2.11, it suffices to work in the PL category.

Certainly the image of ψ is contained in $\{\mathfrak{G} \in \mathcal{S}(M) \mid \mathfrak{G} \leq [F]\}$. If $\mathfrak{G} \leq [F]$, then there are representatives $G' \leq F'$. By Lemma 2.3, there is a PL isotopy sending F' to F . Applying this isotopy to G' gives a representative $G \leq F$. Thus the image of ψ is exactly $\{\mathfrak{G} \in \mathcal{S}(M) \mid \mathfrak{G} \leq [F]\}$. To prove that ψ is injective, suppose $G_1, G_2 \leq F$ are two π_1 -injective embedded surfaces which are homotopic in M . Pick an arc γ from one end of M to the other which intersects G_1, F exactly once. Since G_1, G_2 are homotopic, Lemma 2.12 gives canonical maps $\pi_1(G_1), \pi_1(G_2) \rightarrow \pi_1(M, \gamma)$ with the same image. Now these both factor through $\pi_1((M \setminus F)_-, \gamma) \rightarrow \pi_1(M, \gamma)$, which is injective since $\pi_1(F) \rightarrow \pi_1((M \setminus F)_\pm, \gamma)$ are injective. Thus $\pi_1(G_1), \pi_1(G_2) \rightarrow \pi_1((M \setminus F)_-, \gamma)$ have the same image, so the same obstruction theory argument used in the proof of Lemma 2.14 implies that G_1, G_2 are homotopic in $(M \setminus F)_-$. Thus ψ is injective.

Now it remains to show that ψ preserves \leq . The only nontrivial direction is to show that $\psi(\mathfrak{G}_1) \leq \psi(\mathfrak{G}_2) \implies \mathfrak{G}_1 \leq \mathfrak{G}_2$. If $\psi(\mathfrak{G}_1) \leq \psi(\mathfrak{G}_2)$, then we have representatives $G'_1 \leq G'_2 \leq F'$ (by the argument for transitivity in Lemma 2.14). Now by Lemma 2.3 there is a PL isotopy from F' to F , and applying this isotopy to G'_1, G'_2 , we get $G_1 \leq G_2 \leq F$. Since ψ is injective, we have $[G_i] = \mathfrak{G}_i$ in $\mathcal{S}((M \setminus F)_-)$. Thus $\mathfrak{G}_1 \leq \mathfrak{G}_2$. \square

Lemma 2.16. *Let M be a directed quasicylinder. Let $F_1 \leq F_2$ be two bicollared embedded π_1 -injective surfaces. Then the natural map $\psi : \mathcal{S}((M \setminus F_1)_+ \cap (M \setminus F_2)_-) \rightarrow \mathcal{S}(M)$ is a bijection of $\mathcal{S}((M \setminus F_1)_+ \cap (M \setminus F_2)_-)$ with $\{\mathfrak{G} \in \mathcal{S}(M) \mid [F_1] \leq \mathfrak{G} \leq [F_2]\}$. Furthermore, this bijection satisfies $\mathfrak{G}_1 \leq \mathfrak{G}_2 \iff \psi(\mathfrak{G}_1) \leq \psi(\mathfrak{G}_2)$.*

Proof. This is just two applications of Lemma 2.15. \square

Lemma 2.17. *Let M be a PL directed quasicylinder. Let $G \subseteq M$ be any PL embedded surface (not necessarily π_1 -injective) separating the two ends of M . Then there exists a class $\mathfrak{G} \in \mathcal{S}(M)$ such that for all $\mathfrak{F} \in \mathcal{S}(M)$ with a representative $F \leq G$ (resp. $F \geq G$), we have $\mathfrak{F} \leq \mathfrak{G}$ (resp. $\mathfrak{F} \geq \mathfrak{G}$).*

Proof. As long as G is compressible, we can perform the following operation. Do some compression on G (this may disconnect G), and pick one of the resulting connected components which is nonzero in $H_2(M)$ to keep. This operation does not destroy the property that G can be isotoped to lie in $(M \setminus F)_+$ (resp. $(M \setminus F)_-$) for an incompressible surface F . Since each step decreases the genus of G , we eventually reach an incompressible surface, which by the loop theorem is π_1 -injective, and thus defines a class $\mathfrak{G} \in \mathcal{S}(M)$. \square

Lemma 2.18. *Let M be a directed quasicylinder. Let $A \subseteq \mathcal{S}(M)$ be a finite set. Then there exist elements $\mathfrak{A}_-, \mathfrak{A}_+ \in \mathcal{S}(M)$ such that $\mathfrak{A}_- \leq \mathfrak{A} \leq \mathfrak{A}_+$ for all $\mathfrak{A} \in A$.*

Proof. By Lemma 2.11, it suffices to work in the PL category.

Pick PL transverse representatives A_1, \dots, A_n of all the surfaces in A . Let $((M \setminus A_1)_\pm \cap \dots \cap (M \setminus A_n)_\pm)_0$ denote the unique unbounded component of $(M \setminus A_1)_\pm \cap \dots \cap (M \setminus A_n)_\pm$, and define $A_\pm = \partial(((M \setminus A_1)_\pm \cap \dots \cap (M \setminus A_n)_\pm)_0)$. It is easy to see that A_\pm are connected and separate the two ends of M . Now apply Lemma 2.17 to A_- and A_+ to get the desired classes $\mathfrak{A}_-, \mathfrak{A}_+ \in \mathcal{S}(M)$. \square

Lemma 2.19 (suggested by Ian Agol [1]). *Let M be a directed quasicylinder. Then the partially-ordered set $(\mathcal{S}(M), \leq)$ is a lattice. That is, for $\mathfrak{F}_1, \mathfrak{F}_2 \in \mathcal{S}(M)$, the following set has a least element:*

$$X(\mathfrak{F}_1, \mathfrak{F}_2) = \{\mathfrak{H} \in \mathcal{S}(M) \mid \mathfrak{F}_1, \mathfrak{F}_2 \leq \mathfrak{H}\} \quad (2.2)$$

(and the same holds in the reverse ordering).

Proof. By Lemma 2.11, it suffices to work in the PL category.

First, let us deal with the case where M has tame ends, that is M is the interior of a compact PL manifold with boundary $(M, \partial M)$. Equip $(M, \partial M)$ with a smooth structure and a smooth Riemannian metric so that the boundary is convex.

Now let us recall some facts about area minimizing surfaces. The standard existence theory of Schoen–Yau [35] and Sacks–Uhlenbeck [33, 34] gives the existence of area minimizing

maps in any homotopy class of π_1 -injective surfaces. By Osserman [28, 27] and Gulliver [11], such area minimizing maps are immersions. Now Freedman–Hass–Scott [7] have proved theorems to the effect that area minimizing representatives of π_1 -injective surfaces in irreducible three-manifolds have as few intersections as possible given their homotopy class. Though they state their main results only for closed irreducible three-manifolds, they remark [7, p634 §7] that their results remain true for compact irreducible three-manifolds with convex boundary.

Suppose $\mathfrak{F}_1, \mathfrak{F}_2 \in \mathcal{S}(M)$; let us show that $X(\mathfrak{F}_1, \mathfrak{F}_2)$ has a least element. Pick area minimizing representatives F_1, F_2 in the homotopy classes of $\mathfrak{F}_1, \mathfrak{F}_2$. By [7, p626 Theorem 5.1], F_1, F_2 are embedded. Let $((M \setminus F_1)_+ \cap (M \setminus F_2)_+)_0$ denote the unique unbounded component of $(M \setminus F_1)_+ \cap (M \setminus F_2)_+$, and define $F = \partial(((M \setminus F_1)_+ \cap (M \setminus F_2)_+)_0)$. It is easy to see that F is connected and separates the two ends of M . Note that F_1, F_2 can be isotoped to lie in $(M \setminus F)_-$. Furthermore, if $\mathfrak{G} \in X(\mathfrak{F}_1, \mathfrak{F}_2)$ (and $\mathfrak{G} \neq \mathfrak{F}_1, \mathfrak{F}_2$), then \mathfrak{G} has a least area representative G , which by [7, p630 Theorem 6.2] is disjoint from F_1 and F_2 . Thus $F_1, F_2 \leq G$, so $F \leq G$. Thus we may apply Lemma 2.17 to F and produce an element $\mathfrak{F} \in \mathcal{S}(M)$ which is a least element of $X(\mathfrak{F}_1, \mathfrak{F}_2)$. This finishes the proof for M with tame ends.

Now let us treat the case of general M . Consider the following sets:

$$X(\mathfrak{F}_1, \mathfrak{F}_2; \mathfrak{G}) = \{\mathfrak{H} \in \mathcal{S}(M) \mid \mathfrak{F}_1, \mathfrak{F}_2 \leq \mathfrak{H} \leq \mathfrak{G}\} \subseteq X(\mathfrak{F}_1, \mathfrak{F}_2) \quad (2.3)$$

We claim that it suffices to show that if $\mathfrak{G} \in X(\mathfrak{F}_1, \mathfrak{F}_2)$, then $X(\mathfrak{F}_1, \mathfrak{F}_2; \mathfrak{G})$ has a least element. To see this, we argue as follows. If $\mathfrak{G}_1, \mathfrak{G}_2 \in X(\mathfrak{F}_1, \mathfrak{F}_2)$ and $\mathfrak{G}_1 \leq \mathfrak{G}_2$, then it is easy to see that the natural inclusion $X(\mathfrak{F}_1, \mathfrak{F}_2; \mathfrak{G}_1) \rightarrow X(\mathfrak{F}_1, \mathfrak{F}_2; \mathfrak{G}_2)$ sends the least element of the domain to the least element of the target (assuming both sets have least elements). Since for every $\mathfrak{G}_1, \mathfrak{G}_2$, there exists \mathfrak{G}_3 greater than both (by Lemma 2.18), we see that the least elements of all the sets $\{X(\mathfrak{F}_1, \mathfrak{F}_2; \mathfrak{G})\}_{\mathfrak{G} \in X(\mathfrak{F}_1, \mathfrak{F}_2)}$ are actually the same element $\mathfrak{F} \in X(\mathfrak{F}_1, \mathfrak{F}_2)$. We claim that this \mathfrak{F} is a least element of $X(\mathfrak{F}_1, \mathfrak{F}_2)$. This is trivial: if $\mathfrak{G} \in X(\mathfrak{F}_1, \mathfrak{F}_2)$, then \mathfrak{F} is a least element of $X(\mathfrak{F}_1, \mathfrak{F}_2; \mathfrak{G})$, so *a fortiori* $\mathfrak{F} \leq \mathfrak{G}$. Thus it suffices to show that each set $X(\mathfrak{F}_1, \mathfrak{F}_2; \mathfrak{G})$ has a least element.

Let us show that $X(\mathfrak{F}_1, \mathfrak{F}_2; \mathfrak{G})$ has a least element. By Lemma 2.18 there exists $\mathfrak{F}_0 \in \mathcal{S}(M)$ so that $\mathfrak{F}_0 \leq \mathfrak{F}_1, \mathfrak{F}_2$. Now pick PL representatives F_0, G of $\mathfrak{F}_0, \mathfrak{G}$ with $F_0 \leq G$. Let $M_0 = (M \setminus F_0)_+ \cap (M \setminus G)_-$, which is a directed quasicylinder with tame ends. Thus by the case dealt with earlier, $\mathcal{S}(M_0)$ is a lattice, so $\mathfrak{F}_1, \mathfrak{F}_2$ have a least upper bound in $\mathcal{S}(M_0)$. By Lemma 2.16, $\mathcal{S}(M_0) \rightarrow \mathcal{S}(M)$ is an isomorphism onto the subset $\{\mathfrak{F} \in \mathcal{S}(M) \mid \mathfrak{F}_0 \leq \mathfrak{F} \leq \mathfrak{G}\}$. Thus we get the desired least element of $X(\mathfrak{F}_1, \mathfrak{F}_2; \mathfrak{G})$. \square

Remark 2.20. It should be possible to prove an existence result for area minimizing surfaces in *any* DIFF quasicylinder (with appropriate conditions on the Riemannian metric near the ends). In that case, the argument used for tame quasicylinders would apply in general (the results of Freedman–Hass–Scott [7] extend as long as one has the existence of area minimizing representatives of π_1 -injective surfaces in M).

Remark 2.21. Jaco–Rubinstein [13] have developed a theory of *normal surfaces* in triangulated three-manifolds analogous to that of minimal surfaces in smooth three-manifolds with

a Riemannian metric. In particular, they have results analogous to those of Freedman–Hass–Scott [7]. Thus it is likely that we could eliminate entirely the use of the smooth category and the results in minimal surface theory for the proof of Lemma 2.19.

3 Tools applicable to arbitrary open/closed subsets of manifolds

In our study of a hypothetical action of \mathbb{Z}_p by homeomorphisms on a three-manifold, it will be essential to study certain properties of orbit sets (for example, their homology). However, we do not have the luxury of assuming such sets are at all well behaved; the most we can hope for is that we will be able to construct \mathbb{Z}_p -invariant sets which are either open or closed. Nevertheless, Čech cohomology is still a reasonable object in such situations. The purpose of this section is to develop an elementary theory centered on Alexander duality for arbitrary open and closed subsets of a manifold. We use these tools in the proof of Theorem 1.5, specifically in Sections 4.3–4.4 to construct the set Z and to study its properties.

In this (and all other) sections, we always take homology and cohomology with integer coefficients. We denote by H_* and H^* singular homology and cohomology, and we let \check{H}^* denote Čech cohomology.

3.1 Čech cohomology

Lemma 3.1. *For a compact subset X of a manifold, the natural map below is an isomorphism:*

$$\lim_{\substack{U \supseteq X \\ U \text{ open}}} H^*(U) \xrightarrow{\sim} \check{H}^*(X) \quad (3.1)$$

Proof. This is due to Steenrod [39]; see also Spanier [38, p419], the key fact being that Čech cohomology satisfies the “continuity axiom”. \square

Lemma 3.2. *If X and Y are two compact subsets of a manifold, then there is a (Mayer–Vietoris) long exact sequence:*

$$\cdots \rightarrow \check{H}^*(X \cup Y) \rightarrow \check{H}^*(X) \oplus \check{H}^*(Y) \rightarrow \check{H}^*(X \cap Y) \rightarrow \check{H}^{*+1}(X \cup Y) \rightarrow \cdots \quad (3.2)$$

Proof. For arbitrary open neighborhoods $U \supseteq X$ and $V \supseteq Y$, we have a Mayer–Vietoris sequence of singular cohomology for U and V . Taking the direct limit over all U and V gives a sequence of the form (3.2) by Lemma 3.1, and it is exact since the direct limit is an exact functor. \square

3.2 Alexander duality

Alexander duality for subsets of \mathbb{S}^n is most naturally stated using reduced homology and cohomology, which we denote by \tilde{H} .

Lemma 3.3. *Let $X \subseteq \mathbb{S}^n$ be a compact set. Then:*

$$\tilde{H}^*(X) = \tilde{H}_{n-1-*}(\mathbb{S}^n \setminus X) \quad (3.3)$$

Proof. If $U \subseteq \mathbb{S}^n$ is an open subset with smooth boundary, then it is well-known that there is a natural isomorphism $\tilde{H}^*(U) \xrightarrow{\sim} \tilde{H}_{n-1-*}(\mathbb{S}^n \setminus U)$. Furthermore, for $U \supseteq V$ the following diagram commutes:

$$\begin{array}{ccc} \tilde{H}^*(U) & \xrightarrow{\sim} & \tilde{H}_{n-1-*}(\mathbb{S}^n \setminus U) \\ \downarrow & & \downarrow \\ \tilde{H}^*(V) & \xrightarrow{\sim} & \tilde{H}_{n-1-*}(\mathbb{S}^n \setminus V) \end{array} \quad (3.4)$$

Now consider the direct limit over all such open sets $U \supseteq X$ to get an isomorphism of the direct limits. This family of sets U forms a final system of neighborhoods of X , so by Lemma 3.1 the direct limit on the left is $\tilde{H}^*(X)$. The direct limit on the right is $\tilde{H}_{n-1-*}(\mathbb{S}^n \setminus X)$ by elementary properties of singular homology. \square

3.3 The plus operation

If we have a bounded subset $X \subseteq \mathbb{R}^3$, we want to be able to consider “ X union all of the bounded connected components of $\mathbb{R}^3 \setminus X$ ” (we think of this operation as a way to simplify the set X , which could be very wild). The following definition makes this precise.

Definition 3.4. For a bounded set $X \subseteq \mathbb{R}^3$, we define:

$$X^+ = \bigcap_{\substack{\text{open } U \supseteq X \\ U \text{ bounded} \\ H_2(U)=0}} U \quad (3.5)$$

By Lemma 3.3, for bounded open sets $U \subseteq X$, we have $H_2(U) = 0 \iff \tilde{H}^0(\mathbb{R}^3 \setminus U) = \mathbb{Z}$. With this reformulation, it is easy to see that if U and V both appear in the intersection, then so does $U \cap V$. Let us call $X \mapsto X^+$ the *plus operation*.

Lemma 3.5. *The plus operation satisfies the following properties:*

- i.* $X \subseteq Y \implies X^+ \subseteq Y^+$.
- ii.* $X^{++} = X^+$.
- iii.* If X is closed and bounded, then $\mathbb{R}^3 \setminus X^+$ is the unbounded component of $\mathbb{R}^3 \setminus X$.
- iv.* If X is closed and bounded, then $X = X^+ \iff \tilde{H}^2(X) = 0 \iff \mathbb{R}^3 \setminus X$ is connected.

Proof. Both (i) and (ii) are immediate from the definition.

For (iii), argue as follows. Let V denote the unbounded component of $\mathbb{R}^3 \setminus X$. As V is open and connected, it has an exhaustion by closed connected subsets $V = \bigcup_{i=1}^{\infty} V_i$, where $V_1 \subseteq V_2 \subseteq \dots$ and $\mathbb{R}^3 \setminus V_i$ is bounded for all i . By Lemma 3.3, $H_2(\mathbb{R}^3 \setminus V_i) = 0$, and thus $X^+ \subseteq \bigcap_{i=1}^{\infty} (\mathbb{R}^3 \setminus V_i) = \mathbb{R}^3 \setminus V$. It remains to show the reverse inclusion, namely that if W is a bounded component of $\mathbb{R}^3 \setminus X$, then $W \subseteq X^+$. Suppose $U \supseteq X$ is open, bounded, and $H_2(U) = 0$. Then $\mathbb{R}^3 \setminus U = (\mathbb{R}^3 \setminus (U \cup W)) \cup (W \setminus U)$, which is a disjoint union of closed sets.

However $\mathbb{R}^3 \setminus U$ is connected and $\mathbb{R}^3 \setminus (U \cup W)$ is unbounded so *a fortiori* it is nonempty. Thus we must have $W \setminus U = \emptyset$, that is $U \supseteq W$.

For (iv), argue as follows. Lemma 3.3 gives $\check{H}^2(X) = 0 \iff \mathbb{R}^3 \setminus X$ is connected. By (iii), we have $X = X^+ \iff \mathbb{R}^3 \setminus X$ is connected. \square

Lemma 3.6. *Suppose $X \subseteq \mathbb{R}^3$ is closed and bounded with $X = X^+$. If $\{U_\alpha\}_{\alpha \in A}$ is a final collection of open sets containing X , then so is $\{U_\alpha^+\}_{\alpha \in A}$.*

Proof. Since $\{U_\alpha\}_{\alpha \in A}$ is final, to show that $\{U_\alpha^+\}_{\alpha \in A}$ is final it suffices to show that for every $\alpha \in A$, there exists $\beta \in A$ such that $U_\beta^+ \subseteq U_\alpha$.

Thus let us suppose $\alpha \in A$ is given. As in the proof of Lemma 3.5(iii), there exists an exhaustion by closed connected subsets $\mathbb{R}^3 \setminus X = \bigcup_{i=1}^\infty V_i$, where $V_1 \subseteq V_2 \subseteq \dots$ and $\mathbb{R}^3 \setminus V_i$ is bounded for all i . Then for sufficiently large i , we have $\mathbb{R}^3 \setminus V_i \subseteq U_\alpha$. On the other hand, for every i there exists β such that $U_\beta \subseteq \mathbb{R}^3 \setminus V_i$. But now $U_\beta^+ \subseteq (\mathbb{R}^3 \setminus V_i)^+ = \mathbb{R}^3 \setminus V_i \subseteq U_\alpha$ as needed. \square

Definition 3.7. Let X be a topological space. We say X has dimension zero iff whenever $x \in U \subseteq X$ with U open, there exists a clopen (closed and open) set $V \subseteq X$ with $x \in V \subseteq U$. It is immediate that any subspace of a space of dimension zero also has dimension zero.

Remark 3.8. If M is a manifold with a continuous action of \mathbb{Z}_p , then every orbit is either a finite discrete set (if the stabilizer is $p^k \mathbb{Z}_p$) or a cantor set (if the stabilizer is trivial). In particular, every orbit has dimension zero. It follows that if $A \subseteq M$ is any finite set, then $\mathbb{Z}_p A$ has dimension zero, as does any subset thereof.

Lemma 3.9. *If M is a manifold of dimension $n \geq 2$ and $X \subseteq M$ is closed and has dimension zero, then the map $H_0(M \setminus X) \rightarrow H_0(M)$ is an isomorphism.*

Proof. First, let us deal with the case $M = \mathbb{S}^n$ (thus X is compact). Pick a metric on \mathbb{S}^n and fix $\epsilon > 0$. Since X has dimension zero and is compact, we can cover X with finitely many clopen sets $U_i \subseteq X$ ($1 \leq i \leq N$) of diameter $\leq \epsilon$. Now $\{U_i \setminus \bigcup_{1 \leq j < i} U_j\}_{i=1}^N$ is a cover of X by disjoint open sets of diameter $\leq \epsilon$. Since $\epsilon > 0$ was arbitrary, this implies that $\check{H}^i(X) = 0$ for $i > 0$. Thus by Lemma 3.3, $\check{H}_0(\mathbb{S}^n \setminus X) = \check{H}^{n-1}(X) = 0$ (since $n \geq 2$), which is sufficient.

Second, let us deal with the case $M = \mathbb{R}^n$. Define $\bar{X} = X \cup \{\infty\} \subseteq \mathbb{R}^n \cup \{\infty\} = \mathbb{S}^n$. We claim that \bar{X} has dimension zero; to see this, it suffices to produce small clopen neighborhoods of $\infty \in \bar{X}$. Fix $R < \infty$. Since X has dimension zero and $X \cap \overline{B(\mathbf{0}, R)}$ is compact, there exist finitely many clopen sets $U_i \subseteq X$ ($1 \leq i \leq N$) of diameter ≤ 1 with $X \cap \overline{B(\mathbf{0}, R)} \subseteq \bigcup_{i=1}^N U_i$. Then $\bar{X} \setminus \bigcup_{i=1}^N U_i$ is clopen in \bar{X} and contains ∞ . Since $R < \infty$ was arbitrary, we get arbitrarily small clopen neighborhoods of $\infty \in \bar{X}$. Hence \bar{X} has dimension zero, so by the case $M = \mathbb{S}^n$ dealt with above, we have $H_0(\mathbb{R}^n \setminus X) = H_0(\mathbb{S}^n \setminus \bar{X}) = \mathbb{Z}$, which is sufficient.

Now let us deal with the case of general M . It suffices to show that $H^0(M) \rightarrow H^0(M \setminus X)$ is an isomorphism (recall that H^0 is just the group of locally constant maps to \mathbb{Z}). Now consider any $p \in M$, and pick an open neighborhood $p \in U \cong \mathbb{R}^n$. Since X has dimension zero, so does $U \cap X$. Thus by the case $M = \mathbb{R}^n$ dealt with above, $U \setminus X$ is connected and nonempty. Thus any locally constant function $M \setminus X \rightarrow \mathbb{Z}$ can be extended uniquely to a locally constant function $M \rightarrow \mathbb{Z}$, as needed. \square

4 Nonexistence of faithful actions of \mathbb{Z}_p on three-manifolds

Proof of Theorem 1.5. Let M be a connected three-manifold and suppose there is a continuous injection of topological groups $\mathbb{Z}_p \rightarrow \text{Homeo}(M)$.

4.1 Step 1: Reduction to a local problem in \mathbb{R}^3

Let $B(r)$ denote the open ball of radius r centered at the origin $\mathbf{0}$ in \mathbb{R}^3 . Fix a small positive number $\eta = 2^{-10}$. In this section, we reduce to the case where M is an open subset of \mathbb{R}^3 such that:

- i. $B(1) \subseteq M \subseteq B(1 + \eta)$.
- ii. $d_{\mathbb{R}^3}(x, \alpha x) \leq \eta$ for all $x \in M$ and $\alpha \in \mathbb{Z}_p$.
- iii. No subgroup $p^k \mathbb{Z}_p$ fixes an open neighborhood of $\mathbf{0}$.

This reduction is valid in any dimension.

Definition 4.1. We say that an action of a group G on a topological space X is *locally of finite order* at a point $x \in X$ if and only if some subgroup of finite index $G' \leq G$ fixes an open neighborhood of x . The action is (*globally*) *of finite order* if and only if some subgroup of finite index $G' \leq G$ fixes all of X .

Newman's theorem [24, p6 Theorem 2] (see also Dress [6, p204 Theorem 1] or Smith [37]) implies that on a connected manifold, a group action which is (everywhere) locally of finite order is globally of finite order. Thus if $\mathbb{Z}_p \rightarrow \text{Homeo}(M)$ is injective, then there exists a point $m \in M$ where the action is not locally of finite order. In other words, no subgroup $p^k \mathbb{Z}_p$ fixes an open neighborhood of m .

Now fix some homeomorphism between $B(3) \subseteq \mathbb{R}^3$ and an open set in M containing m , such that $\mathbf{0}$ is identified with m . Note that (by continuity of the map $\mathbb{Z}_p \rightarrow \text{Homeo}(M)$) for sufficiently large k , we have $p^k \mathbb{Z}_p B(2) \subseteq B(3)$ and $d_{\mathbb{R}^3}(x, \alpha x) \leq \eta$ for all $x \in B(2)$ and $\alpha \in p^k \mathbb{Z}_p$. Thus the action of $p^k \mathbb{Z}_p$ (which is isomorphic to \mathbb{Z}_p) on $M' := p^k \mathbb{Z}_p B(1)$ satisfies (i), (ii), and (iii) above. Thus it suffices to replace (M, \mathbb{Z}_p) with $(M', p^k \mathbb{Z}_p)$ and derive a contradiction.

Henceforth we assume that M is an open subset of \mathbb{R}^3 satisfying (i), (ii), and (iii) above.

4.2 Step 2: An invariant metric and the plus operation in M

Equip M with the following \mathbb{Z}_p -invariant metric, which induces the same topology as $d_{\mathbb{R}^3}$:

$$d_{\text{inv}}(x, y) := \int_{\mathbb{Z}_p} d_{\mathbb{R}^3}(\alpha x, \alpha y) d\mu_{\text{Harr}}(\alpha) \quad (4.1)$$

We will never mention d_{inv} again explicitly, but we will often write N_ϵ^{inv} for the open ϵ -neighborhood in M with respect to d_{inv} . Note that if $X \subseteq M$ is \mathbb{Z}_p -invariant, then so is $N_\epsilon^{\text{inv}}(X)$.

Lemma 4.2. *If $X \subseteq B(1) \subseteq M$, then $X^+ \subseteq B(1) \subseteq M$. If in addition X is \mathbb{Z}_p -invariant, then so is X^+ .*




Figure 1: Diagram of K_0 inside $B(1) \subseteq M$.

Proof. The first statement is clear since $B(1)$ appears in the intersection (3.5) defining X^+ .

Now suppose that in addition X is \mathbb{Z}_p -invariant. As remarked in Definition 3.4, the collection of open sets appearing in the intersection (3.5) is closed under finite intersection. In particular, we could consider just those U contained in $B(1)$. Thus *a fortiori* X^+ is the intersection of all open sets U with $X \subseteq U \subseteq M$ and $H_2(U) = 0$. The set of such U is permuted by \mathbb{Z}_p , so X^+ is \mathbb{Z}_p -invariant. \square

4.3 Step 3: Construction of an interesting compact set $Z \subseteq M$

In this section, we construct a compact \mathbb{Z}_p -invariant set $Z \subseteq M$ such that:

1. On a coarse scale, Z looks like a handlebody of genus two.
2. The action of \mathbb{Z}_p on $\check{H}^1(Z)$ is nontrivial.

We follow a natural strategy to construct the set Z . We first let K be the orbit under \mathbb{Z}_p of a closed handlebody of genus two, and we then construct a set L as the orbit under \mathbb{Z}_p of a small arc connecting two points on the boundary of K . Then we let $Z = K \cup L$. This strategy is complicated by the fact that the orbit set K is *a priori* very wild, and we will need to investigate the connectedness properties of it and other sets in order to make the construction work. We will take advantage of the tools we have developed in Section 3.3 and the setup from Sections 4.1–4.2. Even with this preparation, though, careful verification of each step takes us a while.

Let $x_0 \in B(\eta) \setminus \text{Fix } \mathbb{Z}_p$ (such an x_0 exists by Step 1(iii)).

Definition 4.3 (see Figure 1). Let $K_0 \subseteq B(1)$ be a closed handlebody of genus two whose unique point of lowest z -coordinate is x_0 . Note that the closed η -neighborhood (in the Euclidean metric) of K_0 is a larger handlebody isotopic to K_0 . Let $K = (\mathbb{Z}_p K_0)^+$.

We think of K as being illustrated essentially by Figure 1; it's just K_0 plus some wild fuzz of width η . Note that K is compact, \mathbb{Z}_p -invariant (by Lemma 4.2), and $K = K^+$. Also, K is connected; actually, we will need the following stronger fact.

Lemma 4.4. *If $A \subseteq \partial K$ is any finite set, then $K \setminus \mathbb{Z}_p A$ is path-connected.*

Proof. By definition, K_0° is path-connected. Now the action of \mathbb{Z}_p is within $\eta = 2^{-10}$ of the identity, so $\mathbb{Z}_p(K_0^\circ)$ is also path-connected.

Now suppose $x \in \mathbb{Z}_p K_0 \setminus \mathbb{Z}_p A$. Then there exists $\alpha \in \mathbb{Z}_p$ so that $\alpha^{-1}x \in K_0$. From $\alpha^{-1}x$ there is clearly a path inwards to K_0° , and this is disjoint from $\mathbb{Z}_p A$ since $\mathbb{Z}_p A \subseteq \partial K$. Translating this path by α shows that $\mathbb{Z}_p K_0 \setminus \mathbb{Z}_p A$ is path-connected.

Now suppose $x \in K \setminus \mathbb{Z}_p A$. If $x \notin \mathbb{Z}_p K_0$, then let V denote the (open) connected component of $K \setminus \mathbb{Z}_p K_0$ containing x . By Remark 3.8 and Lemma 3.9, we can find a path from x to infinity in $\mathbb{R}^3 \setminus \mathbb{Z}_p A$. There is a time when this path first hits $\partial V \subseteq \mathbb{Z}_p K_0$, thus giving us a path from x to $\mathbb{Z}_p K_0$ contained in $K \setminus \mathbb{Z}_p A$. Thus $K \setminus \mathbb{Z}_p A$ is path-connected, as was to be shown. \square

Lemma 4.5. *There exist points $x_1, x_2 \in K$ with $x_1 \notin \text{Fix } \mathbb{Z}_p$ and $\mathbb{Z}_p x_1 \cap \mathbb{Z}_p x_2 = \emptyset$, along with a compact \mathbb{Z}_p -invariant set $L \subseteq M$ of diameter $\leq 4\eta$ so that $L = L^+$, $K \cap L = \mathbb{Z}_p x_1 \cup \mathbb{Z}_p x_2$, and L contains a path from x_1 to x_2 .*

Proof. Let $x_1 \in K$ be any point of lowest z -coordinate in K . We claim that $x_1 \notin \text{Fix } \mathbb{Z}_p$. First, observe that since $x_1 \in K = (\mathbb{Z}_p K_0)^+$ is a point of lowest z -coordinate, necessarily $x_1 \in \mathbb{Z}_p K_0$. If $x_1 \in K_0$, then $x_1 = x_0$ (recall from Definition 4.3 that x_0 is the unique point of lowest z -coordinate in K_0), and by definition $x_0 \notin \text{Fix } \mathbb{Z}_p$. On the other hand, if $x_1 \in (\mathbb{Z}_p K_0) \setminus K_0$, then certainly $x_1 \notin \text{Fix } \mathbb{Z}_p$. Thus $x_1 \notin \text{Fix } \mathbb{Z}_p$.

Since $x_1 \in \mathbb{Z}_p K_0$, we may pick $x'_2 \in \mathbb{Z}_p(K_0^\circ) \subseteq K^\circ$ which is arbitrarily close to x_1 . Specifically, let us fix $x'_2 \in K^\circ \cap B(x_1, \eta)$. Note that $x'_2 \notin \mathbb{Z}_p x_1$ since $x_1 \notin K^\circ$. Now we claim that there exists a continuous path $\gamma : [0, 1] \rightarrow B(x_1, \eta)$ such that:

- i. $\gamma(0) = x_1$.
- ii. $\gamma(1) = x'_2$.
- iii. $\gamma^{-1}(\mathbb{Z}_p x_1) = \{0\}$.
- iv. 0 is not a limit point of $\gamma^{-1}(K)$.

To construct such a path γ , argue as follows. First, define $\gamma : [0, \frac{1}{2}] \rightarrow B(x_1, \eta)$ to be some path straight downward from x_1 (this takes care of (i)). Since x_1 is a point of smallest z -coordinate in $K \supseteq \mathbb{Z}_p x_1$, this also takes care of (iv) and is consistent with (iii). Now by Remark 3.8 and Lemma 3.9, we know $B(x_1, \eta) \setminus \mathbb{Z}_p x_1$ is path-connected, so we can find a path $\gamma : [\frac{1}{2}, 1] \rightarrow B(x_1, \eta) \setminus \mathbb{Z}_p x_1$ from $\gamma(\frac{1}{2})$ to x'_2 (this takes care of (ii) and (iii)). Splicing these two functions together gives $\gamma : [0, 1] \rightarrow B(x_1, \eta)$ satisfying (i), (ii), (iii), and (iv).

Now by (ii) and (iv) we have that $t := \min(\gamma^{-1}(K) \setminus \{0\})$ exists and is positive. Let $x_2 = \gamma(t)$, and define $L_0 = \gamma([0, t])$. By (iii), we have $\mathbb{Z}_p x_1 \cap \mathbb{Z}_p x_2 = \emptyset$. Define $L = (\mathbb{Z}_p L_0)^+$, which is \mathbb{Z}_p -invariant by Lemma 4.2. By construction, L contains a path from x_1 to x_2 . Certainly $L_0 \subseteq B(x_1, \eta)$ so $L \subseteq B(x_1, 2\eta)$ by Step 1(ii), and so L has diameter $\leq 4\eta$. It remains only to show that $K \cap L = \mathbb{Z}_p x_1 \cup \mathbb{Z}_p x_2$ (certainly the containment \supseteq is given by definition).

Note that $K \setminus \mathbb{Z}_p L_0 = K \setminus (\mathbb{Z}_p x_1 \cup \mathbb{Z}_p x_2)$, which by Lemma 4.4 is path-connected. Thus $K \setminus \mathbb{Z}_p L_0$ lies in a single connected component of $\mathbb{R}^3 \setminus \mathbb{Z}_p L_0$. Now $\mathbb{Z}_p L_0$ has diameter $\leq 4\eta$ as remarked above, and $K \setminus \mathbb{Z}_p L_0$ contains a large handlebody, so it must lie in the unbounded component of $\mathbb{R}^3 \setminus \mathbb{Z}_p L_0$. Hence $K \setminus (\mathbb{Z}_p x_1 \cup \mathbb{Z}_p x_2)$ is disjoint from $(\mathbb{Z}_p L_0)^+ = L$. \square

Definition 4.6. Fix x_1, x_2 , and L as exist by Lemma 4.5, and define $Z = K \cup L$.

4.4 Step 4: The cohomology of Z

Lemma 4.7. $Z = Z^+$, and the action of \mathbb{Z}_p on $\check{H}^1(Z)$ is nontrivial.

Proof. The key to proving both statements is the following Mayer–Vietoris sequence (which exists and is exact by Lemma 3.2):

$$\cdots \rightarrow \check{H}^*(Z) \rightarrow \check{H}^*(K) \oplus \check{H}^*(L) \rightarrow \check{H}^*(K \cap L) \rightarrow \check{H}^{*+1}(Z) \rightarrow \cdots \quad (4.2)$$

Note that \mathbb{Z}_p acts on all of these groups and that the maps are \mathbb{Z}_p -equivariant.

By Lemma 4.5, $K \cap L = \mathbb{Z}_p x_1 \cup \mathbb{Z}_p x_2$, so by Remark 3.8, $\check{H}^i(K \cap L) = 0$ for $i > 0$. Thus we have $\check{H}^2(Z) = \check{H}^2(K) \oplus \check{H}^2(L)$. Now applying Lemma 3.5(iv) twice shows that $K = K^+$ and $L = L^+$ together imply that $Z = Z^+$.

Now let us show that \mathbb{Z}_p acts on $\check{H}^1(Z)$ nontrivially. We use the exact sequence:

$$\check{H}^0(K) \oplus \check{H}^0(L) \rightarrow \check{H}^0(K \cap L) \rightarrow \check{H}^1(Z) \quad (4.3)$$

Note that \check{H}^0 is just the group of locally constant functions to \mathbb{Z} . Thus it suffices to exhibit a locally constant function $q : K \cap L \rightarrow \mathbb{Z}$ and an element $\alpha \in \mathbb{Z}_p$ such that $\alpha q - q$ is not in the image of $\check{H}^0(K) \oplus \check{H}^0(L)$. Let us define:

$$q(r) = \begin{cases} 1 & r \in p\mathbb{Z}_p x_1 \\ 0 & r \in \bigcup_{a \in (\mathbb{Z}/p) \setminus \{0\}} (a + p\mathbb{Z}_p)x_1 \cup \mathbb{Z}_p x_2 \end{cases} \quad (4.4)$$

(observe by Lemma 4.5 that $x_1 \notin \text{Fix } \mathbb{Z}_p$, so the sets on the right hand side are indeed disjoint). Let $\alpha \in \mathbb{Z}_p$ be congruent to 1 mod p . Then we have:

$$(\alpha q - q)(r) = \begin{cases} -1 & r \in p\mathbb{Z}_p x_1 \\ 1 & r \in (1 + p\mathbb{Z}_p)x_1 \\ 0 & r \in \bigcup_{a \in (\mathbb{Z}/p) \setminus \{0,1\}} (a + p\mathbb{Z}_p)x_1 \cup \mathbb{Z}_p x_2 \end{cases} \quad (4.5)$$

Now by Lemma 4.5, x_1, x_2 are in the same component of L , and by Lemma 4.4 (taking $A = \emptyset$), they are in the same component of K . Thus every function in the image of $\check{H}^0(L) \oplus \check{H}^0(K)$ assigns the same value to x_1 and x_2 . Clearly this is not the case for $\alpha q - q$, so we are done. \square

We just proved that $Z = Z^+$, so by Lemma 3.6, $N_\epsilon^{\text{inv}}(Z)^+$ is a final system of neighborhoods of Z . Thus by Lemma 3.1 we have $\check{H}^1(Z) = \varinjlim H^1(N_\epsilon^{\text{inv}}(Z)^+)$ as abelian groups with an action of \mathbb{Z}_p (\mathbb{Z}_p acts on the latter since $N_\epsilon^{\text{inv}}(Z)^+$ is \mathbb{Z}_p -invariant by Lemma 4.2). Since the action on the limit group is nontrivial, the following definition makes sense.

Definition 4.8. Fix $\epsilon \in (0, \eta)$ such that the \mathbb{Z}_p action on the image of the map $H^1(N_\epsilon^{\text{inv}}(Z)^+) \rightarrow \check{H}^1(Z)$ is nontrivial. Denote by $(N_\epsilon^{\text{inv}}(Z)^+)_0$ the connected component containing Z , and define $U = (N_\epsilon^{\text{inv}}(Z)^+)_0 \setminus Z$, which is \mathbb{Z}_p -invariant by Lemma 4.2.

4.5 Step 5: Special elements of $\mathcal{S}(U)$

Lemma 4.9. *The set U is a quasicylinder in the sense of Definition 2.4.*

Proof. We have $\mathbb{R}^3 \setminus U = Z \cup (\mathbb{R}^3 \setminus (N_\epsilon^{\text{inv}}(Z)^+)_0)$; the latter two sets are connected and disjoint, so by Lemma 3.3, we have $H_2(U) = \mathbb{Z}$.

Suppose $F \subseteq U$ is a PL embedded sphere; then $\text{int}(F)$ (the bounded component of $\mathbb{R}^3 \setminus F$) is a ball [2]. If $Z \subseteq \text{int}(F)$, then we have a factorization $H^1(N_\epsilon^{\text{inv}}(Z)^+) \rightarrow H^1(\text{int}(F)) \rightarrow \check{H}^1(Z)$, which implies the composition is trivial, contradicting the definition of U . Thus $Z \not\subseteq \text{int}(F)$, so $\text{int}(F) \subseteq U$, that is F bounds ball in U . Thus U is irreducible.

Certainly U has at least two ends, and since $H_2(U) = \mathbb{Z}$ it has at most two ends.

Since U is an open subset of \mathbb{R}^3 , it is orientable. \square

Recall the definition of $(\mathcal{S}(U), \leq)$ from Section 2. Certainly the action of \mathbb{Z}_p on U induces an action on $\mathcal{S}_{\text{TOP}}(U)$.

Lemma 4.10. *There exists an element $\mathfrak{F} \in \mathcal{S}(U)$ which is fixed by \mathbb{Z}_p .*

Proof. Observe that if F is any surface in U , then for sufficiently large k , we have that αF is homotopic to F for all $\alpha \in p^k \mathbb{Z}_p$. Thus \mathbb{Z}_p acts on $\mathcal{S}(U)$ with finite orbits. Now any group acting with finite orbits on a nonempty lattice has a fixed point, namely the least upper bound of any orbit. Recall that $\mathcal{S}(U)$ is nonempty by Remark 2.8. \square

Definition 4.11. Fix a PL π_1 -injective surface $F \subseteq U$ such that $[F] \in \mathcal{S}(U)$ is fixed by \mathbb{Z}_p . Denote by $\text{int}(F)$ and $\text{ext}(F)$ the two connected components of $\mathbb{R}^3 \setminus F$.

Lemma 4.12. *The \mathbb{Z}_p action on U induces a homomorphism $\mathbb{Z}_p \rightarrow \text{MCG}(F)$, as well as actions on all the homology and cohomology groups appearing in (4.6)–(4.9). These actions are compatible with the maps in (4.6)–(4.9), as well as with the map $\text{MCG}(F) \rightarrow \text{Aut}(H_1(F))$.*

$$\begin{array}{ccccc} H^1(N_\epsilon^{\text{inv}}(Z)^+) & \longrightarrow & H^1(\text{int}(F)) & \longrightarrow & \check{H}^1(Z) \\ & & \downarrow & & \\ & & H^1(F) & & \end{array} \quad (4.6)$$

$$H_1(Z) \rightarrow H_1(\text{int}(F)) \quad (4.7)$$

$$H_1(M \setminus N_\epsilon^{\text{inv}}(Z)^+) \rightarrow H_1(\text{ext}(F)) \quad (4.8)$$

$$H_1(F) \xrightarrow{\sim} H_1(\text{int}(F)) \oplus H_1(\text{ext}(F)) \quad (4.9)$$

Proof. For every $\alpha \in \mathbb{Z}_p$, we know by Lemma 2.3 that there is a compactly supported isotopy of U sending αF to F . Now such an isotopy can clearly be extended to all of M as the constant isotopy on $M \setminus U$. Thus for every $\alpha \in \mathbb{Z}_p$, let us pick an isotopy:

$$\psi_\alpha^t : M \rightarrow M \quad (t \in [0, 1]) \quad (4.10)$$

so that $\psi_\alpha^0 = \text{id}_M$, $\psi_\alpha^1(\alpha F) = F$, and $\psi_\alpha^t(x) = x$ for $x \in M \setminus U$. Denote by $T_\alpha : M \rightarrow M$ the action of $\alpha \in \mathbb{Z}_p$.

Now observe that for every $\alpha \in \mathbb{Z}_p$, the map $\psi_\alpha^1 \circ T_\alpha$ fixes Z , $N_\epsilon^{\text{inv}}(Z)^+$, and F . Thus the map $\psi_\alpha^1 \circ T_\alpha$ induces an automorphism of each of the diagrams (4.6)–(4.9) and gives a compatible element of $\text{MCG}(F)$. It remains only to show that this is a homomorphism from \mathbb{Z}_p .

We need to show that for all $\alpha, \beta \in \mathbb{Z}_p$, the two maps $\psi_\beta^1 \circ T_\beta \circ \psi_\alpha^1 \circ T_\alpha$ and $\psi_{\beta\alpha}^1 \circ T_{\beta\alpha}$ induce the same action on (4.6)–(4.9) and give the same element of $\text{MCG}(F)$. It of course suffices to show that $\psi_\beta^1 \circ T_\beta \circ \psi_\alpha^1 \circ T_\alpha \circ (\psi_{\beta\alpha}^1 \circ T_{\beta\alpha})^{-1}$ induces the trivial action on (4.6)–(4.9) and gives the trivial element of $\text{MCG}(F)$. Now we write:

$$\begin{aligned} \psi_\beta^1 \circ T_\beta \circ \psi_\alpha^1 \circ T_\alpha \circ (\psi_{\beta\alpha}^1 \circ T_{\beta\alpha})^{-1} &= \psi_\beta^1 \circ T_\beta \circ \psi_\alpha^1 \circ T_\alpha \circ T_{\beta\alpha}^{-1} \circ (\psi_{\beta\alpha}^1)^{-1} \\ &= \psi_\beta^1 \circ T_\beta \circ \psi_\alpha^1 \circ T_\beta^{-1} \circ (\psi_{\beta\alpha}^1)^{-1} \end{aligned} \quad (4.11)$$

This is the identity map on Z and on $M \setminus N_\epsilon^{\text{inv}}(Z)^+$, so the action on their (co)homology is trivial. The map (4.11) is isotopic to the identity map via $\psi_\beta^t \circ T_\beta \circ \psi_\alpha^t \circ T_\beta^{-1} \circ (\psi_{\beta\alpha}^t)^{-1}$ for $t \in [0, 1]$, which only moves points in a compact subset of U . Thus the action on the cohomology of $N_\epsilon^{\text{inv}}(Z)^+$ is trivial as well. By Lemma 2.13, this isotopy induces the trivial element in $\text{MCG}(F)$, and this means that the action on the (co)homology of F , $\text{int}(F)$, and $\text{ext}(F)$ is trivial as well. \square

Lemma 4.13. *The map $\mathbb{Z}_p \rightarrow \text{MCG}(F)$ annihilates an open subgroup of \mathbb{Z}_p and has non-trivial image.*

Proof. The action of \mathbb{Z}_p on U is continuous, so for sufficiently large k , we have that αF and F are homotopic as maps $F \rightarrow U$ for all $\alpha \in p^k \mathbb{Z}_p$. Thus the homomorphism $\mathbb{Z}_p \rightarrow \text{MCG}(F)$ annihilates a neighborhood of the identity in \mathbb{Z}_p .

To prove that the image is nontrivial, it suffices to show that the \mathbb{Z}_p action on $H^1(F)$ is nontrivial. For this, consider equation (4.6). By Definition 4.8, there exists an element of $H^1(N_\epsilon^{\text{inv}}(Z)^+)$ whose image in $\check{H}^1(Z)$ is not fixed by \mathbb{Z}_p . Thus the action of \mathbb{Z}_p on $H^1(\text{int}(F))$ is nontrivial. Now the vertical map $H^1(\text{int}(F)) \rightarrow H^1(F)$ is injective (otherwise the Mayer–Vietoris sequence applied to $\mathbb{R}^3 = \text{int}(F) \cup_F \text{ext}(F)$ would imply $H^1(\mathbb{R}^3) \neq 0$). Thus the action of \mathbb{Z}_p on $H^1(F)$ is nontrivial as well. \square

Lemma 4.14. *There is a rank four submodule of $H_1(F)^{\mathbb{Z}_p}$ on which the intersection form is:*

$$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \oplus \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad (4.12)$$

Proof. There are two obvious loops in K_0 (Definition 4.3) generating its homology, and two obvious dual loops in $M \setminus N_\epsilon^{\text{inv}}(Z)^+$. Since the \mathbb{Z}_p action is within $\eta = 2^{-10}$ of the identity, it is easy to see that their classes in homology are fixed by \mathbb{Z}_p . Thus we have well-defined classes $\alpha_1, \alpha_2 \in H_1(Z)^{\mathbb{Z}_p}$ and $\beta_1, \beta_2 \in H_1(M \setminus N_\epsilon^{\text{inv}}(Z)^+)^{\mathbb{Z}_p}$ with linking numbers $\text{lk}(\alpha_i, \beta_j) = \delta_{ij}$. Now the maps from equations (4.7), (4.8), and the isomorphism (4.9) give us corresponding elements $\alpha_1, \alpha_2, \beta_1, \beta_2 \in H_1(F)^{\mathbb{Z}_p}$. The intersection form on $H_1(F)$ coincides with the linking pairing between $H_1(\text{int}(F))$ and $H_1(\text{ext}(F))$. Thus the intersection form on the submodule of $H_1(F)^{\mathbb{Z}_p}$ generated by $\alpha_1, \alpha_2, \beta_1, \beta_2$ is indeed given by (4.12). \square

By Lemma 4.13, the image of \mathbb{Z}_p in $\text{MCG}(F)$ is a nontrivial cyclic p -group. It has a (unique) subgroup isomorphic to \mathbb{Z}/p , and by Lemma 4.14 this subgroup $\mathbb{Z}/p \subseteq \text{MCG}(F)$ has the property that $H_1(F)^{\mathbb{Z}/p}$ has a submodule on which the intersection form is given by (4.12). This contradicts Lemma 5.1 below, and thus completes the proof of Theorem 1.5. \square

Remark 4.15. One expects that if we make U thick enough around most of Z (but still thin near L), then away from L the surface F can be made to look like a punctured surface of genus two. However, it is not clear how to prove this stronger, more geometric property about F .

If we could prove this, then the final analysis is more robust. Instead of homology classes $\alpha_1, \alpha_2, \beta_1, \beta_2 \in H_1(F)^{\mathbb{Z}_p}$, we now have simple closed curves $\alpha_1, \alpha_2, \beta_1, \beta_2$ in F which are fixed up to isotopy by the \mathbb{Z}_p action. Now the finite image of \mathbb{Z}_p in $\text{MCG}(F)$ is realized as a subgroup of $\text{Isom}(F, g)$ for some hyperbolic metric g on F . A hyperbolic isometry fixing $\alpha_1, \alpha_2, \beta_1, \beta_2$ fixes their unique length-minimizing geodesic representatives, and thus fixes their intersections, forcing the isometry to be the identity map. This contradicts the nontriviality of the homomorphism $\mathbb{Z}_p \rightarrow \text{MCG}(F)$. As the reader will readily observe, it is enough that α_1, β_1 be fixed, so we could let K_0 be a handlebody of genus one and this argument would go through.

5 Actions of \mathbb{Z}/p on closed surfaces

Lemma 5.1. *Let F be a closed connected oriented surface, and let $\mathbb{Z}/p \subseteq \text{MCG}(F)$ be some cyclic subgroup of prime order. Then the intersection form of F restricted to $H_1(F)^{\mathbb{Z}/p}$ and taken modulo p has rank at most two.*

The proof below follows Symonds [40, pp389–390 Theorem B] and Chen–Glover–Jensen [5, Theorem 1.1], who make explicit a classification due to Nielsen [25] (in fact, Symonds’ result actually implies ours for $p > 2$).

Proof. We seek simply to classify all subgroups $\mathbb{Z}/p \subseteq \text{MCG}(F)$ and prove that in each case, the conclusion of the lemma is satisfied. This classification is essentially due to Nielsen [25], who showed that finite cyclic subgroups $G \subseteq \text{MCG}(F)$ are classified up to conjugacy by their “fixed point data”.

By work of Nielsen [26], any $\mathbb{Z}/p \subseteq \text{MCG}(F)$ is realized by a genuine action of \mathbb{Z}/p on F by isometries in some metric (see Thurston [42] or Kerckhoff [15] for a modern perspective on this fact and its generalization to any finite group). Now let us switch notation and write $\tilde{S} = F$, $\mathcal{S} = F/(\mathbb{Z}/p)$ (the quotient orbifold), and S for the coarse space of \mathcal{S} . The cover $\tilde{S} \rightarrow \mathcal{S}$ is classified by an element $\alpha \in H^1(\mathcal{S}, \mathbb{Z}/p)$, which is nonzero since \tilde{S} is connected. Let g denote the genus of S , and n the number of orbifold points of \mathcal{S} .

Now for $(\tilde{S}, \mathcal{S}, \alpha, g, n)$ as above, we prove the following more precise statement (which certainly implies the lemma):

$$\text{the intersection form on } H_1(\tilde{S})^{\mathbb{Z}/p} \cong \begin{cases} \left(\begin{smallmatrix} 0 & p \\ -p & 0 \end{smallmatrix} \right)^{\oplus(g-1)} \oplus \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} & n = 0 \\ \left(\begin{smallmatrix} 0 & p \\ -p & 0 \end{smallmatrix} \right)^{\oplus g} & n > 0 \end{cases} \quad (5.1)$$

First, let us treat the case $n = 0$ (so $\mathcal{S} = S$). It is well-known that both compositions $\text{MCG}(\Sigma_g) \rightarrow \text{Sp}(2g, \mathbb{Z}) \rightarrow \text{Sp}(2g, \mathbb{Z}/p)$ are surjective, and that $\text{Sp}(2g, \mathbb{Z}/p)$ acts transitively on the nonzero elements of $(\mathbb{Z}/p)^{\oplus 2g}$. Thus without loss of generality, it suffices to consider one particular nonzero $\alpha \in H^1(S, \mathbb{Z}/p)$. Thus, let us suppose that α is the Poincaré dual of a nonseparating simple closed curve ℓ on S . Then the cover \tilde{S} is obtained by cutting S along ℓ and gluing together in a circle p copies of the resulting cut surface. Now one easily observes that there is a direct sum decomposition respecting intersection forms $H_1(\tilde{S}) \cong H_1(\Sigma_1) \oplus H_1(\Sigma_{g-1})^{\oplus p}$, where \mathbb{Z}/p acts trivially on $H_1(\Sigma_1)$ and acts by rotation of the summands on $H_1(\Sigma_{g-1})^{\oplus p}$. Then $H_1(\tilde{S})^{\mathbb{Z}/p}$ is $H_1(\Sigma_1)$ with its usual intersection form, plus a copy of $H_1(\Sigma_{g-1})$ with its intersection form multiplied by p . Hence equation (5.1) holds in the case $n = 0$.

Now, let us treat the case $n > 0$. Label the orbifold points $p_1, \dots, p_n \in \mathcal{S}$, and define $r_i : H^1(\mathcal{S}, \mathbb{Z}/p) \rightarrow \mathbb{Z}/p$ to be the evaluation on a small loop around p_i . By construction, the orbifold points are precisely the ramification points of $\tilde{S} \rightarrow S$, and hence $r_i(\alpha) \neq 0$ for all i . On the other hand, if $\mathcal{D} \subseteq \mathcal{S}$ is a disk containing all n orbifold points, then $\partial\mathcal{D} = \partial(\mathcal{S} \setminus \mathcal{D})$ is null-homologous in \mathcal{S} , so $\sum_{i=1}^n r_i(\alpha) = 0$. Thus in particular $n \geq 2$.

Next, let us show how to reduce to the case where either $g = 0$ or $n = 2$. So, suppose $g \geq 1$ and $n \geq 3$, and let $\mathcal{D} \subseteq \mathcal{S}$ be a disk containing p_1, \dots, p_{n-1} . Take \mathcal{D} and $\mathcal{S} \setminus \mathcal{D}$ and glue in a disk with a single \mathbb{Z}/p orbifold point to each, and call the resulting closed orbifolds \mathcal{S}' and \mathcal{S}'' respectively. The class $\alpha \in H^1(\mathcal{S}, \mathbb{Z}/p)$ naturally induces $\alpha' \in H^1(\mathcal{S}', \mathbb{Z}/p)$ and $\alpha'' \in H^1(\mathcal{S}'', \mathbb{Z}/p)$, and so we get covers \tilde{S}' and \tilde{S}'' . Since $\langle \alpha, \partial\mathcal{D} \rangle = \sum_{i=1}^{n-1} r_i(\alpha) = -r_n(\alpha) \neq 0$, it follows that $\partial\mathcal{D}$ lifts to a single separating loop in \tilde{S} , which exhibits \tilde{S} as a connected sum of \tilde{S}' and \tilde{S}'' . Hence $H_1(\tilde{S}) = H_1(\tilde{S}') \oplus H_1(\tilde{S}'')$ respecting the intersection form and the \mathbb{Z}/p action. Now $(g(\mathcal{S}'), n(\mathcal{S}')) = (0, n(\mathcal{S}))$ and $(g(\mathcal{S}''), n(\mathcal{S}'')) = (g(\mathcal{S}), 2)$, so knowing equation (5.1) for \tilde{S}' and \tilde{S}'' implies it for \tilde{S} . Hence it suffices to deal with the two cases $g = 0$ and $n = 2$.

Case $g = 0$. Embed a tree $T = (V, E)$ in S , where $V = \{p_1, \dots, p_{n-1}\}$ and where no edge passes through p_n . This gives a cell decomposition of S , which we lift to a (\mathbb{Z}/p) -equivariant cell decomposition of \tilde{S} . Now let us consider the cellular chains computing $H_1(\tilde{S})$. There is exactly one 2-cell, and it has zero boundary. Thus $H_1(\tilde{S})$ is simply the group of 1-cycles. Now a 1-chain is fixed by \mathbb{Z}/p iff for all $e \in E$ it assigns the same weight to each of the p lifts of e . Now it is easy to observe that since T is a tree, such a 1-chain cannot be a 1-cycle unless it vanishes. Thus $H_1(\tilde{S})^{\mathbb{Z}/p} = 0$, so equation (5.1) holds in the case $g = 0$.

Case $n = 2$. There is a natural exact sequence:

$$0 \rightarrow H^1(S, \mathbb{Z}/p) \rightarrow H^1(\mathcal{S}, \mathbb{Z}/p) \xrightarrow{(r_1, r_2)} (\mathbb{Z}/p)^{\oplus 2} \xrightarrow{+} \mathbb{Z}/p \rightarrow 0 \quad (5.2)$$

Now, multiplying α by a nonzero element of \mathbb{Z}/p yields an equivalent problem, so we assume without loss of generality that $\alpha \in A := (r_1, r_2)^{-1}(1, -1)$. We claim that $\text{MCG}(\mathcal{S} \text{ rel } \{p_1, p_2\})$ acts transitively on A . Let $[\ell] \in A$ be the Poincaré dual of a simple arc ℓ from p_1 to p_2 . We observed earlier that $\text{MCG}(S \text{ rel } \ell)$ acts transitively on the nonzero elements of $H^1(S, \mathbb{Z}/p)$; hence $\text{MCG}(\mathcal{S} \text{ rel } \{p_1, p_2\})$ acts transitively on $A \setminus \{[\ell]\}$ (by exactness). On the other hand, since $g \geq 1$, there certainly exists $\gamma \in \text{MCG}(\mathcal{S} \text{ rel } \{p_1, p_2\})$ for which $\gamma[\ell] - [\ell] \in H^1(S, \mathbb{Z}/p)$ is nonzero (for instance, γ could be a Dehn twist around a nonseparating simple closed curve which intersects ℓ exactly once), and so $[\ell]$ is in the same orbit as $A \setminus \{[\ell]\}$. Thus

$\text{MCG}(\mathcal{S} \text{ rel } \{p_1, p_2\})$ acts transitively on A . Hence we may assume without loss of generality that $\alpha = [\ell]$. Now we can see the cover $\tilde{S} \rightarrow S$ explicitly: we cut S along ℓ and glue together p copies in the relevant fashion. One then easily observes that there is an isomorphism respecting intersection forms $H_1(\tilde{S}) \cong H_1(\Sigma_g)^{\oplus p}$, where \mathbb{Z}/p acts by rotation of the summands. Then $H_1(\tilde{S})^{\mathbb{Z}/p}$ is a copy of $H_1(\Sigma_g)$ with its intersection form multiplied by p . Hence equation (5.1) holds in the case $n = 2$ as well, and the proof is complete. \square

References

- [1] Agol (mathoverflow.net/users/1345). Incompressible surfaces in an open subset of \mathbb{R}^3 . MathOverflow. <http://mathoverflow.net/questions/74935> (version: 2011-09-07).
- [2] J. W. Alexander. On the subdivision of 3-space by a polyhedron. *Proceedings of the National Academy of Sciences of the United States of America*, 10(1):pp. 6–8, 1924.
- [3] R. H. Bing. An alternative proof that 3-manifolds can be triangulated. *Ann. of Math. (2)*, 69:37–65, 1959.
- [4] Salomon Bochner and Deane Montgomery. Locally compact groups of differentiable transformations. *Ann. of Math. (2)*, 47:639–653, 1946.
- [5] Yu Qing Chen, Henry H. Glover, and Craig A. Jensen. Prime order subgroups of mapping class groups. *JP J. Geom. Topol.*, 11(2):87–99, 2011.
- [6] Andreas Dress. Newman’s theorems on transformation groups. *Topology*, 8:203–207, 1969.
- [7] Michael Freedman, Joel Hass, and Peter Scott. Least area incompressible surfaces in 3-manifolds. *Invent. Math.*, 71(3):609–642, 1983.
- [8] A. M. Gleason. The structure of locally compact groups. *Duke Math. J.*, 18:85–104, 1951.
- [9] Andrew M. Gleason. Groups without small subgroups. *Ann. of Math. (2)*, 56:193–212, 1952.
- [10] W. H. Gottschalk. Minimal sets: an introduction to topological dynamics. *Bull. Amer. Math. Soc.*, 64:336–351, 1958.
- [11] Robert D. Gulliver, II. Regularity of minimizing surfaces of prescribed mean curvature. *Ann. of Math. (2)*, 97:275–305, 1973.
- [12] A. J. S. Hamilton. The triangulation of 3-manifolds. *Quart. J. Math. Oxford Ser. (2)*, 27(105):63–70, 1976.
- [13] William Jaco and J. Hyam Rubinstein. PL minimal surfaces in 3-manifolds. *J. Differential Geom.*, 27(3):493–524, 1988.

- [14] Osamu Kakimizu. Finding disjoint incompressible spanning surfaces for a link. *Hiroshima Math. J.*, 22(2):225–236, 1992.
- [15] Steven P. Kerckhoff. The Nielsen realization problem. *Ann. of Math. (2)*, 117(2):235–265, 1983.
- [16] Robion C. Kirby and Laurence C. Siebenmann. *Foundational essays on topological manifolds, smoothings, and triangulations*. Princeton University Press, Princeton, N.J., 1977. With notes by John Milnor and Michael Atiyah, Annals of Mathematics Studies, No. 88.
- [17] Joo Sung Lee. Totally disconnected groups, p -adic groups and the Hilbert-Smith conjecture. *Commun. Korean Math. Soc.*, 12(3):691–699, 1997.
- [18] İozhe Maleshich. The Hilbert-Smith conjecture for Hölder actions. *Uspekhi Mat. Nauk*, 52(2(314)):173–174, 1997.
- [19] Gaven J. Martin. The Hilbert-Smith conjecture for quasiconformal actions. *Electron. Res. Announc. Amer. Math. Soc.*, 5:66–70 (electronic), 1999.
- [20] Mahan Mj. Pattern rigidity and the Hilbert–Smith conjecture. *Geom. Topol.*, 16(2):1205–1246, 2012.
- [21] Edwin E. Moise. Affine structures in 3-manifolds. V. The triangulation theorem and Hauptvermutung. *Ann. of Math. (2)*, 56:96–114, 1952.
- [22] Deane Montgomery and Leo Zippin. Small subgroups of finite-dimensional groups. *Ann. of Math. (2)*, 56:213–241, 1952.
- [23] Deane Montgomery and Leo Zippin. *Topological transformation groups*. Interscience Publishers, New York-London, 1955.
- [24] M. H. A. Newman. A theorem on periodic transformations of spaces. *Quart. J. Math.*, os-2(1):1–8, 1931.
- [25] J. Nielsen. Die struktur periodischer transformationen von flächen. *Danske Vid. Selsk. Mat.-Fys. Medd.*, 15:1–77, 1937.
- [26] Jakob Nielsen. Abbildungsklassen endlicher Ordnung. *Acta Math.*, 75:23–115, 1943.
- [27] Robert Osserman. *A survey of minimal surfaces*. Van Nostrand Reinhold Co., New York, 1969.
- [28] Robert Osserman. A proof of the regularity everywhere of the classical solution to Plateau’s problem. *Ann. of Math. (2)*, 91:550–569, 1970.
- [29] C. D. Papakyriakopoulos. On Dehn’s lemma and the asphericity of knots. *Ann. of Math. (2)*, 66:1–26, 1957.

- [30] Piotr Przytycki and Jennifer Schultens. Contractibility of the Kakimizu complex and symmetric Seifert surfaces. *Trans. Amer. Math. Soc.*, 364(3):1489–1508, 2012.
- [31] Frank Raymond and R. F. Williams. Examples of p -adic transformation groups. *Ann. of Math. (2)*, 78:92–106, 1963.
- [32] Dušan Repovš and Evgenij Ščepin. A proof of the Hilbert-Smith conjecture for actions by Lipschitz maps. *Math. Ann.*, 308(2):361–364, 1997.
- [33] J. Sacks and K. Uhlenbeck. The existence of minimal immersions of 2-spheres. *Ann. of Math. (2)*, 113(1):1–24, 1981.
- [34] J. Sacks and K. Uhlenbeck. Minimal immersions of closed Riemann surfaces. *Trans. Amer. Math. Soc.*, 271(2):639–652, 1982.
- [35] R. Schoen and Shing Tung Yau. Existence of incompressible minimal surfaces and the topology of three-dimensional manifolds with nonnegative scalar curvature. *Ann. of Math. (2)*, 110(1):127–142, 1979.
- [36] Jennifer Schultens. The Kakimizu complex is simply connected. *J. Topol.*, 3(4):883–900, 2010. With an appendix by Michael Kapovich.
- [37] P. A. Smith. Transformations of finite period. III. Newman’s theorem. *Ann. of Math. (2)*, 42:446–458, 1941.
- [38] Edwin H. Spanier. Cohomology theory for general spaces. *Ann. of Math. (2)*, 49:407–427, 1948.
- [39] Norman E. Steenrod. Universal Homology Groups. *Amer. J. Math.*, 58(4):661–701, 1936.
- [40] Peter Symonds. The cohomology representation of an action of C_p on a surface. *Trans. Amer. Math. Soc.*, 306(1):389–400, 1988.
- [41] Terence Tao. Hilbert’s fifth problem and related topics. Manuscript, 2012.
<http://terrytao.wordpress.com/books/hilberts-fifth-problem-and-related-topics/>.
- [42] William P. Thurston. On the geometry and dynamics of diffeomorphisms of surfaces. *Bull. Amer. Math. Soc. (N.S.)*, 19(2):417–431, 1988.
- [43] Friedhelm Waldhausen. On irreducible 3-manifolds which are sufficiently large. *Ann. of Math. (2)*, 87:56–88, 1968.
- [44] John J. Walsh. Light open and open mappings on manifolds. II. *Trans. Amer. Math. Soc.*, 217:271–284, 1976.
- [45] David Wilson. Open mappings on manifolds and a counterexample to the Whyburn conjecture. *Duke Math. J.*, 40:705–716, 1973.

- [46] Hidehiko Yamabe. On the conjecture of Iwasawa and Gleason. *Ann. of Math. (2)*, 58:48–54, 1953.
- [47] Hidehiko Yamabe. A generalization of a theorem of Gleason. *Ann. of Math. (2)*, 58:351–365, 1953.
- [48] Chung-Tao Yang. p -adic transformation groups. *Michigan Math. J.*, 7:201–218, 1960.