

ZELEVINSKY DUALITY ON BASIC LOCAL SHIMURA VARIETIES

LINUS HAMANN

Abstract. We give a simple proof of a general result describing the action of the Zelevinsky involution on the cohomology of certain basic local Shimura varieties, using the machinery of Fargues-Scholze [FS21]. As an application, we generalize earlier results of Fargues [Far06] and Mieda [Mie14] on the action of the Zelevinsky involution on the cohomology of GL_n and GSp_4 type basic local Shimura varieties, respectively.

CONTENTS

Acknowledgements	1
Notation	1
1. Introduction	2
2. Proof of the Main Theorem	6
3. Applications	9
References	10

ACKNOWLEDGEMENTS

I would like to thank David Hansen for helpful discussions, Peter Scholze and the referee for pointing out some small errors, and the MPIM Bonn for their hospitality during part of the completion of this project.

NOTATION

- (1) Let $\ell \neq p$ be distinct primes.
- (2) Let G/\mathbb{Q}_p be a quasi-split connected reductive group over the p -adic numbers.
- (3) Fix $T \subset B \subset G$ a maximal torus and Borel of G .
- (4) Let ρ_G denote the half sum of all positive roots of G .
- (5) Let $Z(G)$ denote the center of G .
- (6) Set $\check{\mathbb{Q}}_p$ to be the completion of the maximal unramified extension of \mathbb{Q}_p , with Frobenius σ .
- (7) Set $\mathbb{C}_p := \widehat{\check{\mathbb{Q}}_p}$ to be the completion of the algebraic closure of \mathbb{Q}_p .
- (8) Let $B(G) = G(\check{\mathbb{Q}}_p)/(g \sim hg\sigma(h)^{-1})$ denote the Kottwitz set of G .
- (9) For an element $b \in B(G)$, we let J_b denote the σ -centralizer of b .
- (10) We let $\mathbb{C}_p^\flat := \lim_{x^i \rightarrow x^p} \mathbb{C}_p$ be the tilt of \mathbb{C}_p , we write X for the (adic) Fargues-Fontaine curve defined by \mathbb{C}_p^\flat .
- (11) To an element $b \in B(G)$, we let \mathcal{E}_b be the associated G -bundle on X .
- (12) Let $\mathrm{Perf}_{\overline{\mathbb{F}}_p}$ denote the category of perfectoid spaces in characteristic p over the algebraic closure $\overline{\mathbb{F}}_p$ of \mathbb{F}_p equipped with the v -topology. For $S \in \mathrm{Perf}_{\overline{\mathbb{F}}_p}$, we write X_S for the relative (adic) Fargues-Fontaine curve over S .
- (13) We let Bun_G denote the v -stack sending $S \in \mathrm{Perf}_{\overline{\mathbb{F}}_p}$ to the groupoid of G -bundles on X_S .
- (14) Let $\mathrm{D}_{\mathrm{lis}}(\mathrm{Bun}_G, \overline{\mathbb{Q}}_\ell)$ (resp. $\mathrm{D}_{\mathrm{lis}}(\mathrm{Bun}_G, \overline{\mathbb{Q}}_\ell)^\omega$) be the category of (resp. compact) lisse-étale $\overline{\mathbb{Q}}_\ell$ -sheaves, as defined in [FS21, Section VII.6].

- (15) For a finite extension E/\mathbb{Q}_p , set W_E to be the Weil group of E . We let $\mathrm{D}_{\mathrm{lis}}(\mathrm{Bun}_G, \overline{\mathbb{Q}}_\ell)^{BW_E}$ be the set of objects in $\mathrm{D}_{\mathrm{lis}}(\mathrm{Bun}_G, \overline{\mathbb{Q}}_\ell)$ with continuous W_E -action, as defined in [FS21, Section IX.1].
- (16) We fix a square root of p in $\overline{\mathbb{Q}}_\ell$. All half Tate twists will be defined with respect to this choice.

1. INTRODUCTION

Let μ be a geometric dominant cocharacter of G with reflex field E/\mathbb{Q}_p , and set $b \in B(G, \mu)$ to be the unique basic element in the μ -admissible locus (Definition 2.1). The triple (G, b, μ) defines a diamond [SW20]

$$\mathrm{Sht}(G, b, \mu)_\infty \rightarrow \mathrm{Spd}(\check{E})$$

parametrizing modifications $\mathcal{E}_b \rightarrow \mathcal{E}_0$ with meromorphy bounded by μ on X , where \check{E} is the compositum $E\overline{\mathbb{Q}}_p$. The space $\mathrm{Sht}(G, b, \mu)_\infty$ carries an action of $G(\mathbb{Q}_p) \times J_b(\mathbb{Q}_p)$ and moreover carries a (non-effective) descent datum from \check{E} down to E . This allows us to consider the tower of quotients

$$\mathrm{Sht}(G, b, \mu)_\infty / \underline{K} =: \mathrm{Sht}(G, b, \mu)_K$$

for varying open compact $K \subset G(\mathbb{Q}_p)$. If μ is minuscule, we say that the triple (G, b, μ) defines a basic local Shimura datum. In this case, the members of this tower are represented by rigid analytic spaces and define local Shimura varieties in the sense of Rapoport-Viehmann [RV14]. Let $\overline{\mathbb{Q}}_\ell$ be the algebraic closure of the ℓ -adic numbers. The geometric Satake equivalence of Fargues-Scholze defines a \mathbb{Z}_ℓ -sheaf \mathcal{S}_μ [FS21, Chapter VI] attached to μ on the spaces $\mathrm{Sht}(G, b, \mu)_K$ compatible with the actions of $G(\mathbb{Q}_p)$ and $J_b(\mathbb{Q}_p)$, and if μ is minuscule \mathcal{S}_μ is (up to a shift and twist) just the constant sheaf. Letting $\mathrm{Sht}(G, b, \mu)_{K, \mathbb{C}_p}$ denote the base-change of these spaces to \mathbb{C}_p , we can now define the complex

$$R\Gamma_c(G, b, \mu) := \mathrm{colim}_{K \rightarrow \{1\}} R\Gamma_c(\mathrm{Sht}(G, b, \mu)_{K, \mathbb{C}_p}, \mathcal{S}_\mu) \otimes \overline{\mathbb{Q}}_\ell$$

of $G(\mathbb{Q}_p) \times J_b(\mathbb{Q}_p) \times W_E$ -modules. It is an important problem to describe the cohomology of these complexes as they are known to realize instances of the local Langlands correspondences and can be related to the cohomology of global Shimura varieties via uniformization in the case that μ is minuscule. More specifically, for π (resp. ρ) a smooth irreducible representation of $G(\mathbb{Q}_p)$ (resp. $J_b(\mathbb{Q}_p)$) one wants to understand the π (resp. ρ)-isotypic part. I.e the complexes

$$R\Gamma_c(G, b, \mu)[\pi] := R\Gamma_c(G, b, \mu) \otimes_{\mathcal{H}(G)}^{\mathbb{L}} \pi$$

and

$$R\Gamma_c(G, b, \mu)[\rho] := R\Gamma_c(G, b, \mu) \otimes_{\mathcal{H}(J_b)}^{\mathbb{L}} \rho$$

where $\mathcal{H}(G) := C_c^\infty(G(\mathbb{Q}_p), \overline{\mathbb{Q}}_\ell)$ (resp. $\mathcal{H}(J_b)$) is the usual smooth Hecke algebra of G (resp. J_b). It is a recent result of Fargues-Scholze that $R\Gamma_c(G, b, \mu)[\pi]$ (resp. $R\Gamma_c(G, b, \mu)[\rho]$) are valued in admissible $J_b(\mathbb{Q}_p)$ (resp. $G(\mathbb{Q}_p)$)-representations of finite length carrying a smooth action of W_E [FS21, Page 317]. Moreover, it is concentrated in degrees $-d \leq i \leq d$, where $d = \dim(\mathrm{Sht}(G, b, \mu)_\infty) = \langle 2\rho_G, \mu \rangle$.

Let ${}^L G$ be the Langlands dual group of G . In instances where the local Langlands correspondences are known and the L -parameter $\phi : W_E \times \mathrm{SL}_2(\mathbb{C}) \rightarrow {}^L G(\mathbb{C})$ associated to π and ρ is supercuspidal (i.e the restriction to the $\mathrm{SL}_2(\mathbb{C})$ -factor acts trivially and ϕ does not factor through a proper Levi subgroup of ${}^L G$) it is expected that $R\Gamma_c(G, b, \mu)[\pi]$ (resp. $R\Gamma_c(G, b, \mu)[\rho]$) should be concentrated in middle degree 0 [Han20, Theorem 1.1] and valued in representations lying in the L -packet $\Pi_\phi(J_b)$ (resp. $\Pi_\phi(G)$). Even in this case, verifying this expectation is a very difficult problem and forms the content of the Kottwitz conjecture, [RV14, Conjecture 7.3].

In this note, we would like to prove a result that concerns the structure of the cohomology of $R\Gamma_c(G, b, \mu)[\pi]$ and $R\Gamma_c(G, b, \mu)[\rho]$ when π or ρ is a supercuspidal, but do not necessarily have supercuspidal L -parameter. This is the case in which the L -parameter is discrete but the SL_2 -factor acts non-trivially. This is a much more difficult problem. Even for $G = \mathrm{GL}_n$, J_b an inner form of GL_n , and $\rho = \mathbf{1}$ the trivial representation, the cohomology of the complex $R\Gamma_c(G, b, \mu)[\rho]$ is concentrated in multiple different degrees [SS91; Dat06], and seems, in general, to be valued in representations lying in both the L -packet $\Pi_\phi(G)$ and the A -packet $\Pi_\psi(G)$ [Ito13; IM22], where ψ is the A -parameter:

$$\psi : W_{\mathbb{Q}_p} \times \mathrm{SL}_2(\mathbb{C}) \times \mathrm{SL}_2(\mathbb{C}) \rightarrow W_{\mathbb{Q}_p} \times \mathrm{SL}_2(\mathbb{C}) \times \mathrm{SL}_2(\mathbb{C}) \xrightarrow{\phi} {}^L G(\mathbb{C})$$

Here the first map is given by swapping the SL_2 -factors and the second map is given by trivially extending ϕ on the second factor. To aid in understanding the cohomology in this case, one considers the (derived) Zelevinsky involution.

Definition 1.1. For G/\mathbb{Q}_p a connected reductive group, let $\mathcal{D}_c(G)$ be the convolution algebra of compactly supported $\overline{\mathbb{Q}_\ell}$ -valued functions on $G(\mathbb{Q}_p)$. It contains the usual smooth Hecke algebra $\mathcal{H}(G)$ as a sub-algebra. We let $\mathrm{D}(\mathcal{D}_c(G), \overline{\mathbb{Q}_\ell})$ (resp. $\mathrm{D}(G(\mathbb{Q}_p), \overline{\mathbb{Q}_\ell})$) denote the unbounded derived category of $\mathcal{D}_c(G)$ -modules (resp. smooth $G(\mathbb{Q}_p)$ -modules). If we let $e_K \in \mathcal{H}(G) \subset \mathcal{D}_c(G)$ be the idempotent for the Hecke algebra defined by a compact open subgroup $K \subset G(\mathbb{Q}_p)$ then we can consider the derived functor

$$\infty_{\mathcal{D}} : \mathrm{D}(\mathcal{D}_c(G), \overline{\mathbb{Q}_\ell}) \rightarrow \mathrm{D}(G(\mathbb{Q}_p), \overline{\mathbb{Q}_\ell})$$

$$A \mapsto \mathrm{colim}_{K \rightarrow \{1\}} e_K A$$

which is in particular exact and so has no higher derived functors (See [Mie14, Section 2] and [Far06, Section 2] for details). Then we define the (derived) Zelevinsky involution to be

$$\mathbb{D}_G : \mathrm{D}(G(\mathbb{Q}_p), \overline{\mathbb{Q}_\ell})^{op} \rightarrow \mathrm{D}(G(\mathbb{Q}_p), \overline{\mathbb{Q}_\ell})$$

$$A \mapsto \infty_{\mathcal{D}} \circ \mathcal{R}Hom_{\mathcal{H}(G)}(A, \mathcal{H}(G))$$

where $\mathcal{H}(G)$ is viewed as a Hecke bi-module.

On the subcategory of compact objects, this defines a contravariant equivalence and has the property that $\mathbb{D}_G^2 \simeq Id$. In this note, we seek to describe the complex $\mathbb{D}_G(R\Gamma_c(G, b, \mu)[\rho])$ and $\mathbb{D}_{J_b}(R\Gamma_c(G, b, \mu)[\pi])$. This is desirable since (up to contragredients and shifts) \mathbb{D}_G should exchange the L -packet $\Pi_\phi(G)$ and the A -packet $\Pi_\psi(G)$ [Hir04], so better understanding the structure of this complex should give insight into the complicated way these different representations appear and interact with each other in the cohomology of $R\Gamma_c(G, b, \mu)[\rho]$. As a first step towards this goal, we would like to prove the existence of isomorphisms

$$\mathbb{D}_{J_b}(R\Gamma_c(G, b, \mu)[\pi]) \simeq R\Gamma_c(G, b, \mu)[\mathbb{D}_G(\pi)]$$

and

$$\mathbb{D}_G(R\Gamma_c(G, b, \mu)[\rho]) \simeq R\Gamma_c(G, b, \mu)[\mathbb{D}_{J_b}(\rho)]$$

as complexes of $J_b(\mathbb{Q}_p) \times W_E$ and $G(\mathbb{Q}_p) \times W_E$ -modules, respectively.

In general, this shouldn't be true, but, under a Hodge-Newton reducibility condition, we show that it is. In particular, our main theorem is as follows.

Theorem 1.1. *For a basic local Shimura datum (G, b, μ) such that the set $B(G, \mu)$ is Hodge-Newton reducible (Definition 2.2) and π (resp. ρ) a supercuspidal representation of $G(\mathbb{Q}_p)$ (resp. $J_b(\mathbb{Q}_p)$), we have isomorphisms*

$$\mathbb{D}_{J_b}(R\Gamma_c(G, b, \mu)[\pi]) \simeq R\Gamma_c(G, b, \mu)[\mathbb{D}_G(\pi)]$$

and

$$\mathbb{D}_G(R\Gamma_c(G, b, \mu)[\rho]) \simeq R\Gamma_c(G, b, \mu)[\mathbb{D}_{J_b}(\rho)]$$

as complexes of $J_b(\mathbb{Q}_p) \times W_E$ and $G(\mathbb{Q}_p) \times W_E$ -modules, respectively.

Remark 1.1. A similar result should be provable through our methods with integral or torsion coefficients, for ρ any supercuspidal ℓ -complete or ℓ -modular representation, respectively. We note that in the above we have assumed μ is minuscule; however, we conjecture that this is true even in the non-minuscule case.

The condition that the set $B(G, \mu)$ is Hodge-Newton reducible roughly says that, for every non-basic element $b' \in B(G, \mu)$, the Newton polygon of b' seen as an element of a positive Weyl chamber touches the Hodge polygon defined by μ away from the endpoints. We note that there is a complete group-theoretic classification of pairs (G, μ) such that $B(G, \mu)$ is Hodge-Newton reducible [GHN19, Theorem 2.5].

We explain the idea of proof for the π -isotypic part with the proof for the ρ -isotypic part being analogous. The key insight is to consider the geometric analogue of the Zelevinsky involution acting on sheaves on the moduli stack Bun_G

$$\mathbb{D}_{BZ} : \text{D}_{\text{lis}}(\text{Bun}_G, \overline{\mathbb{Q}}_\ell)^{\omega, \text{op}} \rightarrow \text{D}_{\text{lis}}(\text{Bun}_G, \overline{\mathbb{Q}}_\ell)^\omega$$

as defined in [FS21, Section V.5, Section VII.7]. It defines a contravariant equivalence of $\text{D}_{\text{lis}}(\text{Bun}_G, \overline{\mathbb{Q}}_\ell)^\omega$ and has the property that \mathbb{D}_{BZ}^2 is naturally isomorphic to the identity. Moreover, when restricted to the subcategories $\text{D}(J_b(\mathbb{Q}_p), \overline{\mathbb{Q}}_\ell)^\omega$ coming from basic Harder-Narasimhan (=HN)-strata of Bun_G , it is given by \mathbb{D}_{J_b} . Now, with this in hand, it is useful to reinterpret the complex $R\Gamma_c(G, b, \mu)[\pi]$ as

$$j_b^* T_{\mu^{-1}} j_{1!}(\mathcal{F}_\pi) \simeq R\Gamma_c(G, b, \mu)[\pi]$$

where j_1 and j_b are the inclusions of the HN-strata of Bun_G corresponding to the trivial element $1 \in B(G)$ and $b \in B(G)$, respectively, $\mathcal{F}_\pi \in \text{D}(G(\mathbb{Q}_p), \overline{\mathbb{Q}}_\ell)$ is the sheaf on the basic HN-stratum $\text{Bun}_G^1 \simeq [*/G(\mathbb{Q}_p)]$ defined by π , and $T_{\mu^{-1}}$ is the Hecke operator associated to μ^{-1} a dominant inverse of μ acting on $\text{D}_{\text{lis}}(\text{Bun}_G, \overline{\mathbb{Q}}_\ell)$. Given this isomorphism, Theorem 1.1 would now follow from knowing that \mathbb{D}_{BZ} commutes with all operations applied to \mathcal{F}_π on the LHS. For j_b and the Hecke operator $T_{\mu^{-1}}$, this is shown by Fargues-Scholze. The key point is to show that it commutes with the restriction functor j_1^* . In general, this is not true if the complex $T_{\mu^{-1}} j_{1!}(\mathcal{F}_\pi)$ is supported on non-basic HN-strata, but in the case that $B(G, \mu)$ is Hodge-Newton reducible and π is supercuspidal we show that this cannot occur. The key point is that the restriction of $T_{\mu^{-1}} j_{1!}(\mathcal{F}_\pi)$ to any non-basic HN-strata can be described in terms of the π -isotypic part of $R\Gamma_c(G, b', \mu)$ for non-basic $b' \in B(G, \mu)$. However, by Hodge-Newton reducibility, the space $\text{Sht}(G, b', \mu)_\infty$ is parabolically induced as a space with $G(\mathbb{Q}_p)$ -action from a proper parabolic of G , realizing some form of the Harris-Viehmann conjecture [RV14, Conjecture 8.5]. Therefore, since π is supercuspidal, it cannot contribute to the cohomology of these spaces, so the restriction vanishes.

With Theorem 1.1 in hand, we can start to derive some useful cohomological consequences. Let χ denote the central character of our fixed supercuspidal π . Then the complex $R\Gamma_c(G, b, \mu)[\pi]$ has cohomology valued in $J_b(\mathbb{Q}_p)$ -representations with central character equal to χ , where we recall that J_b is an extended pure inner form of G and therefore we have an isomorphism: $Z(J_b) \simeq Z(G)$. Let $I_\chi(J_b)$ denote the set of equivalence classes of inertial cuspidal data $\mathfrak{s} = (M, \sigma)$, where $M \subset J_b$ is a Levi subgroup of J_b and σ is a supercuspidal representation with $\sigma|_{Z(G)(\mathbb{Q}_p)} = \chi$. We can consider the Bernstein decomposition

$$R\Gamma_c(G, b, \mu)[\pi] \simeq \bigoplus_{\mathfrak{s} \in I_\chi(G)} R\Gamma_c(G, b, \mu)[\pi]_{\mathfrak{s}}$$

where the cohomology of $R\Gamma_c(G, b, \mu)[\pi]_{\mathfrak{s}}$ takes values in representations with supercuspidal support specified by \mathfrak{s} . On the other hand, since π is supercuspidal, $R\Gamma_c(G, b, \mu)[\mathbb{D}_{J_b}(\pi)]$ is, up to a shift, isomorphic to $R\Gamma_c(G, b, \mu)[\pi^*]$, where π^* is the contragradient of π . Therefore, this complex takes values in representations with central character equal to χ^{-1} . Hence, we again obtain a Bernstein decomposition:

$$R\Gamma_c(G, b, \mu)[\pi^*] \simeq \bigoplus_{\mathfrak{s} \in I_{\chi^{-1}}(J_b)} R\Gamma_c(G, b, \mu)[\pi]_{\mathfrak{s}}$$

Fixing now $\mathfrak{s} = (M, \sigma) \in I_{\chi}(J_b)$, we let $\mathfrak{s}^* = (M, \sigma^*) \in I_{\chi^{-1}}(J_b)$. Using Theorem 1.1, one can show that one has (up to a shifts) an isomorphism

$$\mathbb{D}_{J_b}(R\Gamma_c(G, b, \mu)[\pi]_{\mathfrak{s}}) \simeq R\Gamma_c(G, b, \mu)[\pi^*]_{\mathfrak{s}^*}$$

of complexes of $J_b(\mathbb{Q}_p) \times W_E$ -modules. To a given \mathfrak{s} , we let $i(\mathfrak{s}) := r_G - r_M$, where r_G (resp. G) is the split semisimple rank of G (resp. M). If one has an irreducible smooth representation ρ , then the cohomology of $\mathbb{D}_{J_b}(\rho)$ sits in a single degree determined by $i(\mathfrak{s})$. We can define $Zel(\rho)$ (Definition 3.1) to be the representation determined by the non-zero cohomology. As a consequence of this, we can deduce the following result from Theorem 1.1 by passing to cohomology.

Theorem 1.2. *For a basic local Shimura datum (G, b, μ) such that $B(G, \mu)$ is Hodge-Newton reducible, let π be a supercuspidal representation of $G(\mathbb{Q}_p)$ with central character χ , $\mathfrak{s} \in I_{\chi}(J_b)$ with associated integer $i(\mathfrak{s})$, ρ a smooth irreducible representation with supercuspidal support given by \mathfrak{s} , and σ an irreducible representation of W_E . Then $\pi \boxtimes \rho \boxtimes \sigma$ occurs as a subquotient of $H^q(R\Gamma_c(G, b, \mu))$ if and only if $\pi^* \boxtimes Zel(\rho) \boxtimes \sigma^*$ occurs as a subquotient of $H^{-q+i(\mathfrak{s})}(R\Gamma_c(G, b, \mu))$. The analogous statement is true with the roles of π and ρ interchanged.*

Remark 1.2. (1) Using this result, one should be able to make some more precise claims about the cohomological degrees that the complex $R\Gamma_c(G, b, \mu)[\pi]$ (resp. $R\Gamma_c(G, b, \mu)[\rho]$) sits in for π (resp. ρ) supercuspidal. For example, conjecturally the spaces $\text{Sht}(G, b, \mu)_K$ should be Stein (see [Han20, Section 1.3] for a discussion of this) in the sense that the complex $R\Gamma_c(\text{Sht}(G, b, \mu)_K, \mathcal{S}_{\mu})$ should be concentrated in degrees $0 \leq i \leq d$.¹ As a consequence, since π (resp. ρ) supercuspidal is injective/projective in the category of representations with fixed central character, we know that $R\Gamma_c(G, b, \mu)[\pi]$ (resp. $R\Gamma_c(G, b, \mu)[\rho]$) is concentrated in degrees $0 \leq i \leq d$, but then, by Theorem 1.2, this is enough to conclude that $R\Gamma_c(G, b, \mu)[\pi]_{\mathfrak{s}}$ (resp. $R\Gamma_c(G, b, \mu)[\rho]_{\mathfrak{s}}$) can only be concentrated in degrees $0 \leq i \leq i(\mathfrak{s})$.

- (2) In the case that L/\mathbb{Q}_p is a finite extension, $G = \text{Res}_{L/\mathbb{Q}_p} \text{GL}_n$ is the restriction of scalars of GL_n , and μ the Lubin-Tate cocharacter, this is a result of Fargues [Far06, Theorem 4.6]. Similarly, for $G = \text{GSp}_4/\mathbb{Q}_p$, and μ the Siegel cocharacter, this is a result of Mieda [Mie14, Corollary 1.2 (ii)].
- (3) Unlike the arguments of Fargues and Mieda, we completely avoid considering integral models of the local Shimura varieties, working solely on the generic fiber. This substantially simplifies the proof, allowing one to completely avoid using nearby cycles and studying the special fiber, albeit at the cost of using much more sophisticated geometric tools.
- (4) In the case that $G = \text{GSp}_4/\mathbb{Q}_p$, $\mu = (1, 1, 0, 0)$, ϕ is a discrete (not necessarily supercuspidal) L -parameter with associated A -parameter ψ as above, and ρ a (not necessarily supercuspidal) representation in the L -packet or A -packet over ϕ or ψ , respectively, Ito and Mieda [IM22] have leveraged this result in conjunction with basic uniformization and global results to fully understand the cohomology of the summand $R\Gamma_c(G, b, \mu)[\rho]_{sc}$ of $R\Gamma_c(G, b, \mu)[\rho]$, where $G(\mathbb{Q}_p)$ acts via a supercuspidal representation, in terms of members of the L -packet and A -packet $\Pi_{\phi}(G)$ and $\Pi_{\psi}(G)$. It would be interesting to carry out their analysis in other Hodge-Newton reducible situations. For example, in the case where $G = \text{GU}(1, n-1)/\mathbb{Q}_p$

¹We note that in the minuscule case $\mathcal{S}_{\mu} \simeq \mathbb{Z}_{\ell}[d](\frac{d}{2})$.

is an unramified unitary group and $\mu = (1, 0, \dots, 0, 0)$. The form of the cohomology in the case that $n = 3$ is partially described by [Ito13].

In section 2, we discuss Hodge-Newton reducibility and prove Theorem 1.1. In section 3, we discuss the Zelevinsky involution in greater detail and show how to deduce Theorem 1.2 from Theorem 1.1.

2. PROOF OF THE MAIN THEOREM

In this section, we prove Theorem 1.1. We first address the notion of Hodge-Newton reducibility and derive a key geometric consequence. We recall that the Kottwitz set $B(G)$ is equipped with two maps.

- The slope homomorphism

$$\begin{aligned} \nu : B(G) &\rightarrow X_*(T_{\overline{\mathbb{Q}_p}})_{\mathbb{Q}}^{+, \Gamma} \\ b &\mapsto \nu_b \end{aligned}$$

where $\Gamma := \text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$ and $X_*(T_{\overline{\mathbb{Q}_p}})_{\mathbb{Q}}^{+}$ is the set of rational dominant cocharacters of G .

- The Kottwitz invariant

$$\kappa : B(G) \rightarrow \pi_1(G)_{\Gamma}$$

where $\pi_1(G) = X_*(T_{\overline{\mathbb{Q}_p}})/X_*(T_{\overline{\mathbb{Q}_p, sc}})$ for $T_{\overline{\mathbb{Q}_p, sc}}$ the maximal simply connected subtorus of $T_{\overline{\mathbb{Q}_p}}$.

Now, given a geometric cocharacter μ of G with reflex field E , we can define the element:

$$\tilde{\mu} := \frac{1}{[E : \mathbb{Q}_p]} \sum_{\gamma \in \text{Gal}(E/\mathbb{Q}_p)} \gamma(\mu) \in X_*(T_{\overline{\mathbb{Q}_p}})_{\mathbb{Q}}^{+, \Gamma}$$

We let μ^b be the image of μ in $\pi_1(G)_{\Gamma}$. This allows us to make the following definition.

Definition 2.1. Let b be an element in the Kottwitz set $B(G)$ of G , we define the set $B(G, \mu, b)$ to be set of $b' \in B(G)$ for which $\nu_{b'} - \nu_b \leq \tilde{\mu}$ with respect to the Bruhat ordering and $\kappa(b') - \kappa(b) = \mu^b$. If b is the trivial element we will denote this by $B(G, \mu)$.

With this in hand, we can define the relevant notion of Hodge-Newton reducibility.

Definition 2.2. For $b \in B(G)$, we say that $B(G, \mu, b)$ is Hodge-Newton reducible if, for all non-basic $b' \in B(G, \mu, b)$, there exists a proper Levi subgroup $M \subset G$ and $b_M, b'_M \in B(M)$ such that:

- b and b' are in the images of b_M and b'_M under the natural map $B(M) \rightarrow B(G)$, respectively,
- $b_M \in B(M, \mu, b_M)$, for some fixed choice of conjugacy class of μ regarded as a cocharacter inside M^2 .

Remark 2.1. We note that, since we assumed G is quasi-split, by [CFS21, Lemma 4.11], $B(G, \mu)$ being Hodge-Newton reducible is equivalent to saying that $B(G, \mu)$ is fully Hodge-Newton decomposable in the sense of [CFS21, Definition 4.10]. If $b \in B(G, \mu)$ denotes the unique basic element and μ^{-1} is a dominant inverse of μ then it follows that, if $B(G, \mu)$ is Hodge-Newton reducible, then $B(G, \mu^{-1}, b)$ is also Hodge-Newton reducible in our sense by [CFS21, Corollary 4.15] and [CFS21, Lemma 4.11] (See the first part of the proof of [CFS21, Theorem 6.1] for details).

²We note that the conjugacy class $\{\mu\}$ does not define a unique conjugacy class inside the Levi M , and it is important to fix some choice.

We now seek to derive a key geometric consequence of this condition. Fix an element $b \in B(G)$ and let $b' \in B(G, \mu, b)$. We define the Shtuka space

$$\mathrm{Sht}(G, b, b', \mu)_\infty \rightarrow \mathrm{Spd}(\check{E})$$

to be the space parametrizing modifications

$$\mathcal{E}_{b'} \dashrightarrow \mathcal{E}_b$$

of G -bundles on the Fargues-Fontaine curve X with meromorphy bounded by μ . This has commuting actions of $J_b(\mathbb{Q}_p)$ and $J_{b'}(\mathbb{Q}_p)$ coming from automorphisms of \mathcal{E}_b and $\mathcal{E}_{b'}$, respectively. We define as before the tower

$$\mathrm{Sht}(G, b, b', \mu)_K := \mathrm{Sht}(G, b, b', \mu)/K$$

for varying open compact subgroups $K \subset J_b(\mathbb{Q}_p)$. Assume that b is basic from now on. Then J_b is an extended pure inner form of G , it follows that one has an isomorphism $\mathrm{Bun}_{G, \bar{\mathbb{F}}_p} \simeq \mathrm{Bun}_{J_b, \bar{\mathbb{F}}_p}$ by mapping a G -bundle \mathcal{E} on X to the J_b -torsor of isomorphisms $\mathcal{I}som(\mathcal{E}_b, \mathcal{E})$. This implies that the space of Shtukas $\mathrm{Sht}(G, b, b', \mu)_\infty$ is isomorphic to the space of Shtukas $\mathrm{Sht}(J_b, b'b^{-1}, \mu)_\infty$ for the group J_b . This isomorphism means that the cocharacter μ and the Geometric Satake correspondence for J_b [FS21, Chapter VI] will give rise to a \mathbb{Z}_ℓ -sheaf \mathcal{S}_μ on the space $\mathrm{Sht}(J_b, b'b^{-1}, \mu)_\infty$ and in turn on $\mathrm{Sht}(G, b, b', \mu)_\infty$ compatible with the actions of $J_b(\mathbb{Q}_p)$ and $J_{b'}(\mathbb{Q}_p)$. This allows us to define the complex

$$R\Gamma_c(G, b, b', \mu) := \mathrm{colim}_{K \rightarrow \{1\}} R\Gamma_c(\mathrm{Sht}(G, b, b', \mu)_{K, \mathbb{C}_p}, \mathcal{S}_\mu) \otimes \bar{\mathbb{Q}}_\ell$$

of $J_b(\mathbb{Q}_p) \times J_{b'}(\mathbb{Q}_p) \times W_E$ -modules, where $\mathrm{Sht}(G, b, b', \mu)_{K, \mathbb{C}_p}$ is the base change of $\mathrm{Sht}(G, b, b', \mu)_K$ to \mathbb{C}_p . We have the following key result concerning this complex.

Proposition 2.1. *Suppose that μ is minuscule and $B(G, \mu, b)$ is Hodge-Newton reducible and let $b' \in B(G, \mu, b)$ be a non-basic element, then the complex $R\Gamma_c(G, b, b', \mu)$ is parabolically induced as a complex of $J_b(\mathbb{Q}_p)$ -modules.*

Proof. In the case that μ is minuscule, we have an isomorphism: $\mathcal{S}_\mu \simeq \mathbb{Z}_\ell[d](\frac{d}{2})$. It follows from [GI16, Theorem 4.23] that the cohomology of $R\Gamma_c(G, b, b', \mu)$ is parabolically induced as a $J_b(\mathbb{Q}_p)$ -module. However, since parabolic induction is exact by second adjointness, it follows that this is also true at the level of complexes. \square

To μ and its dominant inverse μ^{-1} , we consider the Hecke operators

$$T_\mu, T_{\mu^{-1}} : \mathrm{D}_{\mathrm{lis}}(\mathrm{Bun}_G, \bar{\mathbb{Q}}_\ell) \rightarrow \mathrm{D}_{\mathrm{lis}}(\mathrm{Bun}_G, \bar{\mathbb{Q}}_\ell)^{BW_E}$$

defined by mutually dual highest weight representations of highest weight μ and μ^{-1} of the Langlands dual group, as defined in [FS21, Section IX.2]. We can deduce the following consequence for the support of these Hecke operators.

Corollary 2.2. *For a basic local Shimura datum (G, b, μ) such that $B(G, \mu)$ is Hodge-Newton reducible and π (resp. ρ) a supercuspidal representation of $G(\mathbb{Q}_p)$ (resp. $J_b(\mathbb{Q}_p)$), the objects*

$$T_{\mu^{-1}} j_{1!}(\mathcal{F}_\pi)$$

and

$$T_\mu j_{b!}(\mathcal{F}_\rho)$$

in $\mathrm{D}_{\mathrm{lis}}(\mathrm{Bun}_G, \bar{\mathbb{Q}}_\ell)^{BW_E}$ only have non-zero restriction to the open HN-stratum Bun_G^b and Bun_G^1 defined by b and the trivial element 1, respectively. Here $j_b : \mathrm{Bun}_G^b \hookrightarrow \mathrm{Bun}_G$ and $j_1 : \mathrm{Bun}_G^1 \hookrightarrow \mathrm{Bun}_G$ are open immersions of the HN-strata defined by b and 1, respectively, and \mathcal{F}_π (resp. \mathcal{F}_ρ) are the lisse-étale $\bar{\mathbb{Q}}_\ell$ -sheaves defined by π (resp. ρ) on Bun_G^1 (resp. Bun_G^b).

Proof. We start with the statement for π . It follows by [Rap18, Proposition A.9] that the locus where $T_{\mu^{-1}}j_{1!}(\mathcal{F}_\pi)$ is supported is precisely $B(G, \mu) \subset B(G) \simeq |\text{Bun}_G|$, where $|\text{Bun}_G|$ is the underlying topological space of Bun_G . For any $b' \in B(G)$ it follows by the analysis in [FS21, Section IX.3] that we have an isomorphism (up to shifts and twists)

$$j_{b'}^* T_{\mu^{-1}} j_{1!}(\mathcal{F}_\pi) \simeq R\Gamma_c(G, b', \mu)[\pi]$$

as complexes of $J_{b'}(\mathbb{Q}_p) \times W_E$ -modules. However, by Proposition 2.1, we know that the complex $R\Gamma_c(G, b', \mu)$ is parabolically induced as a $G(\mathbb{Q}_p)$ -module. Therefore, since π is supercuspidal, it follows that $R\Gamma_c(G, b', \mu)[\pi] := R\Gamma_c(G, b, \mu) \otimes^{\mathbb{L}} \pi$ must be trivial, by the adjunction between the Jacquet module and parabolic induction in the derived category. For ρ , since b is basic, the sheaf

$$T_\mu j_{b!}(\mathcal{F}_\rho)$$

will be supported on $B(G, \mu^{-1}, b)$ by [Rap18, Corollary A.10]. It then follows by [CFS21, Corollary 4.15] that this is also Hodge-Newton reducible, as in Remark 2.1. The restrictions of $T_\mu j_{b!}(\mathcal{F}_\rho)$ to all non-basic HN-stratum $b' \in B(G, \mu^{-1}, b)$ are isomorphic to $R\Gamma_c(G, b, b', \mu^{-1}) \otimes_{\mathcal{H}(J_b)}^{\mathbb{L}} \rho$. However, by Proposition 2.1, we know that the cohomology of $R\Gamma_c(G, b, b', \mu^{-1})$ is parabolically induced as a $J_b(\mathbb{Q}_p)$ -module and therefore the restriction vanishes, since ρ is supercuspidal. \square

From this, we conclude Theorem 1.1.

Theorem 2.3. *For a basic local Shimura datum (G, b, μ) such that $B(G, \mu)$ is Hodge-Newton reducible and π (resp. ρ) a supercuspidal representation of $G(\mathbb{Q}_p)$ (resp. $J_b(\mathbb{Q}_p)$), we have isomorphisms*

$$\mathbb{D}_{J_b}(R\Gamma_c(G, b, \mu)[\pi]) \simeq R\Gamma_c(G, b, \mu)[\mathbb{D}_G(\pi)]$$

and

$$\mathbb{D}_G(R\Gamma_c(G, b, \mu)[\rho]) \simeq R\Gamma_c(G, b, \mu)[\mathbb{D}_{J_b}(\rho)]$$

as complexes of $J_b(\mathbb{Q}_p) \times W_E$ and $G(\mathbb{Q}_p) \times W_E$ -modules, respectively.

Proof. We explain the proof for the π -isotypic part, with the proof for the ρ -isotypic part being analogous. As in the previous proof, we start with the isomorphism

$$j_b^* T_{\mu^{-1}} j_{1!}(\mathcal{F}_\pi) \simeq R\Gamma_c(G, b, \mu)[\pi]$$

of complexes of $J_b(\mathbb{Q}_p) \times W_E$ -modules. Acting by \mathbb{D}_{J_b} , we have an isomorphism:

$$\mathbb{D}_{J_b} j_b^*(T_{\mu^{-1}} j_{1!}(\mathcal{F}_\pi)) \simeq \mathbb{D}_{J_b} R\Gamma_c(G, b, \mu)[\pi]$$

Let $\mathbb{D}_{BZ} : \text{D}_{\text{lis}}(\text{Bun}_G, \overline{\mathbb{Q}}_\ell)^{\omega, \text{op}} \rightarrow \text{D}_{\text{lis}}(\text{Bun}_G, \overline{\mathbb{Q}}_\ell)^\omega$ be the geometric incarnation of the Zelevinsky involution, as defined in [FS21, Section V.5, Section VII.7]. By Corollary 2.2 and [FS21, Proposition VII.7.6], it follows that LHS of the previous equation is isomorphic to:

$$j_b^*(\mathbb{D}_{BZ} T_{\mu^{-1}} j_{1!}(\mathcal{F}_\pi))$$

Now, by [FS21, Theorem I.7.2] and [FS21, Proposition VII.7.6], \mathbb{D}_{BZ} also commutes with the Hecke operator $T_{\mu^{-1}}$ and acts on $j_{1!}(\mathcal{F}_\pi)$ via \mathbb{D}_G , respectively; therefore, we can rewrite the LHS as

$$j_b^* T_{\mu^{-1}} j_{1!}(\mathcal{F}_{\mathbb{D}_G(\pi)})$$

which, as before, is isomorphic to

$$R\Gamma_c(G, b, \mu)[\mathbb{D}_G(\pi)]$$

as a complex of $J_b(\mathbb{Q}_p) \times W_E$ -modules. All in all, we obtain the desired identification. \square

3. APPLICATIONS

We start by recalling some facts about the Zelevinsky involution. Set r to be the split rank of the center of G . We let $\chi : Z(G) \rightarrow \overline{\mathbb{Q}}_\ell^*$ be a central character and $I_\chi(G)$ be the set of equivalence classes of inertial types (M, σ) , where M is a proper Levi subgroup of G and σ is a supercuspidal representation of M with $\sigma|_{Z(G)} \simeq \chi$. We write $\text{Rep}_\chi(G)$ for the category of finite-length smooth representations with central character equal to χ . We consider the Bernstein-decomposition [Ren10, Theorem VI.7.2]:

$$\text{Rep}_\chi(G) = \bigoplus_{\mathfrak{s} \in I_\chi(G)} \text{Rep}_\mathfrak{s}(G)$$

For $\mathfrak{s} \in I_\chi(G)$, we let $i(\mathfrak{s})$ be the integer defined in Section 1. We then have the following key result of Schneider-Stuhler.

Proposition 3.1. [SS97, Theorem III.3.1] *For $\mathfrak{s} \in I_\chi(G)$ and $\pi \in \text{Rep}_\mathfrak{s}(G)$ viewed as an element of $D(G(\mathbb{Q}_p), \overline{\mathbb{Q}}_\ell)$ sitting in degree 0, the cohomology of the complex $\mathbb{D}_G(\pi)$ is concentrated only in degrees $-i(\mathfrak{s}) + r$. Moreover, if $\mathfrak{s}^* := (M, \sigma^*) \in I_{\chi^{-1}}(G)$ then it takes values in the Bernstein component $\text{Rep}_{\mathfrak{s}^*}(G)$ of $\text{Rep}_{\chi^{-1}}(G)$.*

This gives us the following key definition.

Definition 3.1. For $\pi \in \text{Rep}_\mathfrak{s}(G)$ as above, we define $Zel(\pi) \in \text{Rep}_{\mathfrak{s}^*}(G)$ to be the $-i(\mathfrak{s}) + r$ -cohomology of $\mathbb{D}_G(\pi)$. When pre-composed with the contragradient map, $Zel(-)$ defines an exact categorical equivalence of $\text{Rep}_\mathfrak{s}(G)$. In particular, it takes irreducible objects to irreducible objects.

One can verify Proposition 3.1 by computing \mathbb{D}_G explicitly in terms of the following Deligne-Luzstig complex

$$\text{DL}(\pi) := 0 \rightarrow \pi \rightarrow \bigoplus_P i_P^G r_P^G(\pi) \rightarrow \dots \rightarrow i_B^G r_B^G(\pi) \rightarrow 0$$

where i_P^G and r_P^G are the parabolic induction and Jacquet functors associated to a parabolic subgroup P , and the i th term of the complex runs over conjugacy classes of parabolics of corank i . The differentials of the complex are an appropriate alternating sum of adjunction maps (via transitivity of parabolic induction). If one has a complex $A \in D(G(\mathbb{Q}_p), \overline{\mathbb{Q}}_\ell)$ then one gets a bi-complex $\text{DL}(A)$, and thereby a functor $\text{DL} : D(G(\mathbb{Q}_p), \overline{\mathbb{Q}}_\ell) \rightarrow D(G(\mathbb{Q}_p), \overline{\mathbb{Q}}_\ell)$. We have the following theorem.

Theorem 3.2. [SS97; R04; BBK18] *For $A \in D(G(\mathbb{Q}_p), \overline{\mathbb{Q}}_\ell)$ with admissible cohomology we have a quasi-isomorphism*

$$\text{DL}(A^*) \simeq \mathbb{D}_G(A)[r]$$

where A^* denotes the smooth contragradient.

For example, using this, we can immediately deduce the following lemma.

Lemma 3.3. *Let π be a supercuspidal representation of $G(\mathbb{Q}_p)$ then we have an isomorphism*

$$\mathbb{D}_G(\pi) \simeq \pi^*[-r]$$

as $G(\mathbb{Q}_p)$ -modules.

Now let's consider the isomorphism supplied by Theorem 1.1

$$(1) \quad \mathbb{D}_{J_b}(R\Gamma_c(G, b, \mu)[\pi]) \simeq R\Gamma_c(G, b, \mu)[\mathbb{D}_G(\pi)] \simeq R\Gamma_c(G, b, \mu)[\pi^*][-r]$$

where we have used Lemma 3.3. Now if χ denotes the central character of π it follows that all smooth irreducible $J_b(\mathbb{Q}_p)$ -representations occurring in the complex $R\Gamma_c(G, b, \mu)[\pi]$ (resp. $R\Gamma_c(G, b, \mu)[\pi^*]$) have central character χ (resp. χ^{-1}). This follows since the diagonally embedded center of

$Z(J_b)(\mathbb{Q}_p) \simeq Z(G)(\mathbb{Q}_p)$ acts trivially on the space of modifications $\mathcal{E}_b \dashrightarrow \mathcal{E}_0$ with its $J_b(\mathbb{Q}_p) \times G(\mathbb{Q}_p)$ -action by automorphisms of bundles. Moreover, we know both sides of (1) take values in admissible representations of finite length, therefore we obtain a Bernstein decomposition of both sides:

$$\bigoplus_{\mathfrak{s} \in I_\chi(J_b)} \mathbb{D}_{J_b}(R\Gamma_c(G, b, \mu)[\pi]_{\mathfrak{s}}) \simeq \bigoplus_{\mathfrak{s} \in I_{\chi^{-1}}(J_b)} R\Gamma_c(G, b, \mu)[\pi^*]_{\mathfrak{s}^*}[-r]$$

Then, by Proposition 3.1, we see that, for a fixed $\mathfrak{s} = (M, \sigma) \in I_\chi(J_b)$ with $\mathfrak{s}^* = (M, \sigma^*) \in I_{\chi^{-1}}(J_b)$, this induces an isomorphism

$$\mathbb{D}_{J_b}(R\Gamma_c(G, b, \mu)[\pi]_{\mathfrak{s}}) \simeq R\Gamma_c(G, b, \mu)[\pi^*]_{\mathfrak{s}^*}[-r]$$

of $J_b(\mathbb{Q}_p) \times W_E$ -modules. Similarly, we deduce such an isomorphism for the ρ -isotypic part. We record this as a Corollary now.

Corollary 3.4. *For a basic local Shimura datum (G, b, μ) such that $B(G, \mu)$ is Hodge-Newton reducible and π (resp. ρ) a supercuspidal representation of $G(\mathbb{Q}_p)$ (resp. $J_b(\mathbb{Q}_p)$) with central character χ , we have, for all $\mathfrak{s} = (M, \sigma) \in I_\chi(J_b)$ (resp. $I_\chi(G)$) with $\mathfrak{s}^* = (M, \sigma^\vee) \in I_{\chi^{-1}}(J_b)$ (resp. $I_{\chi^{-1}}(G)$), isomorphisms*

$$\mathbb{D}_{J_b}(R\Gamma_c(G, b, \mu)[\pi]_{\mathfrak{s}}) \simeq R\Gamma_c(G, b, \mu)[\pi^*]_{\mathfrak{s}^*}[-r]$$

and

$$\mathbb{D}_G(R\Gamma_c(G, b, \mu)[\rho]_{\mathfrak{s}}) \simeq R\Gamma_c(G, b, \mu)[\rho^*]_{\mathfrak{s}^*}[-r]$$

of $J_b(\mathbb{Q}_p) \times W_E$ and $G(\mathbb{Q}_p) \times W_E$ -modules, respectively.

Now, with this in hand, we can deduce Theorem 1.2 by passing to cohomology.

Theorem 3.5. *For a basic local Shimura datum (G, b, μ) such that $B(G, \mu)$ is Hodge-Newton reducible, let π a supercuspidal representation of $G(\mathbb{Q}_p)$ with central character χ , $\mathfrak{s} \in I_\chi(J_b)$ with associated integer $i(\mathfrak{s})$, ρ a smooth irreducible representation with supercuspidal support given by \mathfrak{s} , and σ an irreducible representation of W_E . Then $\pi \boxtimes \rho \boxtimes \sigma$ occurs as a subquotient of $H^q(R\Gamma_c(G, b, \mu))$ if and only if $\pi^* \boxtimes \text{Zel}(\rho) \boxtimes \sigma^*$ occurs as a subquotient of $H^{-q+i(\mathfrak{s})}(R\Gamma_c(G, b, \mu))$. The analogous statement is true with the roles of π and ρ interchanged.*

Proof. We explain the proof for π supercuspidal with the proof for ρ supercuspidal being the same. We apply Corollary 3.4 to the contragradient π^* and the Bernstein-component \mathfrak{s}^* , this gives us an identification:

$$\mathbb{D}_{J_b}(R\Gamma_c(G, b, \mu)[\pi^*]_{\mathfrak{s}^*}) \simeq R\Gamma_c(G, b, \mu)[\pi]_{\mathfrak{s}}[-r]$$

Now, since π is supercuspidal and therefore injective/projective in the category of representations with fixed central character, it follows that $\pi \boxtimes \rho \boxtimes \sigma$ occurring as a subquotient of $H^q(R\Gamma_c(G, b, \mu))$ is equivalent to $\rho \boxtimes \sigma$ occurring as a sub-quotient of $H^q(R\Gamma_c(G, b, \mu)[\pi])$. By the above isomorphism, this is equivalent to $\rho \boxtimes \sigma$ occurring as a sub-quotient of $H^{q+r}(\mathbb{D}_{J_b}(R\Gamma_c(G, b, \mu)[\pi^*]_{\mathfrak{s}^*}))$. However, since \mathbb{D}_{J_b} is contravariant, it follows, by Proposition 3.1, that this occurs if and only if $\text{Zel}(\rho) \boxtimes \sigma^*$ occurs as a sub-quotient of $H^{-q+i(\mathfrak{s})}(R\Gamma_c(G, b, \mu)[\pi^*])$, which is in turn equivalent to $\pi^* \boxtimes \text{Zel}(\rho) \boxtimes \sigma^*$ occurring as a sub-quotient of $H^{-q+i(\mathfrak{s})}(R\Gamma_c(G, b, \mu))$, as desired. \square

REFERENCES

- [BBK18] Joseph Bernstein, Roman Bezrukavnikov, and David Kazhdan. “Deligne-Lusztig duality and wonderful compactification”. In: *Selecta Math. (N.S.)* 24.1 (2018), pp. 7–20. ISSN: 1022-1824. DOI: 10.1007/s00029-018-0391-5. URL: <https://doi.org/10.1007/s00029-018-0391-5>.
- [CFS21] Miaofen Chen, Laurent Fargues, and Xu Shen. “On the structure of some p -adic period domains”. In: *Cambridge Journal of Mathematics* 9.1 (2021), pp. 213–267.

- [Dat06] Jean François Dat. “Espaces symétriques de Drinfeld et correspondance de Langlands locale”. In: *Ann. Sci. École Norm. Sup. (4)* 39.1 (2006), pp. 1–74. ISSN: 0012-9593. DOI: 10.1016/j.ansens.2005.11.002. URL: <https://doi.org/10.1016/j.ansens.2005.11.002>.
- [Far06] L. Fargues. “Dualité de Poincaré et involution de Zelevinsky dans la cohomologie étale équivariante des espaces analytiques rigides”. In: *Preprint* (2006). Available here.
- [FS21] L. Fargues and P. Scholze. “Geometrization of the local Langlands Correspondence”. In: *Preprint* (2021). arXiv:2102.13459.
- [GHN19] Ulrich Görtz, Xuhua He, and Sian Nie. “Fully Hodge-Newton decomposable Shimura varieties”. In: *Peking Math. J.* 2.2 (2019), pp. 99–154. ISSN: 2096-6075. DOI: 10.1007/s42543-019-00013-2. URL: <https://doi.org/10.1007/s42543-019-00013-2>.
- [GI16] I. Gaisan and N. Imai. “Non-semi-stable loci in Hecke stacks and Fargues’ Conjecture”. In: *Preprint* (2016). arXiv:1608.07446.
- [Han20] D. Hansen. “On the supercuspidal cohomology of basic local Shimura varieties”. In: (2020). Available at home page of first named author.
- [Hir04] Kaoru Hiraga. “On functoriality of Zelevinski involutions”. In: *Compos. Math.* 140.6 (2004), pp. 1625–1656. ISSN: 0010-437X. URL: <https://doi.org/10.1112/S0010437X04000892>.
- [IM22] Tetsushi Ito and Yoichi Mieda. “Local Saito-Kurokawa A -packets and ℓ -adic cohomology of Rapoport-Zink tower for GSp_4 ”. In: *Preparation* (2022).
- [Ito13] Tetsushi Ito. “Supercuspidal representations in the cohomology of the Rapoport-Zink space for the unitary group in three variables”. In: (2013). Available here.
- [Mie14] Y. Mieda. “Zelevinsky involution and ℓ -adic cohomology of the Rapoport-Zink tower”. In: *Preprint* (2014). arXiv:1401.5135.
- [R04] Bezrukavnikov R. “Homological properties of representations of p -adic groups related to geometry of the group at infinity”. In: *Preprint* (2004). arXiv:math/0406223.
- [Rap18] Michael Rapoport. “Appendix to On the p -adic cohomology of the Lubin-Tate tower”. In: *Ann. Sci. Éc. Norm. Supér. (4)* 51.4 (2018), pp. 811–863. ISSN: 0012-9593. DOI: 10.24033/asens.2367. URL: <https://doi.org/10.24033/asens.2367>.
- [Ren10] David Renard. *Représentations des groupes réductifs p -adiques*. Vol. 17. Cours Spécialisés [Specialized Courses]. Société Mathématique de France, Paris, 2010, pp. vi+332. ISBN: 978-2-85629-278-5.
- [RV14] Michael Rapoport and Eva Viehmann. “Towards a theory of local Shimura varieties”. In: *Münster J. Math.* 7.1 (2014), pp. 273–326. ISSN: 1867-5778.
- [SS91] P. Schneider and U. Stuhler. “The cohomology of p -adic symmetric spaces”. In: *Invent. Math.* 105.1 (1991), pp. 47–122. ISSN: 0020-9910. DOI: 10.1007/BF01232257. URL: <https://doi.org/10.1007/BF01232257>.
- [SS97] Peter Schneider and Ulrich Stuhler. “Representation theory and sheaves on the Bruhat-Tits building”. In: *Inst. Hautes Études Sci. Publ. Math.* 85 (1997), pp. 97–191. ISSN: 0073-8301. URL: http://www.numdam.org/item?id=PMIHES_1997__85__97_0.
- [SW20] P. Scholze and J. Weinstein. *Berkeley Lectures on p -adic Geometry*. Vol. 389. Annals of Mathematics Studies. Princeton University Press, 2020.