

# DIRECT SUMMAND THEOREM BY $\text{arc}$ COHOMOLOGY

LONGKE TANG

The derived direct summand theorem is

**Theorem 1.** *Let  $R$  be a regular ring and  $f: X \rightarrow \text{Spec } R$  be proper surjective. Then the natural map  $f^*: R \rightarrow \text{R}\Gamma(X, \mathcal{O}_X)$  splits in  $\text{D}(R)$ .*

In this note we give a quick proof of it with the theory developed in [BS19].

*Proof.* We first make some reductions. Let  $C_f = \text{cofib}(f^*) \in \text{D}(R)$ , then  $f$  determines a class  $\alpha_f \in \text{Ext}^1(C_f, R)$ , and the theorem just says that  $\alpha_f = 0$ . From this we know that the theorem is fpqc-local, so we can assume  $R$  is a complete regular local ring with residue field  $k$  algebraically closed.

Now if  $\text{char } k = 0$ , then by Cohen structure theorem  $R = k[[x_1, \dots, x_n]]$ . By Artin–Popescu approximation,  $R$  is a colimit of smooth  $k$ -algebras; also  $k$  is a colimit of regular rings of finite type over  $\mathbb{Z}$ ; so  $R$  is a colimit of regular rings of finite type over  $\mathbb{Z}$ . The theorem can be obviously passed to limit, so it is reduced to the case that  $R$  is finite type over  $\mathbb{Z}$ ; but now all the maximal ideals of  $R$  has residue fields of positive characteristic, so we have reduced  $\text{char } k = 0$  case to  $\text{char } k = p$  case.

We also need some reductions about  $X$ . For  $g: Y \rightarrow X$  with  $fg$  surjective, one easily sees that the class  $\alpha_{fg}$  is carried to  $\alpha_f$  under the natural map  $C_f \rightarrow C_{fg}$ , so the theorem about  $fg$  implies that about  $f$ . By this we can assume that  $X$  is integral and  $f$  is generically finite. We can further assume that  $f$  is generically étale: if  $R = k[[x_1, \dots, x_n]]$  is of equal characteristic  $p$ , just base change to  $k[[x_1^{p^{-N}}, \dots, x_n^{p^{-N}}]]$  and then take the reduced scheme structure, for a sufficiently large  $N$ .

Now we arrive at the following situation:  $R$  is a complete regular local ring with residue field  $k$  algebraically closed with characteristic  $p$ .  $X$  is an integral scheme and  $f: X \rightarrow \text{Spec } R$  is proper surjective and generically étale. We need to prove that the class  $\alpha = \alpha_f \in \text{Ext}^1(C_f, R) = 0$ . By passage to limit one can take a nonzero  $r \in R$  so that  $X_{R_r} \rightarrow \text{Spec } R_r$  is already finite étale. Let  $X \rightarrow \text{Spec } S \rightarrow \text{Spec } R$  denote the Stein factorization of  $f$ ; then on  $\text{Spec } R_r$ , the first arrow is an isomorphism and the second is finite étale.

By Cohen structure theorem,  $R = W(k)[[x_1, \dots, x_n]]/E$  where  $E - p \in (x_1, \dots, x_n)$ . Let  $R_N = W(k)[[x_1^{p^{-N}}, \dots, x_n^{p^{-N}}]]/E$  and  $R_\infty = (\text{colim}_N R_N)^\wedge_p$ . Then  $R_\infty$  is a perfectoid ring  $p$ -completely faithfully flat over  $R$ . By [BS19, Theorem 7.14], there exists an absolutely integrally closed  $R'$  that is  $p$ -completely faithfully flat over  $R_\infty$ . Denote by  $X', S', f', \alpha'$  the base change to  $R'$  of  $X, S, f, \alpha$ , e.g.  $f'$  is the composition  $X' \rightarrow \text{Spec } S' \rightarrow \text{Spec } R'$ . Take compatible  $p^N$ -th roots of  $r$  in  $R'$  denoted  $r^{p^{-N}}$ , and let  $(r^{p^{-\infty}})$  denote the ideal generated by them. Then  $(R', (r^{p^{-\infty}}))$  is an almost setting, denoted  $r$ -almost.

I claim that it suffices to prove that  $\alpha'$  is  $r$ -almost zero. In fact, by the faithful flatness of  $R \rightarrow R'$  and Grothendieck coherence which implies that  $C_f \in \text{D}(R)$  is

bounded with finitely generated cohomology, we have

$$\mathrm{Ext}_{R'}^1(C_{f'}, R') = \mathrm{Ext}_R^1(C_f, R') = \mathrm{Ext}_R^1(C_f, R) \otimes_R R',$$

so  $\mathrm{Ann}_{R'}(\alpha') = \mathrm{Ann}_R(\alpha)R'$ . If  $\alpha'$  is  $r$ -almost zero, then  $\mathrm{Ann}_{R'}(\alpha') \supseteq (r^{p^{-\infty}})$ , which means that  $\mathrm{Ann}_{R'}(\alpha')^{p^N} \supseteq rR'$  and thus  $r \in \mathrm{Ann}_R(\alpha)^{p^N}$  for every  $N$ . Since  $r \neq 0$ , the Krull's intersection theorem forces  $\mathrm{Ann}_R(\alpha) = 1$ , which means  $\alpha = 0$ .

Finally we prove that  $\alpha'$  is  $r$ -almost zero, i.e.  $R' \rightarrow \mathrm{R}\Gamma(X', \mathcal{O}_{X'})$  splits in the category of  $p$ -complete  $r$ -almost  $R'$ -complexes. We use the  $p$ -complete arc topology. Since it is obviously finer than Zariski topology, we have a natural map  $\mathrm{R}\Gamma(X', \mathcal{O}_{X'}) \rightarrow \mathrm{R}\Gamma(X'_{\mathrm{arc}}, \mathcal{O}_{\mathrm{arc}})$ . Now it suffices to prove that  $R' \rightarrow \mathrm{R}\Gamma(X'_{\mathrm{arc}}, \mathcal{O}_{\mathrm{arc}})$  is  $p$ -completely  $r$ -almost split. Let  $I$  denotes the ideal  $(r^{p^{-\infty}})S'$ . At this stage, note that since  $X' \rightarrow \mathrm{Spec} S'$  is proper and an isomorphism outside  $V(I)$ , the proof of [BS19, Corollary 8.11] actually shows that the diagram

$$\begin{array}{ccc} S'_{\mathrm{perfd}} & \longrightarrow & \mathrm{R}\Gamma(X'_{\mathrm{arc}}, \mathcal{O}_{\mathrm{arc}}) \\ \downarrow & & \downarrow \\ (S'/I)_{\mathrm{perfd}} & \longrightarrow & \mathrm{R}\Gamma((X' \times_{\mathrm{Spec} S'} \mathrm{Spec} S'/I)_{\mathrm{arc}}, \mathcal{O}_{\mathrm{arc}}) \end{array}$$

is a pullback. The two lower modules are over  $S'/I$ , thus  $r$ -almost zero, so the upper arrow is an  $r$ -almost isomorphism. Now it remains to prove that  $R' \rightarrow S'_{\mathrm{perfd}}$  is  $p$ -completely  $r$ -almost split, but this follows from [BS19, Theorem 10.9].  $\square$

#### REFERENCES

- [BS19] Bhargav Bhatt and Peter Scholze. *Prisms and Prismatic Cohomology*. 2019. URL: <https://arxiv.org/abs/1905.08229>.