

Subdivided Claws and the Clique-Stable Set Separation Property

Maria Chudnovsky* and Paul Seymour†

Abstract Let \mathcal{C} be a class of graphs closed under taking induced subgraphs. We say that \mathcal{C} has the *clique-stable set separation property* if there exists $c \in \mathbb{N}$ such that for every graph $G \in \mathcal{C}$ there is a collection \mathcal{P} of partitions (X, Y) of the vertex set of G with $|\mathcal{P}| \leq |V(G)|^c$ and with the following property: if K is a clique of G , and S is a stable set of G , and $K \cap S = \emptyset$, then there is $(X, Y) \in \mathcal{P}$ with $K \subseteq X$ and $S \subseteq Y$. In 1991 M. Yannakakis conjectured that the class of all graphs has the clique-stable set separation property, but this conjecture was disproved by M. Göös in 2014. Therefore it is now of interest to understand for which classes of graphs such a constant c exists. In this paper we define two infinite families \mathcal{S}, \mathcal{K} of graphs and show that for every $S \in \mathcal{S}$ and $K \in \mathcal{K}$, the class of graphs with no induced subgraph isomorphic to S or K has the clique-stable set separation property.

1 Introduction

All graphs in this paper are finite and simple. Let G be a graph. A *clique* in G is a set of pairwise adjacent vertices, and a *stable set* is a set of pairwise non-adjacent vertices. Let \mathcal{C} be a class of graphs closed under taking induced subgraphs. We say that \mathcal{C} has the *clique-stable set separation property* if there exists $c \in \mathbb{N}$ such that for every graph $G \in \mathcal{C}$ there is a collection \mathcal{P} of partitions

Maria Chudnovsky
Princeton University, Princeton, NJ 08544, e-mail: mchudnov@math.princeton.edu

Paul Seymour
Princeton University, Princeton, NJ 08544, e-mail: pds@math.princeton.edu

* This material is based upon work supported in part by the U. S. Army Research Office under grant number W911NF-16-1-0404, and by NSF grant DMS-1763817.

† Partially supported by NSF grant DMS-1800053 and AFOSR grant A9550-19-1-0187.

(X, Y) of the vertex set of G with $|\mathcal{P}| \leq |V(G)|^c$ and with the following property: if K is a clique of G , and S is a stable set of G , and $K \cap S = \emptyset$, then there is $(X, Y) \in \mathcal{P}$ with $K \subseteq X$ and $S \subseteq Y$. This property plays an important role in a large variety of fields: communication complexity, combinatorial optimization, constraint satisfaction and others (for a comprehensive survey of these connections see [3]).

In 1991 Mihalis Yannakakis conjectured that the class of all graphs has the clique-stable set separation property [5], but this conjecture was disproved by Mika Göös in 2014 [2]. Therefore it is now of interest to understand for which classes of graphs such a constant c exists; our main result falls into that category.

Let G be a graph and let X, Y be disjoint subsets of $V(G)$. We denote by $G[X]$ the subgraph of G induced by X , by $N(X)$ the set of all vertices of $V(G) \setminus X$ with a neighbor in X , and by $N[X]$ the set $N(X) \cup X$. We say that X is *complete* to Y if every vertex of X is adjacent to every vertex of Y , and that X is *anticomplete* to Y if every vertex of X is non-adjacent to every vertex of Y . We say that X and Y are *matched* if every vertex of X has exactly one neighbor in Y , and every vertex of Y has exactly one neighbor in X (and therefore $|X| = |Y|$). For a graph H , we say that G is *H -free* if no induced subgraph of G is isomorphic to H .

Next we define two types of graphs. Let $p, q \in \mathbb{N}$. We define the graph $F_S^{p,q}$ as follows:

- $V(F_S^{p,q}) = K \cup S_1 \cup S_2 \cup S_3$ where K is a clique, S_1, S_2, S_3 are stable sets, and the sets K, S_1, S_2, S_3 are pairwise disjoint;
- $|K| = |S_1| = p$, and K and S_1 are matched;
- $|S_2| = |S_3| = q$, and S_2 and S_3 are matched;
- K is complete to S_2 ;
- there are no other edges in $F_S^{p,q}$.

The graph $F_K^{p,q}$ is obtained from $F_S^{p,q}$ by making all pairs of vertices of S_3 adjacent.




Fig. 1 The graphs $F_S^{3,3}$ and $F_K^{3,3}$

Let $\mathcal{F}^{p,q}$ be the class of all graphs that are both $F_S^{p,q}$ -free and $F_K^{p,q}$ -free. We can now state our main result:

Theorem 1. *For all $p, q > 0$ the class $\mathcal{F}^{p,q}$ has the clique-stable set separation property.*

Since the clique-stable set separation property is preserved under taking complements, we immediately deduce:

Theorem 2. *For all $p, q > 0$ the class of graphs whose complements are in $\mathcal{F}^{p,q}$ has the clique-stable set separation property.*

2 The Proof

In this section we prove 1. The idea of the proof comes from [1]. Let $G \in \mathcal{F}^{p,q}$. Define \mathcal{P}_1 to be the set of all partitions $(N[X], V(G) \setminus N[X])$ and $(N(X), V(G) \setminus N(X))$ where X is a subset of $V(G)$ with $|X| < p$. Clearly $|\mathcal{P}_1| \leq 2|V(G)|^p$.

Write $R = R(q, q)$ to mean the smallest positive integer R such that every 2-coloring of the edges of the complete graph on R vertices contains a monochromatic complete graph on q vertices. Ramsey's Theorem [4] implies:

Theorem 3. $R(q, q) \leq 2^{2q}$.

For $a, b \in \mathbb{N}$ let the graph $F_{a,b}$ be defined as follows:

- $V(F_{a,b}) = K_1 \cup S_1 \cup S_2 \cup W$ where K_1 is a clique, S_1, S_2 are stable sets, and the sets K_1, S_1, S_2, W are pairwise disjoint;
- $|K_1| = |S_1| = a$, and K_1 and S_1 are matched;
- $|S_2| = |W| = b$, and S_2 and W are matched;
- K_1 is complete to S_2 ;
- there is no restriction on the adjacency of pairs of vertices of W ;
- there are no other edges in $F_{a,b}$.

From the definition of R we immediately deduce:

Theorem 4. G is $F_{p,R}$ -free.

For every triple $X = (K_1, S_1, S_2)$ of pairwise disjoint non-empty subsets of $V(G)$ such that $|K_1| = |S_1| = p$ and $|S_2| < R$ we define the partition P_X of $V(G)$ as follows. Let Z be the set of all vertices of G that are anticomplete to $K_1 \cup S_1$. Let A_X be the set of all vertices v of G such that

- either $v \in K_1$, or v is complete to K_1 , and
- either v has a neighbor in S_1 , or v has a neighbor in $Z \setminus N(S_2)$.

Note that, since S_1 is a stable set and Z is anticomplete to S_1 , A_X is disjoint from $S_1 \cup Z$. Define $P_X = (A_X, V(G) \setminus A_X)$, and let \mathcal{P}_2 be the set of all such partitions P_X . Since $|K_1 \cup S_1 \cup S_2| \leq 2p + R - 1$, and since by 3 $R \leq 2^{2q}$, we deduce that $|\mathcal{P}_2| < |V(G)|^{2p+2^{2q}}$.

In order to complete the proof of 1 we will prove the following:

Theorem 5. *For every clique K and stable set S of G such that $K \cap S = \emptyset$, there exists $(X, Y) \in \mathcal{P}_1 \cup \mathcal{P}_2$ with $K \subseteq X$ and $S \subseteq Y$.*

Proof. Let K and S be as in the statement of 5.

- (1) *We may assume that K is a maximal clique of G , and S is a maximal stable set of G .*

Let K' be a maximal clique of G with $K \subseteq K'$, and let S' be a maximal stable set of G with $S \subseteq S'$. If $K' \cap S' = \emptyset$, then the existence of the desired partition for K, S follows from the existence of such a partition for K', S' ; thus we may assume that $K' \cap S' \neq \emptyset$. Since K' is a clique and S' is a stable set, it follows that $|K' \cap S'| = 1$, say $K' \cap S' = \{v\}$. But now the partitions $(N[\{v\}], V(G) \setminus N[\{v\}])$ and $(N(\{v\}), V(G) \setminus N(\{v\}))$ are both in \mathcal{P}_1 , and at least one of them has the desired property. This proves (1).

In view of (1) from now on we assume that K is a maximal clique of G , and S is a maximal stable set of G . Consequently every vertex of K has a neighbor in S . Let $S'_1 \subseteq S$ be a minimal subset of S such that every vertex of K has a neighbor in S'_1 . It follows from the minimality of S'_1 that there is a subset K'_1 of K such that S'_1 and K'_1 are matched. If $|S'_1| < p$, then the partition $(N(S'_1), V(G) \setminus N(S'_1)) \in \mathcal{P}_1$ has the desired property, so we may assume that $|S'_1| \geq p$.

Let S_1 be a subset of S'_1 with $|S_1| = p$, and let $K_1 = N(S_1) \cap K'_1$. Then S_1 and K_1 are matched, and so $|K_1| = p$. Let Z be the set of vertices of G that are anticomplete to $S_1 \cup K_1$. Then $S'_1 \setminus S_1 \subseteq Z \cap S$, and in particular every vertex of K has a neighbor either in S_1 or in $Z \cap S$. Let S' be the subset of vertices of $S \setminus S_1$ that are complete to K_1 . Note that $S' \cap Z = \emptyset$. Let S_2 be a minimal subset of S' such that $N(S_2) \cap Z = N(S') \cap Z$. It follows from the minimality of S_2 that there is a subset $W \subseteq Z \cap N(S')$ such that W and S_2 are matched. Observe that $G[K_1 \cup S_1 \cup S_2 \cup W]$ is isomorphic to $F_{p, |S_2|}$ (with K_1, S_1, S_2, W as in the definition of $F_{a,b}$). It follows from 4 that $|S_2| < R$.

Let $X = (K_1, S_1, S_2)$. We claim that the partition $P_X \in \mathcal{P}_2$ has the desired property for the pair K, S . Recall that $P_X = (A_X, V(G) \setminus A_X)$, where A_X is the set of all vertices v of G such that

- either $v \in K_1$, or v is complete to K_1 , and
- either v has a neighbor in S_1 , or v has a neighbor in $Z \setminus N(S_2)$.

We need to show that $K \subseteq A_X$, and $S \cap A_X = \emptyset$.

(2) $K \subseteq A_X$.

Let $k \in K$. Clearly either $k \in K_1$ or k is complete to K_1 . Moreover, k has a neighbor in S'_1 , and $S'_1 \subseteq S_1 \cup (Z \cap S)$. Since S is a stable set, it follows that $Z \cap S \subseteq Z \setminus N(S_2)$, and thus k has a neighbor either in S_1 , or in $Z \setminus N(S_2)$. This proves (2).

(3) $S \cap A_X = \emptyset$.

Suppose that $s \in S \cap A_X$. Then $s \notin K_1$; therefore s is complete to K_1 , and so $s \in S'$. Since S is a stable set, it follows that s is anticomplete to S_1 , and therefore s has a neighbor in $Z \setminus N(S_2)$. But $N(S') \cap Z = N(S_2) \cap Z$, a contradiction. This proves (3).

Now 5 follows from (2) and (3). \square

This completes the proof of 1.

Acknowledgment

This work was done during the Structural Graph Theory Downunder Matrix Program, that was held in 2019 at the Creswick Campus of the University of Melbourne. The authors express their gratitude to the Matrix Center for the funding it provided and the use of its facilities, and to the organizers of the program for the invitation.

References

1. T. Abrishami, M. Chudnovsky, M. Pilipczuk, P. Rzazewski and P. Seymour: Induced subgraphs of bounded tree-width and the container method, *in preparation*.
2. M. Göös: Lower bounds for clique vs. independent Set. In: Proc. 56th Foundations of Computer Science (FOCS), 2015: 1066–1077.
3. A. Lagoutte: Interactions entre les cliques et les stables dans un graphe. PhD thesis, ENS de Lyon, 2015.
4. F.P. Ramsey: On a problem of formal logic. Proc. London Math. Soc. **30**, 264–286 (1930).
5. M. Yannakakis: Expressing combinatorial optimization problems by linear programs. J. Comput. Syst. Sci. **43**, 441–466 (1991).