

INDUCED SUBGRAPHS AND TREE DECOMPOSITIONS XVI. COMPLETE BIPARTITE INDUCED MINORS

MARIA CHUDNOVSKY^{†*}, SEPEHR HAJEBI[§], AND SOPHIE SPIRKL^{§||}

ABSTRACT. We prove that for every graph G with a sufficiently large complete bipartite induced minor, either G has an induced minor isomorphic to a large wall, or G contains a large *constellation*; that is, a complete bipartite induced minor model such that on one side of the bipartition, each branch set is a singleton, and on the other side, each branch set induces a path.

We further refine this theorem by characterizing the unavoidable induced subgraphs of large constellations as two types of highly structured constellations. These results will be key ingredients in several forthcoming papers of this series.

1. INTRODUCTION

The set of all positive integers is denoted by \mathbb{N} . For each $k \in \mathbb{N}$, we write \mathbb{N}_k for the set of all positive integers not greater than k . Graphs in this paper have finite vertex sets, no loops and no parallel edges. Given a graph $G = (V(G), E(G))$ and $X \subseteq V(G)$, we use X to denote both the set X and the subgraph $G[X]$ of G induced by X . For a set \mathcal{X} of subsets of $V(G)$, we write $V(\mathcal{X}) = \bigcup_{X \in \mathcal{X}} X$. We say that G is H -free, for another graph H , if G has no induced subgraph isomorphic to H .

This series of papers studies the interplay between induced subgraphs and *treewidth* (where the treewidth of a graph G is denoted by $\text{tw}(G)$; see [10] for a definition).

Specifically, one of the main goals is to answer the following question:

Question 1.1. *What are the unavoidable induced subgraphs of graphs with large treewidth?*

The answer is known if we replace “induced subgraphs” by “subgraphs” or “minors” as shown by Robertson and Seymour [19] (where $W_{r \times r}$ denotes the r -by- r hexagonal grid, also known as the r -by- r wall; see Figure 1):

Date: 15th November, 2024.

[†] Princeton University, Princeton, NJ, USA.

[§] Department of Combinatorics and Optimization, University of Waterloo, Waterloo, Ontario, Canada.

* Supported by NSF-EPSC Grant DMS-2120644, AFOSR grant FA9550-22-1-0083 and NSF Grant DMS-2348219.

|| We acknowledge the support of the Natural Sciences and Engineering Research Council of Canada (NSERC), [funding reference number RGPIN-2020-03912]. Cette recherche a été financée par le Conseil de recherches en sciences naturelles et en génie du Canada (CRSNG), [numéro de référence RGPIN-2020-03912]. This project was funded in part by the Government of Ontario. This research was conducted while Spirkl was an Alfred P. Sloan Fellow.




FIGURE 1. The basic obstructions of treewidth 4: $K_{4,4}$ (top left), K_5 (bottom left), a subdivision of the 4×4 wall (middle) and the line graph of a subdivision of the 4×4 wall (right).

Theorem 1.2 (Robertson and Seymour [19]). *For every integer $r \in \mathbb{N}$, there is a constant $f_{1.2} = f_{1.2}(r) \in \mathbb{N}$ such that every graph G with $\text{tw}(G) \geq f_{1.2}$ has a subgraph isomorphic to a subdivision of $W_{r \times r}$.*

The answer to Question 1.1 is necessarily more complicated, as it needs to include all *basic obstructions*: complete graphs, complete bipartite graphs, subdivided walls, and the line graphs of subdivided walls (see Figure 1). Moreover, basic obstructions do not form a complete answer to Question 1.1, as shown by several examples of “non-basic” obstructions such as the Pohoata-Davies graphs [9, 16], “occultations” [2, 6], and “layered wheels” [21].

It appears that the key to answering Question 1.1 is to understand the unavoidable induced subgraphs of graphs with large dense minors. In all non-basic obstructions, large treewidth is caused by the presence of a large complete minor (and that is for a good reason: it has been shown several times [1, 20, 22] that every graph with large enough treewidth and no induced subgraph isomorphic to a basic obstruction of large treewidth has a large complete minor). In fact, the first two non-basic obstructions mentioned above, the Pohoata-Davies graphs and the occultations (see Figure 2), have large complete bipartite induced minors (while the third, layered wheels, do not [8]. This also follows directly from our main result, Theorem 2.3.)

It is therefore natural to ask for a characterization of the unavoidable induced subgraphs of graphs with large complete bipartite induced minors (at least in the absence of basic obstructions of large treewidth). Our main result is exactly this characterization. As it turns out, the Pohoata-Davies graphs and the occultations are essentially the only (non-basic) outcomes.

Let us make all this precise, beginning with a few definitions. Let G be a graph. We say that $X, Y \subseteq V(G)$ are *anticomplete in G* if $X \cap Y = \emptyset$ and there is no edge in G with an end in X and an end in Y (if $X = \{x\}$ is a singleton, then we also say x is *anticomplete to Y in G*). Given a graph H , we say G has an *induced minor isomorphic to H* if there are pairwise disjoint connected induced subgraphs ($G_v : v \in V(H)$) of G , which we call the *branch sets*, such that for all distinct $u, v \in V(G)$, we have $uv \in E(H)$ if and only if $G_u, G_v \subseteq V(G)$ are not anticomplete in G . Both the Pohoata-Davies graphs and the occultations contain




FIGURE 2. A Pohoata-Davies graph (left) and an occultation (right), both of treewidth 4.




FIGURE 3. A $(3, 3)$ -constellation.

complete bipartite induced minors of a special type: On one side of the bipartition, each branch set is a singleton, and on the other side, each branch set induces a path. We call such graphs *constellations*, and analyzing the structure of constellations was one of the main tools in many of our earlier papers [2, 3, 4, 5, 14]. Formally, a *constellation* is a graph \mathfrak{c} in which there is a stable set $S_{\mathfrak{c}}$ such that every component of $\mathfrak{c} \setminus S_{\mathfrak{c}}$ is a path, and each vertex $x \in S_{\mathfrak{c}}$ has at least one neighbor in each component of $\mathfrak{c} \setminus S_{\mathfrak{c}}$. We denote by $\mathcal{L}_{\mathfrak{c}}$ the set of all components $\mathfrak{c} \setminus S_{\mathfrak{c}}$ (each of which is a path), and denote the constellation \mathfrak{c} by the pair $(S_{\mathfrak{c}}, \mathcal{L}_{\mathfrak{c}})$. For $l, s \in \mathbb{N}$, by an (s, l) -constellation we mean a constellation \mathfrak{c} with $|S_{\mathfrak{c}}| = s$ and $|\mathcal{L}_{\mathfrak{c}}| = l$ (see Figure 3). Given a graph G , by an (s, l) -constellation in G we mean an induced subgraph of G which is an (s, l) -constellation.

Note that an (s, l) -constellation has treewidth at least $\min\{s, l\}$, because it has an induced minor isomorphic to the complete bipartite graph $K_{s,l}$ (obtained by contracting all edges of the paths in $\mathcal{L}_{\mathfrak{c}}$). Our first result shows that, in the absence of basic obstructions, the converse is also true:

Theorem 1.3. *For all $l, r, s \in \mathbb{N}$, there is a constant $f_{1.3} = f_{1.3}(l, r, s) \in \mathbb{N}$ such that if G is a graph with an induced minor isomorphic to $K_{f_{1.3}, f_{1.3}}$, then one of the following holds.*

- (a) *There is an induced minor of G isomorphic to $W_{r \times r}$.*
- (b) *There is an (s, l) -constellation in G .*

It remains to describe the unavoidable induced subgraphs of large constellations, which we will do in our second result. The exact statement will be given as Theorem 2.1, which roughly says the following: for all $l, r, s \in \mathbb{N}$, given a sufficiently large constellation \mathbf{c} , either a basic obstruction of treewidth r is present in \mathbf{c} as an induced subgraph, or there is an (s, l) -constellation \mathbf{b} in \mathbf{c} along with a linear order x_1, \dots, x_s of the vertices in $S_{\mathbf{b}}$ such that one of the following holds:

- For all $1 \leq i < j \leq s$ and every path P from x_i to x_j with interior in $V(\mathcal{L})$, every vertex x_k with $j < k \leq s$ has a neighbor in P ; or
- There is a constant $c = c(r)$ (actually, $c(r) = 2r^2$ works) such that for all $1 \leq i < k \leq s$ and every path P from x_i to x_k with interior in $V(\mathcal{L})$, we have

$$|\{j \in \{i+1, \dots, k-1\} : N(x_j) \cap V(P) = \emptyset\}| < c.$$

In particular, the first of these is a slight generalization of occultations. The second is a relatively substantial generalization of the Pohoata-Davies graphs, but we will show in Section 3 that this outcome cannot be replaced with (straightforward variants of) the Pohoata-Davies graphs. We will also prove in the final section that both outcomes are necessary.

2. UNAVOIDABLE CONSTELLATIONS AND THE MAIN RESULT

Here we give the exact statement of our second result, Theorem 2.1. Then we state our main result, Theorem 2.3, as a direct combination of Theorems 1.3 and 2.1.

We need several definitions. Let P be a graph which is a path. Then we write $P = p_1 \cdots p_k$ to mean $V(P) = \{p_1, \dots, p_k\}$ for $k \in \mathbb{N}$, and $E(P) = \{p_i p_{i+1} : i \in \mathbb{N}_{k-1}\}$. We call p_1, p_k the *ends* of P , and we call $P \setminus \{p_1, p_k\}$ the interior of P , denoted P^* . Given a graph G , by a *path in G* we mean an induced subgraph of G that is a path.

Given a graph G , a *subdivision of G* is a graph obtained from G by replacing each edge $e = uv$ of G by a path P_e of length at least 1 from u to v such that the interiors of the paths are pairwise disjoint and anticomplete. For $d \in \mathbb{N}$, this is a *d -subdivision* (*$(\leq d)$ -subdivision*, *$(\geq d)$ -subdivision*) if each path P_e has length exactly $d+1$ (at most $d+1$, at least $d+1$) for all $e \in E(G)$. A subdivision is *proper* if it is a (≥ 1) -subdivision. For a set X , a linear order \preceq on X , and $x, y \in X$, we write $x \prec y$ to mean $x \preceq y$ and x and y are distinct. For an element $x \in X$ and a subset $Y \subseteq X$, we write $x \prec Y$ to mean $x \prec y$ for every $y \in Y$. Similarly, we write $Y \prec x$ to mean $y \prec x$ for every $y \in Y$.

Let $\mathbf{c} = (S_{\mathbf{c}}, \mathcal{L}_{\mathbf{c}})$ be a constellation. By a \mathbf{c} -*route* we mean a path R in \mathbf{c} with ends in $S_{\mathbf{c}}$ and with $R^* \subseteq V(\mathcal{L}_{\mathbf{c}})$, or equivalently, with $R^* \subseteq V(L)$ for some $L \in \mathcal{L}_{\mathbf{c}}$. For $d \in \mathbb{N}$, we say \mathbf{c} is *d -ample* if there is no \mathbf{c} -route of length at most $d+1$. We also say \mathbf{c} is *ample* if \mathbf{c} is 1-ample (that is, no two vertices in $S_{\mathbf{c}}$ have a common neighbor in $V(\mathcal{L}_{\mathbf{c}})$; see Figure 4).




FIGURE 4. A \mathfrak{c} -route of length four in the $(3,3)$ -constellation \mathfrak{c} from Figure 3, which is 2-ample.




FIGURE 5. A $(4,1)$ -constellation which is interrupted with $x_1 \preccurlyeq x_2 \preccurlyeq x_3 \preccurlyeq x_4$.

We say a constellation \mathfrak{c} is *interrupted* if there is a linear order \preccurlyeq on $S_{\mathfrak{c}}$ such that for all $x, y, z \in S_{\mathfrak{c}}$ with $x \prec y \prec z$ and every \mathfrak{c} -route R from x to y , the vertex z has a neighbor in R (see Figure 5). We remark that occultations as in [2, 6] are a special case of this (compare Figure 2 (right) and Figure 5).

For $q \in \mathbb{N}$, we say a constellation \mathfrak{c} is q -zigzagged if there is a linear order \preccurlyeq on $S_{\mathfrak{c}}$ such that for all $x, y \in S_{\mathfrak{c}}$ with $x \prec y$ and every \mathfrak{c} -route R from x to y , there are fewer than q vertices $z \in S_{\mathfrak{c}}$ where $x \prec z \prec y$ and z has no neighbor in R (see Figure 6).

We also need a canonical “containment” relation on constellations, which we define next. For constellations \mathfrak{b} and \mathfrak{c} , we say \mathfrak{b} *sits in* \mathfrak{c} if

- $S_{\mathfrak{b}} \subseteq S_{\mathfrak{c}}$;
- for every $L \in \mathcal{L}_{\mathfrak{b}}$, there exists $M \in \mathcal{L}_{\mathfrak{c}}$ such that $L \subseteq M$; and
- for every $M \in \mathcal{L}_{\mathfrak{c}}$, there is at most one path $L \in \mathcal{L}_{\mathfrak{b}}$ such that $L \subseteq M$.

See Figure 7. It follows that for constellations \mathfrak{b} and \mathfrak{c} where \mathfrak{b} sits in \mathfrak{c} , the useful properties of \mathfrak{c} are also inherited by \mathfrak{b} . In particular,




FIGURE 6. Left: A $(4, 1)$ -constellation which is 1-zigzagged with $x_1 \preceq x_2 \preceq x_3 \preceq x_4$. Right: A $(6, 1)$ -constellation which is 1-zigzagged with $x_1 \preceq x_2 \preceq x_3 \preceq x_4 \preceq x_5 \preceq x_6$.




FIGURE 7. A $(2, 2)$ -constellation sitting in the $(3, 3)$ -constellation from Figure 3.

- \mathfrak{b} is an induced subgraph of \mathfrak{c} ;
- if \mathfrak{c} is d -ample for some $d \in \mathbb{N}$, then so is \mathfrak{b} ;
- if \mathfrak{c} is interrupted, then so is \mathfrak{b} ; and
- if \mathfrak{c} is q -zigzagged for some $q \in \mathbb{N}$, then so is \mathfrak{b} .

We are now ready to state our second result:

Theorem 2.1. *For all $d, l, l', r, s, s', t \in \mathbb{N}$, there are constants $f_{2.1} = f_{2.1}(d, l, l', r, s, s', t) \in \mathbb{N}$ and $g_{2.1} = g_{2.1}(d, l, l', r, s, s', t) \in \mathbb{N}$ such that for every $(f_{2.1}, g_{2.1})$ -constellation \mathfrak{c} , one of the following holds.*

- There is an induced subgraph of \mathfrak{c} isomorphic to either $K_{r,r}$ or a proper subdivision of K_{2t+2} .*
- There is a d -ample interrupted (s, l) -constellation which sits in \mathfrak{c} .*
- There is a d -ample $2t$ -zigzagged (s', l') -constellation which sits in \mathfrak{c} .*

We also need the next result:

Lemma 2.2 (Aboulker, Adler, Kim, Sintuari, Trotignon; see Lemma 3.6 in [1]). *For every $r \in \mathbb{N}$, there is a constant $f_{2.2} = f_{2.2}(r)$ such that if G is a graph with an induced minor isomorphic to a subdivision of $W_{f_{2.2} \times f_{2.2}}$, then G has an induced subgraph isomorphic to either a subdivision of $W_{r \times r}$ or the line graph of a subdivision of $W_{r \times r}$.*

We can now state the main result of this paper, which follows directly from combining Theorems 1.3 and 2.1 with Lemma 2.2.

Theorem 2.3. *For all $d, l, l', r, s, s' \in \mathbb{N}$, there are constants $f_{2.3} = f_{2.3}(d, l, l', r, s, s') \in \mathbb{N}$ and $g_{2.3} = g_{2.3}(d, l, l', r, s, s') \in \mathbb{N}$ with the following property. Let G be a graph which has an induced minor isomorphic to $K_{f_{2.3}, g_{2.3}}$. Then one of the following holds.*

- (a) *There is an induced subgraph of G isomorphic to either $K_{r,r}$, a subdivision of $W_{r \times r}$, or the line graph of a subdivision of $W_{r \times r}$.*
- (b) *There is a d -ample and interrupted (s, l) -constellation G .*
- (c) *There is a d -ample and $2r^2$ -zigzagged (s', l') -constellation in G .*

It remains to prove Theorems 1.3 and 2.1, which we will do in Sections 7 and 8.

This paper is organized as follows. In Section 3, we will give a construction showing that the third outcome of Theorem 2.3 cannot be strengthened to yield the Pohoata-Davies graphs. Then, we proceed to the proof of Theorem 2.1: In Section 4, we show that we can restrict our attention to d -ample constellations. In Section 5, we obtain a precursor to outcomes (b) and (c) of Theorem 2.1 in which a “mixture” of the two cases holds. Section 6 recovers these two distinct outcomes, and Section 7 finishes the proof of Theorem 2.1. In Section 8, we prove Theorem 1.3, hence completing the proof of Theorem 2.3. Finally, in Section 9, we will prove that interrupted constellations and zigzagged constellations are both unavoidable as outcomes of Theorems 2.1 and 2.3.

3. THE ZIGZAG GRAPH

Before we continue to our main proofs, let us expand on Theorem 2.1(c). One might hope that this outcome could be refined further to coincide with the Pohoata-Davies graphs [9, 16] or more precisely, with “arrays,” which are slight generalizations of Pohoata-Davies graphs [4]. Let \mathfrak{c} be a constellation. We say that \mathfrak{c} is *aligned* if there is a linear order \preceq on $S_{\mathfrak{c}}$ such that every path L in $\mathcal{L}_{\mathfrak{c}}$ traverses neighbors of $S_{\mathfrak{c}}$ in order, that is, there is a labelling v_1, \dots, v_t of the vertices of L such that for all $i, j \in \mathbb{N}_t$:

- (1) $v_i v_j \in E(L)$ if and only if $|i - j| = 1$; and
- (2) if $s, s' \in S_{\mathfrak{c}}$ with $s \prec s'$ such that $v_i \in N(s)$ and $v_j \in N(s')$, then $i < j$.

Arrays are aligned constellations; Pohoata-Davies graphs have some further restrictions.

Writing x_1, \dots, x_s for the vertices of $S_{\mathfrak{c}}$, each path L of $\mathcal{L}_{\mathfrak{c}}$ gives rise to a sequence A_L (unique up to reversal) obtained by recording, as we traverse L , the indices of vertices in $S_{\mathfrak{c}}$ whose neighbors we encounter. For example, in Figure 6, this sequence would read 1, 1, 1, 2, 2, 3, 4, 4 for the graph of the left, and begin with 1, 2, 3, 2, 3, 4 for the graph on the right. A constellation is aligned (with the linear order $x_1 \preceq \dots \preceq x_s$), then, if for every $L \in \mathcal{L}_{\mathfrak{c}}$, we have $A_L = 1, 2, \dots, s$ after omitting consecutive occurrences of the same number (and up to reversal).

Let us say that a sequence $A = a_1, \dots, a_t$ of integers is *smooth* if consecutive entries differ by at most 1. Thus, a constellation \mathbf{c} is 1-zigzagged if and only if there is a linear order on \mathbf{c} such that A_L is smooth for every $L \in \mathcal{L}_{\mathbf{c}}$.

One might hope that if \mathbf{c} is a sufficiently large 1-zigzagged constellation, then there is a large aligned constellation which sits in \mathbf{c} . Translating this to sequences, we would hope that for every smooth sequence $A = a_1, \dots, a_t$ with $|\{a_1, \dots, a_t\}|$ sufficiently large, there is a large subset $B \subseteq \{a_1, \dots, a_t\}$ as well as $i, j \in \mathbb{N}_t$ such that the sequence $A^{B,i,j}$, obtained from a_i, \dots, a_j by skipping all entries which are not in B and then omitting consecutive entries that are equal, is a sequence containing each element of B exactly once. Let us say that B is an *alignment in A* in this case; we call $|B|$ the *size* of the alignment. We will show that for every n , there is a smooth sequence containing the numbers $1, \dots, n$ with no alignment of size 4. We call this sequence the *zigzag sequence*. The associated constellation is 1-zigzagged, but no aligned constellation with a stable set of size 4 sits in it.

To define the zigzag sequence, we need some further definitions. Given a sequence $A = a_1, \dots, a_t$, we define $A+1$ to be the sequence $a_1+1, a_2+1, \dots, a_t+1$. Moreover, we write A^{-1} for the *reverse of A* , that is, the sequence a_t, a_{t-1}, \dots, a_1 . We define A^* to be the sequence A_1, \dots, A_{t-1} .

We are now ready to define the *zigzag sequence* Z_n for $n \in \mathbb{N}_{\geq 2}$. We let $Z_2 = 1, 2$ and $Z_3 = 1, 2, 3$. For $n \geq 4$, we define to be the sequence $Z_n = Z_{n-1}^*, ((Z_{n-2} + 1)^{-1})^*, (Z_{n-1}) + 1$. Thus,

$$Z_4 = 1, 2, 3, 2, 3, 4$$

and

$$Z_5 = 1, 2, 3, 2, 3, 4, 3, 2, 3, 4, 3, 4, 5.$$

From the definition, it follows that Z_n is a smooth sequence containing the numbers $1, \dots, n$. In addition, replacing each entry a_i of Z_n with $n+1-a_i$ yields the sequence $(Z_n)^{-1}$. We write l_n for the length of Z_n .

Lemma 3.1. *Z_n contains no 4-alignment.*

Proof. Suppose for a contradiction that $\{p, q, r, s\}$ is an alignment in the sequence $A = z_i, \dots, z_j$ for some $i < j$, where $Z_n = z_1, \dots, z_{l_n}$. We choose this counterexample with n minimum. We may further choose this sequence with $j-i$ minimum, and so we may assume that $z_i = p$ and $z_j = s$ and $z_k \notin \{p, s\}$ for $k \in \{i+1, \dots, j-1\}$. By symmetry, we may assume that $p < s$. It follows that $z_{i+1}, \dots, z_{j-1} \in \{p+1, \dots, s-1\}$.

We claim that $p = 1$. Suppose not, so $p \geq 2$. It follows that all entries of A except possibly z_i are strictly larger than 2, and so A is contained in either $(Z_{n-1}) + 1$ or $Z_{n-1}^*, ((Z_{n-2} + 1)^{-1})^*, 2 = Z_{n-1}^*, ((Z_{n-2} + 1)^{-1})$. The former contradicts the minimality of n , and so the latter case holds. In particular, A does not contain n , and so $s \leq n-1$. Since the first entry of $((Z_{n-2} + 1)^{-1})$ is $n-1$, it follows that A is contained in one of $Z_{n-1}^*, n-1 = Z_{n-1}$ or $((Z_{n-2} + 1)^{-1})$; however, both contradict the minimality of n . Thus, $p = 1$, and by symmetry, $s = n$. It follows that $i = 1$ and $j = l_n$, and so $A = Z_n$.

By symmetry, we may assume that $q < r$. Then,

$$A = Z_n = Z_{n-1}^*, ((Z_{n-2} + 1)^{-1})^*, (Z_{n-1}) + 1$$

and Z_{n-1}^* contains q (and possibly r), $((Z_{n-2} + 1)^{-1})^*$ contains r (and possibly q), and $(Z_{n-1}) + 1$ contains q (and possibly r); but this contradicts the assumption that $\{p, q, r, s\}$ is an alignment in A , hence completing the proof. \blacksquare

The *zigzag graph* GZ_n is defined as follows. Let $Z_n = z_1, \dots, z_{l_n}$. We let $V(GZ_n) = \{p_1, \dots, p_{l_n}, x_1, \dots, x_n\}$ and

$$E(GZ_n) = \{p_i p_{i+1} : i \in \{1, \dots, l_n - 1\}\} \cup \{x_i p_j : i \in \{1, \dots, n\}, j \in \{1, \dots, l_n\}, z_j = i\}.$$

This is an $(n, 1)$ -constellation which is 1-zigzagged (Figure 6 (right) shows GZ_6); by making l copies of the path p_1, \dots, p_{l_n} we obtain an (n, l) -constellation GZ_n^l . Using Lemma 3.1, one can show that GZ_n^l contains no large array; we omit the proof. It follows that it is not possible to simplify Theorem 2.1(c) to the case of arrays. We leave open the question of whether “ $2t$ -zigzagged” can be improved to “1-zigzagged.”

4. AMPLIFICATION

The first step in our proof of Theorem 2.1 is to pass to a sufficiently ample constellation; doing so makes it easier to find routes which are pairwise anticomplete. This will be done using the following lemma:

Lemma 4.1. *For all $d, l, r, r', s \in \mathbb{N}$, there are constants $f_{4.1} = f_{4.1}(d, l, r, r', s) \in \mathbb{N}$ and $g_{4.1} = g_{4.1}(d, l, r, r', s) \in \mathbb{N}$ such that for every $(f_{4.1}, g_{4.1})$ -constellation \mathfrak{c} , one of the following holds.*

- (a) *There is an induced subgraph of \mathfrak{c} which is isomorphic to either $K_{r,r}$ or a proper subdivision of $K_{r'}$.*
- (b) *There is a d -ample (s, l) -constellation \mathfrak{d} with $S_{\mathfrak{d}} \subseteq S_{\mathfrak{c}}$ and $\mathcal{L}_{\mathfrak{d}} \subseteq \mathcal{L}_{\mathfrak{c}}$.*

We will deduce Lemma 4.1 from a straightforward application of the next two results:

Lemma 4.2 (Alecú, Chudnosvky, Hajebi and Spirkl; see Lemma 6.2 in [2]). *For all $l, m, s \in \mathbb{N}$, there is a constant $f_{4.2} = f_{4.2}(l, m, s) \in \mathbb{N}$ such that for every $(f_{4.2}, l + m^2 - 1)$ -constellation \mathfrak{c} and every $d \in \mathbb{N}$, one of the following holds.*

- (a) *There is a subgraph of \mathfrak{c} which is isomorphic to a $(\leq d)$ -subdivision of K_m .*
- (b) *There is a d -ample (s, l) -constellation \mathfrak{d} with $S_{\mathfrak{d}} \subseteq S_{\mathfrak{c}}$ and $\mathcal{L}_{\mathfrak{d}} \subseteq \mathcal{L}_{\mathfrak{c}}$.*

Lemma 4.3 (Hajebi [15]; see also Dvořák [12]). *For all $d, r, r' \in \mathbb{N}$, there is a constant $f_{4.3} = f_{4.3}(d, r, r') \in \mathbb{N}$ with the following property. Let G be a K_r -free and $K_{r,r}$ -free graph which has a subgraph isomorphic to a $(< d)$ -subdivision of $K_{f_{4.3}}$. Then G has an induced subgraph isomorphic to a proper $(< d)$ -subdivision of $K_{r'}$.*

Proof of Lemma 4.1. Let $m = f_{4.3}(d, \max\{r, 4\}, r')$, let

$$f_{4.1} = f_{4.1}(d, l, r, r', s) = f_{4.2}(l, m, s)$$

and let

$$g_{4.1} = g_{4.1}(d, l, r, r', s) = l + m^2 - 1.$$

Let \mathfrak{c} be a $(f_{4.1}, g_{4.1})$ -constellation. Apply Lemma 4.2 to \mathfrak{c} . Since 4.1(b) and 4.2(b) are identical, we may assume that 4.2(a) holds, that is, \mathfrak{c} has a subgraph isomorphic to a $(\leq d)$ -subdivision of K_m . Now, observe that \mathfrak{c} is K_4 -free. Moreover, if G has a subgraph

isomorphic to $K_{r,r}$, then 4.2(a) holds. So we may assume that \mathfrak{c} is $K_{r,r}$ -free, as well. This, along with the choice of m , allows for an application of Lemma 4.3. We conclude that \mathfrak{c} has an induced subgraph isomorphic to a proper subdivision of $K_{r,r}$, which in turn implies that 4.1(a) holds, as desired. \blacksquare

5. MIXED CONSTELLATIONS

In this section, we prove a preliminary version of Theorem 2.1 in which the interrupted and the zigzagged outcomes are replaced by a “mixture” of the two.

Recall that for $q \in \mathbb{N}$, a constellation \mathfrak{c} is q -zigzagged if there is a linear order \preceq on $S_{\mathfrak{c}}$ such for all $x, y \in S_{\mathfrak{c}}$ with $x \prec y$, every \mathfrak{c} -route R from x to y and every q -subset Q of $S_{\mathfrak{c}}$ with $x \prec Q \prec y$, some vertex in Q has a neighbor in R . In a “mixed” constellation, we relax this condition by allowing Q to also contain a controlled number of vertices in $S_{\mathfrak{c}}$ that are **not** between the ends of R .

More precisely, for $q \in \mathbb{N}$ and $q^-, q^+ \in \mathbb{N} \cup \{0\}$, we say a constellation \mathfrak{c} is (q^-, q, q^+) -mixed if there is a linear order \preceq on $S_{\mathfrak{c}}$ for which the following holds. Let $x, y \in S_{\mathfrak{c}}$ with $x \prec y$, let R be a \mathfrak{c} -route from x to y and let $Q^-, Q, Q^+ \subseteq S_{\mathfrak{c}}$ such that $|Q^-| = q^-$, $|Q| = q$, $|Q^+| = q^+$ and $Q^- \prec x \prec Q \prec y \prec Q^+$. Then some vertex in $Q^- \cup Q \cup Q^+$ has a neighbor in R . In particular, observe that \mathfrak{c} is $(0, q, 0)$ -mixed if and only if \mathfrak{c} is q -zigzagged. Also, for constellations \mathfrak{b} and \mathfrak{c} where \mathfrak{b} sits in \mathfrak{c} , if \mathfrak{c} is (q^-, q, q^+) -mixed for some $q^-, q^+ \in \mathbb{N} \cup \{0\}$ and $q \in \mathbb{N}$, then so is \mathfrak{b} .

We show that:

Lemma 5.1. *For all $d, l, r, s, t \in \mathbb{N}$, there are constants $f_{5.1} = f_{5.1}(d, l, r, s, t) \in \mathbb{N}$ and $g_{5.1} = g_{5.1}(d, l, r, s, t) \in \mathbb{N}$ with the following property. Let \mathfrak{c} be a $(f_{5.1}, g_{5.1})$ -constellation. Then one of the following holds.*

- (a) *There is an induced subgraph of \mathfrak{c} which isomorphic to either $K_{r,r}$ or a proper subdivision of K_{2t+2} .*
- (b) *There is a d -ample and $(2t, 2t, 2t)$ -mixed (s, l) -constellation \mathfrak{b} which sits in \mathfrak{c} .*

The proof uses the following version of Ramsey’s theorem:

Theorem 5.2 (Ramsey [17]). *For all $l, m, n \in \mathbb{N}$, there is a constant $f_{5.2} = f_{5.2}(l, m, n) \in \mathbb{N}$ with the following property. Let U be a set of cardinality at least $f_{5.2}$ and let F be a non-empty set of cardinality at most l . Let $\Phi : \binom{U}{m} \rightarrow F$ be a map. Then there exist $i \in F$ and an n -subset Z of U such that $\Phi(X) = i$ for all $X \in \binom{Z}{m}$.*

The main idea of the proof is the following: If there is a large X subset of $S_{\mathfrak{c}}$ such that for every two vertices in X , there are many \mathfrak{c} -routes between them avoiding neighbors of the remaining vertices in Q , then this is an induced subdivision of $K_{|Q|}$. So we apply Theorem 5.2, asking for each subset X which \mathfrak{c} -routes exist and avoid neighbors of other vertices.

Proof of Lemma 5.1. Let

$$m = f_{5.2} \left(\binom{2t+2}{2}, \binom{(l+1)\binom{2t+2}{2}}{l}, 2t+2, s \right).$$

Our choice of $f_{5.1}$ and $g_{5.1}$ involves the above value m as well as the constants from Lemma 4.1. Specifically, we claim that:

$$f_{5.1} = f_{5.1}(d, l, r, s, t) = f_{4.1} \left(d, (l+1) \binom{2t+2}{2}, r, 2t+2, m \right)$$

and

$$g_{5.1} = g_{5.1}(d, l, r, s, t) = g_{4.1} \left(d, (l+1) \binom{2t+2}{2}, r, 2t+2, m \right)$$

satisfy the lemma.

Let \mathbf{c} be a $(f_{5.1}, g_{5.1})$ -constellation. Assume that 5.1(a) does not hold. By the choice of $f_{5.1}$ and $g_{5.1}$, we can apply Lemma 4.1 to \mathbf{c} . Note that 4.1(a) implies 5.1(a). So we may assume that 4.1(b) holds, that is, there is a d -ample $(m, (l+1) \binom{2t+2}{2})$ -constellation \mathbf{m} which sits \mathbf{c} .

For a $(2t+2)$ -subset T of $S_{\mathbf{m}}$, we say that a path $L \in \mathcal{L}_{\mathbf{m}}$ is T -friendly if for all distinct $x, y \in T$, there is a \mathbf{c} -route R from x to y with $R^* \subseteq L$ such that $T \setminus \{x, y\}$ is anticomplete to R^* . We claim that:

(1) *For every $(2t+2)$ -subset T of $S_{\mathbf{m}}$, the number of T -friendly paths in $\mathcal{L}_{\mathbf{m}}$ is fewer than $\binom{2t+2}{2}$.*

Suppose not. Then for some $T \subseteq S_{\mathbf{m}}$ with $|T| = 2t+2$, there are at least $\binom{2t+2}{2}$ paths in $\mathcal{L}_{\mathbf{m}}$ that are T -friendly. In particular, one may choose $\binom{2t+2}{2}$ pairwise distinct T -friendly paths $(L_{\{x,y\}} : \{x,y\} \in \binom{T}{2})$ in $\mathcal{L}_{\mathbf{m}}$, each assigned to a 2-subset of T . From the definition of a T -friendly path, it follows that for every $\{x,y\} \in \binom{T}{2}$, there is a \mathbf{c} -route $R_{\{x,y\}}$ from x to y with $R_{\{x,y\}}^* \subseteq L_{\{x,y\}}$ such that $T \setminus \{x,y\}$ is anticomplete to $R_{\{x,y\}}^*$. But then the subgraph of \mathbf{m} (and so of \mathbf{c}) induced by

$$\bigcup_{\{x,y\} \in \binom{T}{2}} R_{\{x,y\}}$$

is isomorphic to a proper subdivision of K_{2t+2} , a contradiction to the assumption that 5.1(a) does not hold. This proves (1).

For the rest of the proof, fix a linear order \prec on $S_{\mathbf{m}}$. From (1), we deduce that:

(2) *For every $T = \{x_1, \dots, x_{2t+2}\} \subseteq S_{\mathbf{m}}$ where $x_1 \prec \dots \prec x_{2t+2}$, there is a 2-subset $\{i(T), j(T)\}$ of \mathbb{N}_{2t+2} with $i(T) < j(T)$ as well as and l -subset $\mathcal{L}(T)$ of $\mathcal{L}_{\mathbf{m}}$ for which the following holds. Let R be a \mathbf{c} -route from $x_{i(T)}$ to $x_{j(T)}$ such that $R^* \subseteq V(\mathcal{L}(T))$. Then some vertex in $T \setminus \{x_{i(T)}, x_{j(T)}\}$ has a neighbor in R .*

Let \mathcal{A}_T be the set of all paths in $\mathcal{L}_{\mathbf{m}}$ that are not T -friendly. Then, for every $L \in \mathcal{A}_T$, there is a 2-subset $\{i_{L,T}, j_{L,T}\}$ of \mathbb{N}_{2t+2} with $i_{L,T} < j_{L,T}$ such that for every \mathbf{c} -route R from $x_{i_{L,T}}$ to $x_{j_{L,T}}$ with $R^* \subseteq L$, some vertex in $T \setminus \{x_{i_{L,T}}, x_{j_{L,T}}\}$ has a neighbor in R . Moreover, by (1), we have

$$|\mathcal{A}_T| > |\mathcal{L}_{\mathbf{m}}| - \binom{2t+2}{2} = l \binom{2t+2}{2}.$$

It follows that there exists $\mathcal{L}(T) \subseteq \mathcal{A}_T \subseteq \mathcal{L}_{\mathbf{m}}$ with $|\mathcal{L}(T)| = l$ such that for all distinct $L, L' \in \mathcal{L}(T)$, we have $i_{L,T} = i_{L',T}$ and $j_{L,T} = j_{L',T}$. This proves (2).

Henceforth, for each $(2t+2)$ -subset T of S_m , let $\{i(T), j(T)\} \subseteq \mathbb{N}_{2t+2}$ and $\mathcal{L}(T) \subseteq \mathcal{L}_m$ be as given by (2). Let

$$\Phi : \binom{S_m}{2t+2} \rightarrow \binom{\mathbb{N}_{2t+2}}{2} \times \binom{\mathcal{L}_m}{l}$$

be the map given by $\Phi(T) = (\{i(T), j(T)\}, \mathcal{L}(T))$. By the choice of m , we can apply Theorem 5.2 and deduce that there exist $i, j \in \mathbb{N}_{2t+2}$ with $i < j$, an l -subset \mathcal{L} of \mathcal{L}_m and an s -subset S of S_m , such that for every $(2t+2)$ -subset T of S , we have $i(T) = i, j(T) = j$ and $\mathcal{L}(T) = \mathcal{L}$.

Now, consider the (s, l) -constellation $\mathfrak{b} = (S, \mathcal{L})$. We wish to prove that \mathfrak{b} satisfies 5.1(b). Note that \mathfrak{b} sits in \mathfrak{m} . In particular, \mathfrak{b} is d -ample because \mathfrak{m} is, and \mathfrak{b} sits in \mathfrak{c} because \mathfrak{m} does. It remains to show that \mathfrak{b} is $(2t, 2t, 2t)$ -mixed. To see this, consider the restriction of \preceq to S . Let $x, y \in S$ with $x \prec y$, let R be \mathfrak{c} -route from x to y and let $Q^-, Q, Q^+ \subseteq S_c$ such that $|Q^-| = |Q| = |Q^+| = 2t$ and $Q^- \prec x \prec Q \prec y \prec Q^+$. Recall that $i, j \in \mathbb{N}_{2t+2}$ with $i < j$, and so we may choose $X^- \subseteq Q^-, X \subseteq Q$ and $X^+ \subseteq Q^+$ such that $|X^-| = i - 1, |X| = j - i - 1$ and $|X^+| = 2t + 2 - j$. In particular, we have $X^- \prec x \prec X \prec y \prec X^+$, and

$$T = X^- \cup \{x\} \cup X \cup \{y\} \cup X^+$$

is a $(2t+2)$ -subset of S . Hence, by the choice of $i, j \in \mathbb{N}_{2t+2}$, some vertex in $T \setminus \{x, y\} = X^- \cup X \cup X^+ \subseteq Q^- \cup Q \cup Q^+$ has a neighbor in R . This completes the proof of Lemma 5.1. \blacksquare

6. FROM MIXED TO ZIGZAGGED

In this section, we show that:

Lemma 6.1. *For all $l, l', s, s', t \in \mathbb{N}$ and $\sigma \in \mathbb{N} \cup \{0\}$, there are constants $f_{6.1} = f_{6.1}(l, l', s, s', t, \sigma) \in \mathbb{N}$ and $g_{6.1} = g_{6.1}(l, l', s, s', t, \sigma) \in \mathbb{N}$ with the following property. Let $q \in \mathbb{N}$ and let $q^-, q^+ \in \mathbb{N} \cup \{0\}$ such that $q^- + q^+ = \sigma$. Let \mathfrak{b} be an ample and (q^-, q, q^+) -mixed $(f_{6.1}, g_{6.1})$ -constellation. Then one of the following holds.*

- (a) *There is an induced subgraph of \mathfrak{b} isomorphic to a proper subdivision of K_{2t+2} .*
- (b) *There is an interrupted (s, l) -constellation which sits in \mathfrak{b} .*
- (c) *There is a q -zigzagged (s', l') -constellation which sits in \mathfrak{b} .*

The proof of Lemma 6.1 uses a result from [11], which we state below. By a *hypergraph* we mean a pair $H = (V(H), E(H))$ where $V(H)$ is a finite set, $E(H) \subseteq 2^{V(H)} \setminus \{\emptyset\}$ and $V(H), E(H) \neq \emptyset$. The elements of H are called the *vertices* of H and the elements of $E(H)$ are called the *hyperedges* of H . We need to define three parameters associated with a hypergraph H :

1. Let $\nu(H)$ be the maximum number of pairwise disjoint hyperedges in H .
2. Let $\tau(H)$ be the minimum cardinality of a set $X \subseteq V(H)$ where $e \cap X \neq \emptyset$ for all $e \in E(H)$.
3. Let $\lambda(H)$ be the maximum $k \in \mathbb{N}$ for which there is a k -subset F of $E(H)$ with the following property: for every 2-subset $\{e, e'\}$ of F , some vertex $v_{\{e, e'\}} \in V(H)$ satisfies $\{f \in F : v_{\{e, e'\}} \in f\} = \{e, e'\}$.

(Observe that if the hyperedges of H are pairwise disjoint, then we have $\lambda(H) = 1$, and otherwise we have $\lambda(H) \geq 2$.)

The following is proved in [11]:

Theorem 6.2 (Ding, Seymour and Winkler [11]). *Let $a, a' \in \mathbb{N}$ and let H be a hypergraph with $\nu(H) \leq a$ and $\lambda(H) \leq a'$. Then we have*

$$\tau(H) \leq 11a^2(a + a' + 3) \binom{a + a'}{a'}^2.$$

Theorem 6.2 is often useful for finding certain induced subgraphs. In our setting, hypergraphs H with $\lambda(H)$ large will translate to induced subdivisions of large complete graphs.

We now turn to the proof of the main result in this section:

Proof of Lemma 6.1. Let $l, l', s', t \in \mathbb{N}$ (so all the variables in the statement except for s and σ) be fixed. First, we define two sequences

$$\{a_{s,\sigma} : s \in \mathbb{N}, \sigma \in \mathbb{N} \cup \{0\}\}$$

and

$$\{b_{s,\sigma} : s \in \mathbb{N}, \sigma \in \mathbb{N} \cup \{0\}\}$$

of positive integers recursively, as follows. For every $\sigma \in \mathbb{N} \cup \{0\}$, let

$$a_{1,\sigma} = 1; \quad b_{1,\sigma} = l.$$

For every $s \in \mathbb{N}$, let

$$a_{s,0} = s'; \quad b_{s,0} = l'.$$

For all $s \geq 2$ and $\sigma \geq 1$, assuming $a_{s-1,\sigma}, a_{s,\sigma-1}, b_{s-1,\sigma}$ and $b_{s,\sigma-1}$ are all defined, let

$$a_{s,\sigma} = 11a_{s-1,\sigma}^2(a_{s-1,\sigma} + 2t + 4) \binom{a_{s-1,\sigma} + 2t + 1}{2t + 1}^2 a_{s,\sigma-1}$$

and let

$$b_{s,\sigma} = \binom{a_{s,\sigma}}{a_{s-1,\sigma}} b_{s-1,\sigma} + \binom{a_{s,\sigma}}{a_{s,\sigma-1}} b_{s,\sigma-1}.$$

This concludes the definition of the two sequences. Back to the proof of 6.1, we will prove by induction on $s + \sigma$ that

$$f_{6.1} = f_{6.1}(l, l', s, s', t, \sigma) = a_{s,\sigma} + 1$$

and

$$g_{6.1} = g_{6.1}(l, l', s, s', t, \sigma) = b_{s,\sigma}$$

satisfy the lemma.

Let $q \in \mathbb{N}$ and let $q^-, q^+ \in \mathbb{N} \cup \{0\}$ such that $q^- + q^+ = \sigma$. Let \mathfrak{b} be an ample and (q^-, q, q^+) -mixed $(f_{6.1}, g_{6.1})$ -constellation. Assume that $s = 1$. From the choice of $a_{1,\sigma}$ and $b_{1,\sigma}$, it follows that \mathfrak{b} is a $(1, l)$ -constellation, and so \mathfrak{b} is interrupted, implying that 6.1(b) holds. Next, assume that $\sigma = 0$. Then we have $q^- = q^+ = 0$. From the choice of $a_{s,0}$ and $b_{s,0}$, it follows that \mathfrak{b} is an (s', l') -constellation which is $(0, q, 0)$ -mixed, and so \mathfrak{b} is q -zigzagged, implying that 6.1(b) holds.

We may assume from now on that $s \geq 2$ and $\sigma \geq 1$. Let \preceq be the linear order on $S_{\mathfrak{b}}$ with respect to which \mathfrak{b} is (q^-, q, q^+) -mixed. Since $q^- + q^+ = \sigma \geq 1$, it follows that either $q^- \geq 1$

or $q^+ \geq 1$. Up to the reversal of \preceq , we may assume that $q^+ \geq 1$. Let v be the unique vertex in $S_{\mathfrak{b}}$ with $S_{\mathfrak{b}} \setminus \{v\} \prec v$.

For $L \in \mathcal{L}_{\mathfrak{b}}$, we denote by $N_L(v)$ the set of all neighbors of v in L , and by \mathcal{K}_L the set of all components of $L \setminus N_L(v)$. Observe that \mathcal{K}_L is a set of pairwise disjoint and anticomplete paths in \mathfrak{b} with $|\mathcal{K}_L| \geq 2$, and that v is anticomplete to $V(\mathcal{K}_L)$ in \mathfrak{b} . For each $L \in \mathcal{L}_{\mathfrak{b}}$ and every $x \in S_{\mathfrak{b}} \setminus \{v\}$, let $\mathcal{K}_{x,L}$ be the set of all paths $K \in \mathcal{K}_L$ such that x has a neighbor in K (note that $N_L(x) \subseteq V(\mathcal{K}_L)$ because \mathfrak{b} is ample). For each $L \in \mathcal{L}_{\mathfrak{b}}$, we define the hypergraph H_L with $V(H_L) = \mathcal{K}_L$ and $E(H_L) = \{\mathcal{K}_{x,L} : x \in S_{\mathfrak{b}} \setminus \{v\}\}$.

Let

$$\mathcal{M} = \{L \in \mathcal{L}_{\mathfrak{b}} : \nu(H_L) \geq a_{s-1,\sigma}\}$$

and let

$$\mathcal{N} = \{L \in \mathcal{L}_{\mathfrak{b}} : \nu(H_L) \leq a_{s-1,\sigma}\}.$$

Since $\mathcal{L}_{\mathfrak{b}} = \mathcal{M} \cup \mathcal{N}$, it follows from the choice of $b_{s,\sigma}$ that either

$$|\mathcal{M}| \geq \binom{a_{s,\sigma}}{a_{s-1,\sigma}} b_{s-1,\sigma}$$

or

$$|\mathcal{N}| \geq \binom{a_{s,\sigma}}{a_{s,\sigma-1}} b_{s,\sigma-1}.$$

Assume that the former holds. Then, since $|S_{\mathfrak{b}} \setminus \{v\}| = a_{s,\sigma}$ and from the definition of \mathcal{M} , it follows that there is an $a_{s-1,\sigma}$ -subset S_1 of $S_{\mathfrak{b}} \setminus \{v\}$ as well as a $b_{s-1,\sigma}$ -subset \mathcal{L}_1 of $\mathcal{M} \subseteq \mathcal{L}_{\mathfrak{b}}$ such that for every $L \in \mathcal{L}_1$, the sets $(\mathcal{K}_{x,L} : x \in S_1)$ are pairwise disjoint. Let \mathfrak{b}_1 be the $(a_{s-1,\sigma}, b_{s-1,\sigma})$ -constellation (S_1, \mathcal{L}_1) . Then \mathfrak{b}_1 sits in \mathfrak{b} ; in particular, \mathfrak{b}_1 is ample and (q^-, q, q^+) -mixed, because \mathfrak{b} is. Therefore, by the induction hypothesis applied to \mathfrak{b}_1 , one of the following holds:

- (A1) There is an induced subgraph of \mathfrak{b}_1 isomorphic to a proper subdivision of K_{2t+2} .
- (B1) There is an interrupted $(s-1, l)$ -constellation which sits in \mathfrak{b}_1 .
- (C1) There is a q -zigzagged (s', l') -constellation which sits in \mathfrak{b}_1 .

Since \mathfrak{b}_1 sits in \mathfrak{b} , it follows that (A1) and (C1) imply 6.1(a) and 6.1(c), respectively. So we may assume that (B1) holds, that is, there is an interrupted $(s-1, l)$ -constellation \mathfrak{a} which sits in \mathfrak{b}_1 . Let \preceq_1 be the linear order on $S_{\mathfrak{a}}$ with respect to which \mathfrak{a} is interrupted. Recall that by the choice of S_1 and \mathcal{L}_1 , the vertex v has a neighbor in every \mathfrak{b}_1 -route. Since \mathfrak{a} sits in \mathfrak{b}_1 , it follows that v has a neighbor in every \mathfrak{a} -route. Consequently, $(S_{\mathfrak{a}} \cup \{v\}, \mathcal{L}_{\mathfrak{a}})$ is an (s, l) -constellation which sits in \mathfrak{b} , and extending \preceq_1 from $S_{\mathfrak{a}}$ to $S_{\mathfrak{a}} \cup \{v\}$ by setting $S_{\mathfrak{a}} \prec_1 \{v\}$, it follows that $(S_{\mathfrak{a}} \cup \{v\}, \mathcal{L}_{\mathfrak{a}})$ is interrupted. But then 6.1(b) holds. This completes the induction in the case $|\mathcal{M}| \geq \binom{a_{s,\sigma}}{a_{s-1,\sigma}} b_{s-1,\sigma}$.

Henceforth, assume that $|\mathcal{N}| \geq \binom{a_{s,\sigma}}{a_{s,\sigma-1}} b_{s,\sigma-1}$. As it turns out, we may also assume that $\lambda(H_L) \leq 2t+1$ for every $L \in \mathcal{N}$. In fact, we have:

(3) Suppose that $\lambda(H_L) \geq 2t + 2$ for some $L \in \mathcal{L}_b$. Then 6.1(a) holds.

By the definition of the hypergraph H_L and the parameter λ , there is a $(2t+2)$ -subset X of $S_b \setminus \{v\}$ with the following property: for every 2-subset $\{x, y\}$ of X , there exists $K_{\{x,y\}} \in \mathcal{K}_L$ such that x and y both have neighbors in $K_{\{x,y\}}$ while $X \setminus \{x, y\}$ and $K_{\{x,y\}}$ are anticomplete in \mathfrak{b} . In particular, for every 2-subset $\{x, y\}$ of X , there is a \mathfrak{b} -route $R_{\{x,y\}}$ from x to y with $R^* \subseteq K_{\{x,y\}}$. But now the subgraph of \mathfrak{b} induced by

$$\bigcup_{\{x,y\} \in \binom{X}{2}} R_{\{x,y\}}$$

is isomorphic to a proper subdivision of K_{2t+2} . This proves (3).

From (3), the definition of \mathcal{N} , and Theorem 6.2, it follows that for every $L \in \mathcal{N}$, we have

$$\tau(H_L) \leq 11a_{s-1,\sigma}^2(a_{s-1,\sigma} + 2t + 4) \binom{a_{s-1,\sigma} + 2t + 1}{2t + 1}^2;$$

which, combined with the choice of $a_{s,\sigma}$, yields

$$|S_b \setminus \{v\}| = a_{s,\sigma} \geq \tau(H_L)a_{s,\sigma-1}.$$

In particular, for every $L \in \mathcal{N}$, there is an $a_{s,\sigma-1}$ -subset $S_{2,L}$ of $S_b \setminus \{v\}$ as well as a path $K_L \in \mathcal{K}_L$ such that

$$K_L \in \bigcap_{x \in S_{2,L}} \mathcal{K}_{x,L};$$

that is, every vertex in $S_{2,L}$ has a neighbor in K_L . This, along with the assumption that $|\mathcal{N}| \geq \binom{a_{s,\sigma}}{a_{s,\sigma-1}} b_{s,\sigma-1}$, implies that there is an $a_{s,\sigma-1}$ -subset S_2 of $S_b \setminus \{v\}$ and a $b_{s,\sigma-1}$ -subset \mathcal{L}_2 of $\mathcal{N} \subseteq \mathcal{L}_b$ such that for every $L \in \mathcal{L}_2$, some path $K_L \in \mathcal{K}_L$ satisfies

$$K_L \in \bigcap_{x \in S_2} \mathcal{K}_{x,L}.$$

Again, this means in words that for every $L \in \mathcal{L}$, there is a path $K_L \in \mathcal{K}_L$ such that every vertex in S_2 has a neighbor in K_L . We deduce that $\mathfrak{b}_2 = (S_2, \{K_L : L \in \mathcal{L}_2\})$ is a $(a_{s,\sigma-1}, b_{s,\sigma-1})$ -constellation.

Now, observe that \mathfrak{b}_2 sits in \mathfrak{b} , and so \mathfrak{b}_2 is ample because \mathfrak{b} is. Moreover, recall that \mathfrak{b} is (q^-, q, q^+) -mixed with respect to the linear order \preceq on S_b , where $q^+ \geq 1$. Since $S_2 \subseteq S_b \setminus \{v\} \prec v$ and since v is anticomplete to $V(\mathcal{L}_2)$, it follows that \mathfrak{b}_2 is $(q^-, q, q^+ - 1)$ -mixed with respect to the restriction of \preceq to S_2 . Therefore, since $q^- + (q^+ - 1) = \sigma - 1$, it follows from the induction hypothesis applied to the constellation \mathfrak{b}_2 that one of the following holds.

- (A2) There is an induced subgraph of \mathfrak{b}_2 isomorphic to a proper subdivision of K_{2t+2} .
- (B2) There is an interrupted (s, l) -constellation which sits in \mathfrak{b}_1 .
- (C2) There is a q -zigzagged (s', l') -constellation which sits in \mathfrak{b}_2 .

Since \mathfrak{b}_2 sits in \mathfrak{b} , it follows that (A2) implies 6.1(a), (B2) implies 6.1(b) and (C2) implies 6.1(c). This completes the induction in the case $|\mathcal{N}| \geq \binom{a_{s,\sigma}}{a_{s,\sigma-1}} b_{s,\sigma-1}$, hence completing the proof of Lemma 6.1. \blacksquare

7. FINISHING THE PROOF OF THEOREM 2.1

Finally, let us put everything together and prove our first main result, which we restate:

Theorem 2.1. *For all $d, l, l', r, s, s', t \in \mathbb{N}$, there are constants $f_{2.1} = f_{2.1}(d, l, l', r, s, s', t) \in \mathbb{N}$ and $g_{2.1} = g_{2.1}(d, l, l', r, s, s', t) \in \mathbb{N}$ such that for every $(f_{2.1}, g_{2.1})$ -constellation \mathfrak{c} , one of the following holds.*

- (a) *There is an induced subgraph of \mathfrak{c} isomorphic to either $K_{r,r}$ or a proper subdivision of K_{2t+2} .*
- (b) *There is a d -ample interrupted (s, l) -constellation which sits in \mathfrak{c} .*
- (c) *There is a d -ample $2t$ -zigzagged (s', l') -constellation which sits in \mathfrak{c} .*

Proof. Let

$$f = f_{6.1}(l, l', s, s', t, 4t)$$

and let

$$g = g_{6.1}(l, l', s, s', t, 4t).$$

We claim that

$$f_{2.1} = f_{2.1}(d, l, l', r, s, s', t) = f_{5.1}(d, g, r, f, t)$$

and

$$g_{2.1} = g_{2.1}(d, l, l', r, s, s', t) = g_{5.1}(d, g, r, f, t)$$

satisfy the theorem.

Let \mathfrak{c} be a $(f_{2.1}, g_{2.1})$ -constellation. By the choice of $f_{2.1}$ and $g_{2.1}$, we can apply Lemma 5.1 to \mathfrak{c} and deduce that either 5.1(a) or 5.1(b) holds. The former is identical to 2.1(a), so we may assume that the latter outcome holds, that is, there is a d -ample and $(2t, 2t, 2t)$ -mixed (f, g) -constellation \mathfrak{b} which sits in \mathfrak{c} .

In particular, \mathfrak{b} is ample. Therefore, by the choices of f and g , we can apply Lemma 6.1 to \mathfrak{b} . Again, note that 6.1(a) is identical to 2.1(a). Also, since \mathfrak{b} is d -ample, it follows that the constellations given by both 6.1(b) and 6.1(c) are also be d -ample. Hence, 6.1(b) implies 2.1(b) and 6.1(c) implies 2.1(c). This completes the proof of Theorem 2.1. \blacksquare

8. COMPLETE BIPARTITE INDUCED MINORS

In this section, we prove Theorem 1.3. This, together with Theorem 2.1, implies Theorem 2.3.

For ease of notation, we define the following. Let G be a graph and let $s, t \in \mathbb{N}$. By an *induced (s, t) -model in G* we mean an $(s + t)$ -tuple $M = (A_1, \dots, A_s; B_1, \dots, B_t)$ of pairwise disjoint connected induced subgraphs of G such that:

- A_1, \dots, A_s are pairwise anticomplete in G ;
- B_1, \dots, B_t are pairwise anticomplete in G ; and
- For all $i \in \mathbb{N}_s$ and $j \in \mathbb{N}_t$, the sets A_i and B_j are not anticomplete in G .

It is readily observed that a graph G has an induced minor isomorphic to $K_{s,t}$ if and only if there is an induced (s, t) -model in G .

Let G be a graph and let $M = (A_1, \dots, A_s; B_1, \dots, B_t)$ be an induced (s, t) -model in G . We call the sets $A_1, \dots, A_s, B_1, \dots, B_t$ the *branch sets* of M . We also define

$$A(M) = \bigcup_{i=1}^s A_i$$

and

$$B(M) = \bigcup_{j=1}^t B_j.$$

We say that M is *A-linear* if A_i is a path in G for every $i \in \mathbb{N}_s$. Similarly, we say that M is *B-linear* if B_j is a path in G for every $j \in \mathbb{N}_t$. We say M is *linear* if it is both *A-linear* and *B-linear*.

The proof of Theorem 1.3 has three steps: First, in Theorems 8.1 and 8.2, we reduce the problem to the linear case. Next, in Lemma 8.3, we further simplify the setup using Lemma 4.1. Finally, in Theorem 8.4, we finish the proof; this is only part of the argument in which we exclude $W_{r \times r}$ as an induced minor, rather than a 1-subdivision of a complete graph.

Theorem 8.1. *For all $r, s, t \in \mathbb{N}$, there are constants $f_{8.1} = f_{8.1}(r, s, t) \in \mathbb{N}$ and $g_{8.1} = g_{8.1}(r, s, t) \in \mathbb{N}$ with the following property. Let G be a graph and let $M = (A_1, \dots, A_{f_{8.1}}; B_1, \dots, B_{g_{8.1}})$ be an induced $(f_{8.1}, g_{8.1})$ -model in G . Then one of the following holds.*

- (a) *There is an A-linear induced (s, t) -model $M' = (A'_1, \dots, A'_s; B'_1, \dots, B'_t)$ in G such that for every $i \in \mathbb{N}_t$, we have $B'_i \in \{B_1, \dots, B_{g_{8.1}}\}$.*
- (b) *There is an induced minor of G isomorphic to a 1-subdivision of K_r .*

Proof. Let

$$g_{8.1} = g_{8.1}(r, s, t) = 11(r + 3)r^2t.$$

Let $q = s \binom{g_{8.1}}{t}$ and let

$$f_{8.1} = f_{8.1}(r, s, t) = q \binom{g_{8.1}}{2}.$$

Let G be a graph and let $M = (A_1, \dots, A_{f_{8.1}}; B_1, \dots, B_{g_{8.1}})$ be an induced $(f_{8.1}, g_{8.1})$ -model in G . Suppose for a contradiction that neither 8.1(a) nor 8.1(b) holds. We first partition $\{1, \dots, f_{8.1}\}$ into $\binom{g_{8.1}}{2}$ sets $(X_{i,j} : i, j \in \mathbb{N}_{g_{8.1}}, i < j)$, each of cardinality q .

Fix $i, j \in \mathbb{N}_{g_{8.1}}$ with $i < j$. For every $k \in X_{i,j}$, we let $P_{i,j,k}$ be a path in A_k with $|P_k|$ minimum such that P_k contains a vertex with a neighbor in B_i and a vertex with a neighbor in B_j . Note that such a choice of $P_{i,j,k}$ is possible because A_k is connected and contains both a vertex with a neighbor in B_i and a vertex with a neighbor in B_j . For every $k \in X_{i,j}$, let

$$S_{i,j,k} = \{l \in \mathbb{N}_{g_{8.1}} : P_{i,j,k} \text{ is not anticomplete to } B_l\}.$$

It follows that $\{i, j\} \subseteq S_{i,j,k}$. Moreover, we deduce that:

- (4) *For all $i, j \in \mathbb{N}_{g_{8.1}}$ with $i < j$, there exists $k_{i,j} \in X_{i,j}$ such that $|S_{k_{i,j}}| < t$.*

For suppose there exist $i, j \in \mathbb{N}_{g_{8.1}}$ with $i < j$ such that $|S_k| \geq t$ for all $k \in X_{i,j}$. For each $k \in X_{i,j}$, choose a t -subset T_k of S_k . From the choice of $q = |X_{i,j}|$, it follows that there is

an s -subset Y of $X_{i,j}$ such that $T_k = T_{k'}$ for all $k, k' \in Y$. Let T be the set with $T = T_k$ for all $k \in Y$. But now $(P_k : k \in Y; B_l : l \in T)$ is an A -linear induced (s, t) -model in G that satisfies 8.1(a), a contradiction. This proves (4).

From now on, for each choice of $i, j \in \mathbb{N}_{g_{8.1}}$ with $i < j$, let $k_{i,j} \in X_{i,j}$ be as given by (4), and write $S_{i,j} = S_{i,j,k_{i,j}}$ and $P_{i,j} = P_{i,j,k_{i,j}}$.

Let H be the hypergraph defined as follows. For each choice of $i, j \in \mathbb{N}_{g_{8.1}}$ with $i < j$, the hypergraph H has a vertex $v_{i,j}$, and for every $l \in \mathbb{N}_{g_{8.1}}$, the hypergraph H has a hyperedge $e_l = \{v_{i,j} : l \in S_{i,j}\}$ (so H has $\binom{g_{8.1}}{2}$ vertices and $g_{8.1}$ hyperedges).

We consider the values of $\tau(H)$, $\nu(H)$ and $\lambda(H)$. Since $|S_{i,j}| < t$ for all $i, j \in \mathbb{N}_{g_{8.1}}$ with $i < j$, it follows that every vertex of H belongs to fewer than t hyperedges. In particular, we have $\tau(H) > g_{8.1}/t$, which along with the choice of $g_{8.1}$ implies that:

(5) *We have $\tau(H) > 11(r+3)r^2$.*

Also, recall that for all $i, j \in \mathbb{N}_{g_{8.1}}$ with $i < j$, we have $i, j \in S_{i,j}$, and so $v_{i,j} \in e_i \cap e_j$. In particular,

(6) *We have $\nu(H) = 1$.*

Let us now show that:

(7) *We have $\lambda(H) < r$.*

Suppose not; that is, there is an r -subset F of $\mathbb{N}_{g_{8.1}}$ such that for each choice of $i, j \in F$ with $i < j$, there exist $\alpha_{i,j}, \beta_{i,j} \in \mathbb{N}_{g_{8.1}}$ with $\alpha_{i,j} < \beta_{i,j}$ such that

$$\{l \in F : v_{\alpha_{i,j}, \beta_{i,j}} \in e_l\} = \{i, j\}.$$

By the definition of H , we obtain that

$$F \cap S_{\alpha_{i,j}, \beta_{i,j}} = \{i, j\}.$$

For each choice of $i, j \in F$ with $i < j$, it follows from the definition of $S_{\alpha_{i,j}, \beta_{i,j}}$ that B_i and B_j are the only sets among $(B_l : l \in F)$ to which the path $P_{\alpha_{i,j}, \beta_{i,j}}$ is not anticomplete in G . Moreover, recall that the sets $(B_l : l \in F)$ are connected and pairwise anticomplete in G , and the sets $(P_{\alpha_{i,j}, \beta_{i,j}} : i, j \in F, i < j)$ are connected and pairwise anticomplete in G . Therefore, G has an induced minor isomorphic to a 1-subdivision of K_r , where the sets $(B_l : l \in F)$ correspond to the vertices of K_r and sets $(P_{\alpha_{i,j}, \beta_{i,j}} : i, j \in F, i < j)$ correspond to the $\binom{r}{2}$ vertices obtained by 1-subdividing each edge of K_r . But now 8.1(b) holds, a contradiction. This proves (7).

From (6), (7) and Theorem 6.2 (plugging in $a = 1$ and $a' = r - 1$), it follows that

$$\tau(H) \leq 11a^2(a + a' + 3) \binom{a + a'}{a'}^2 \leq 11(r + 3)r^2.$$

This is a contradiction to (5), and concludes the proof of Theorem 8.1. ■

Let $s, t \in \mathbb{N}$, let G be a graph and let $M = (A_1, \dots, A_s; B_1, \dots, B_t)$ be an induced (s, t) -model in G . We denote by M^T the induced (t, s) -model $(B_1, \dots, B_t; A_1, \dots, A_s)$ in G .

Applying Theorem 8.1 to both sides of the bipartition, we deduce the following:

Theorem 8.2. *For all $r, s, t \in \mathbb{N}$, there are constants $f_{8.2} = f_{8.2}(r, s, t) \in \mathbb{N}$ and $g_{8.2} = g_{8.2}(r, s, t) \in \mathbb{N}$ with the following property. Let G be a graph and assume that there is an induced $(f_{8.2}, g_{8.2})$ -model M in G . Then one of the following holds.*

- (a) *There is a linear induced (s, t) -model in G .*
- (b) *There is an induced minor of G isomorphic to a 1-subdivision of K_r .*

Proof. Let $f_{8.1} = f_{8.1}(r, s, t)$ and $g_{8.1} = g_{8.1}(r, t)$ be as given by Theorem 8.1. Let

$$f_{8.2} = f_{8.2}(r, s, t) = f_{8.1}(r, g_{8.1}, f_{8.1})$$

and let

$$g_{8.2} = g_{8.2}(r, s, t) = g_{8.1}(r, g_{8.1}, f_{8.1}).$$

We apply Theorem 8.1 to the induced $(f_{8.2}, g_{8.2})$ -model M in G . Note that 8.1(b) and 8.2(b) are identical, and so we may assume that 8.1(a) holds. In particular, there is an A -linear induced $(g_{8.1}, f_{8.1})$ -model $M' = (A'_1, \dots, A'_{g_{8.1}}; B'_1, \dots, B'_{f_{8.1}})$ in G . We now apply Theorem 8.1 to the induced $(f_{8.1}, g_{8.1})$ -model $M'^T = (B'_1, \dots, B'_{f_{8.1}}; A'_1, \dots, A'_{g_{8.1}})$ in G . Again, since 8.1(b) and 8.2(b) are identical, we may assume that 8.1(a) holds; that is, there is an A -linear induced (s, t) -model $M'' = (A''_1, \dots, A''_s; B''_1, \dots, B''_t)$ in G such that for all $i \in \mathbb{N}_t$, we have $B''_i \in \{A_1, \dots, A_{g_{8.1}}\}$. Since $A_1, \dots, A_{g_{8.1}}$ are paths in G (as M' is A -linear), it follows that M'' is B -linear. Hence, M'' is both A -linear and B -linear, and so 8.2(a) holds, as desired. \blacksquare

Let $s, t \in \mathbb{N}$ and let G be a graph. Let $M = (A_1, \dots, A_s; B_1, \dots, B_t)$ be a linear induced (s, t) -model in G . By contracting each set A_i to a vertex a_i , we obtain an induced minor of G which is an (s, t) -constellation. We call this constellation the A -contraction of M , and denote it by $\mathbf{c}(M)$. For convenience, we write S_M for $S_{\mathbf{c}(M)} = \{a_i : i \in \mathbb{N}_s\}$ and \mathcal{L}_M for $\mathcal{L}_{\mathbf{c}(M)} = \{B_j : j \in \mathbb{N}_t\}$; so $\mathbf{c}(M) = (S_M, \mathcal{L}_M)$. The B -contraction of M is defined as the A -contraction $\mathbf{c}(M^T)$ of M^T . Analogous to the notion for constellations, for $d \in \mathbb{N}$, we say that M is d - A -ample if $\mathbf{c}(M)$ is d -ample, and M is d - B -ample if $\mathbf{c}(M^T)$ is d -ample. We say that M is d -ample if M is both d - A -ample and d - B -ample.

Lemma 8.3. *For all $d, l, r, r', s \in \mathbb{N}$, there are constants $f_{8.3} = f_{8.3}(d, l, r, r', s) \in \mathbb{N}$ and $g_{8.3} = g_{8.3}(d, l, r, r', s) \in \mathbb{N}$ with the following property. Let G be a graph and let $M = (A_1, \dots, A_{f_{8.3}}; B_1, \dots, B_{g_{8.3}})$ be a linear induced $(f_{8.3}, g_{8.3})$ -model in G . Then one of the following holds.*

- (a) *There is an (r, r) -constellation in G .*
- (b) *There is an induced minor of G which is isomorphic to a 1-subdivision of $K_{r'}$.*
- (c) *There exist subsets $X \subseteq \mathbb{N}_{f_{8.3}}$ and $Y \subseteq \mathbb{N}_{g_{8.3}}$ such that $|X| = s$, $|Y| = l$ and the induced (s, l) -model $(A_i : i \in X; B_j : j \in Y)$ in G is d -ample.*

Proof. Let

$$f_{4.1} = f_{4.1}(d, l, \max\{r, 3\}, r', s)$$

and

$$g_{4.1} = g_{4.1}(d, l, \max\{r, 3\}, r', s)$$

be as given by Lemma 4.1. Let

$$f_{8.3} = f_{8.3}(d, l, r, r', s) = f_{4.1}(d, f_{4.1}, \max\{r, 3\}, r', g_{4.1}).$$

and

$$g_{8.3} = g_{8.3}(d, l, r, r', s) = g_{4.1}(d, f_{4.1}, \max\{r, 3\}, r', g_{4.1})$$

Let G be a graph and let $M = (A_1, \dots, A_{f_{8.3}}; B_1, \dots, B_{g_{8.3}})$ be a linear induced $(f_{8.3}, g_{8.3})$ -model in G . Suppose that 8.3(a) and 8.3(b) do not hold. We show that:

(8) *Neither $\mathfrak{c}(M)$ nor $\mathfrak{c}(M^T)$ has an induced subgraph isomorphic to $K_{\max\{r, 3\}, \max\{r, 3\}}$.*

Suppose not. By symmetry, we may assume $\mathfrak{c}(M)$ has an induced subgraph isomorphic to $K_{\max\{r, 3\}, \max\{r, 3\}}$. Then there are disjoint stable sets $Q, R \subseteq V(\mathfrak{c}(M))$ in $\mathfrak{c}(M)$ such that $|Q| = |R| = \max\{r, 3\}$, and every vertex in Q is adjacent to every vertex in R in $\mathfrak{c}(M)$. Since S_M is a stable set in $\mathfrak{c}(M)$, it follows that either Q or R is disjoint from S_M ; without loss of generality, assume that $R \cap S_M = \emptyset$. Thus, we have $R \subseteq B(M)$; therefore, $R \subseteq B(M)$ is a stable set of cardinality $\max\{r, 3\}$ in G .

Next, we claim that $Q \subseteq S_M$. For suppose there is a vertex $q \in Q \setminus S_M \subseteq \mathfrak{c}(M) \setminus S_M$. Then, since $R \subseteq \mathfrak{c}(M) \setminus S_M$ and since q is complete to R in $\mathfrak{c}(M)$, it follows that q is a vertex of degree at least $|R| = \max\{r, 3\} \geq 3$ in the graph $\mathfrak{c}(M) \setminus S_M$. However, each component of the graph $\mathfrak{c}(M) \setminus S_M$ is path and so $\mathfrak{c}(M) \setminus S_M$ contains no vertex of degree greater than 2, a contradiction. The claim follows.

Now, for each $i \in \mathbb{N}_{f_{8.3}}$, let A_i be the vertex of $\mathfrak{c}(M)$ obtained from contracting A_i . Let $I = \{i \in \mathbb{N}_{f_{8.3}} : a_i \in Q\}$. Then, by the above claim, we have $|I| = |Q| = \max\{r, 3\}$. Moreover, for every $i \in I$, every vertex in R has a neighbour in the path A_i . But now $(R, \{A_i : i \in I\})$ is a $(\max\{r, 3\}, \max\{r, 3\})$ -constellation in G , and so 8.3(a) holds, a contradiction. This proves (8).

(9) *Neither $\mathfrak{c}(M)$ nor $\mathfrak{c}(M^T)$ has an induced subgraph isomorphic to a proper subdivision of $K_{r'}$.*

Suppose not. By symmetry, we may assume that $\mathfrak{c}(M)$ has an induced subgraph isomorphic to a proper subdivision of $K_{r'}$. Since $\mathfrak{c}(M)$ is an induced minor of G , it follows that G has an induced minor isomorphic to a proper subdivision of $K_{r'}$, which in turn implies that G has an induced minor isomorphic to a 1-subdivision of $K_{r'}$. But then 8.3(b) holds, a contradiction. This proves (9).

We apply Lemma 4.1 to the constellation $\mathfrak{c}(M) = (S_M, \mathcal{L}_M)$. From (8) and (9), it follows that the outcome 4.1(a) does not hold. Therefore, 4.1(b) holds; that is, there are subsets $X' \subseteq \mathbb{N}_{f_{8.3}}$ and $Y' \subseteq \mathbb{N}_{g_{8.3}}$ such that:

- $|X'| = g_{4.1}(d, l, \max\{r, 3\}, r', s)$;
- $|Y'| = f_{4.1}(d, l, \max\{r, 3\}, r', s)$; and
- $M' = (A_i : i \in X'; B_j : j \in Y')$ is d - A -ample.

In particular, the third bullet above implies that M^{TT} is d - B -ample. We again apply Lemma 4.1, this time to $\mathfrak{c}(M^{TT})$. Since $\mathfrak{c}(M^{TT})$ is isomorphic to an induced subgraph of $\mathfrak{c}(M^T)$, it follows from (8) and (9) that the outcome 4.1(a) does not hold. So 4.1(b) holds; that is, there are subsets $X \subseteq X'$ and $Y \subseteq Y'$ such that:

- $|X| = l$ and $|Y| = s$; and
- $M'' = (B_j : j \in Y; A_i : i \in X)$ is d - A -ample.

Furthermore, since M^{TT} is d - B -ample, so is M'' . Hence, M'' is d -ample. But now M''^{TT} satisfies the outcome 8.3(c). This completes the proof of Lemma 8.3. \blacksquare

Theorem 8.4. *For every $r \in \mathbb{N}$, there is a constant $f_{8.4} = f_{8.4}(r) \in \mathbb{N}$ with the following property. Let G be a graph and assume that there is a 2-ample linear induced $(f_{8.4}, f_{8.4})$ -model in G . Then G has an induced minor isomorphic to $W_{r \times r}$.*

Proof. Let $f_{8.4} = f_{8.4}(r) = 2f_{1.2}(r) + 1$.

Let $M = (A_1, \dots, A_{f_{8.4}}; B_1, \dots, B_{f_{8.4}})$ be a 2-ample linear induced $(f_{8.4}, f_{8.4})$ -model in G . Let $G_M = G[A(M) \cup B(M)]$. For every subset $X \subseteq A(M) \cup B(M)$, we define

$$I_X = \{i \in \mathbb{N}_{f_{8.4}} : X \cap A_i \neq \emptyset\}$$

and

$$J_X = \{j \in \mathbb{N}_{f_{8.4}} : X \cap B_j \neq \emptyset\}.$$

Let F be the set of all edges of G_M with an end in $A(M)$ and an end in $B(M)$. Let Z be the set of all vertices in G_M that are not incident with any edge in F . Since M is linear, it follows that every vertex in Z has degree at most two in G_M .

Let H be the spanning subgraph of G_M with edge set F . It follows that H is bipartite with bipartition $(A(M), B(M))$, and that Z is the set of all isolated vertices of H . Let \mathcal{K} be the set of all components of $G_M \setminus Z$. Since every vertex in $(A(M) \cup B(M)) \setminus Z$ is incident with an edge in F , it follows that for every $K \in \mathcal{K}$, we have $|I_K|, |J_K| \geq 1$. But we can indeed prove the following:

(10) *For every $K \in \mathcal{K}$, we have $|I_K| = |J_K| = 1$.*

Suppose not. Let $P = p_1 \cdots p_k$ be a shortest path in K such that $|I_P| > 1$ or $|J_P| > 1$. By symmetry, we may assume that $|I_P| > 1$. By the minimality of P , there exist distinct $i, i' \in I_P$ and $j \in J_P$ such that $p_1 \in A_i$ and $p_k \in A_{i'}$, and $p_2, \dots, p_{k-1} \in B_j$. Since M is 2-ample, it follows that $k \geq 5$, and furthermore, $I(N(p_3)) \subseteq \{i\}$. If p_3 has a neighbor $p'_1 \in A_i$, then $p'_1 \in K$ and $p'_1 p_3 \cdots p_k$ contradicts the minimality of P . It follows that p_3 has no neighbors in $A(M)$, and hence $p_3 \in Z$, contrary to the assumption that $p_3 \in K$. This proves (10).

Now, let G' be the graph obtained from G_M by contracting each set $K \in \mathcal{K}$ to a vertex v_K . Then G' is an induced minor of G_M (and so of G) with $V(G') = \{v_K : K \in \mathcal{K}\} \cup Z$. It also follows that:

(11) *No two vertices of degree at least three in G' are adjacent.*

By construction $\{v_K : K \in \mathcal{K}\}$ is a stable set in G' , and the vertices in Z are of degree at most two in G' because they are of degree at most two in G_M . This proves (11).

Furthermore, we deduce that G' has large treewidth:

(12) G' has treewidth at least $(f_{8.4} - 1)/2 = f_{1.2}(r)$.

Let G'' arise from G' by replacing each vertex v_K for $K \in \mathcal{K}$ with two adjacent copies v_K^A and v_K^B . It is straightforward to observe that $\text{tw}(G'') \leq 2 \text{tw}(G') + 1$. Therefore, it suffices to show that $\text{tw}(G'') \geq f_{8.4}$. To that end, for every $i \in \mathbb{N}_{f_{8.4}}$, let

$$A'_i = \{v_K^A : K \in \mathcal{K}, i \in I_K\} \cup (A_i \cap Z)$$

and let

$$B'_i = \{v_K^B : K \in \mathcal{K}, i \in J_K\} \cup (B_i \cap Z).$$

It follows that:

- The sets $(A'_i : i \in \mathbb{N}_{f_{8.4}})$ are connected. This is because A_i is connected and A'_i arose from A_i by identifying the vertices in $A_i \cap K$ for each $K \in \mathcal{K}$. Likewise, the sets $(B'_i : i \in \mathbb{N}_{f_{8.4}})$ are connected.
- The sets $(A'_i : i \in \mathbb{N}_{f_{8.4}})$ are pairwise disjoint. This is due to (10) and the fact that the sets $(A_i : i \in \mathbb{N}_{f_{8.4}})$ are pairwise disjoint. Likewise, the sets $(B'_i : i \in \mathbb{N}_{f_{8.4}})$ are pairwise disjoint.
- For all $i, j \in \mathbb{N}_{f_{8.4}}$, the sets A'_i and B'_j are disjoint (by construction) and not anticomplete in G . The latter holds because G has an edge $ab \in F$ with $a \in A_i$ and $b \in B_j$ (as M is a induced $(f_{8.4}, f_{8.4})$ -model), and so there is a component $K \in \mathcal{K}$ with $a, b \in K$. But now $i \in I_K$ and $j \in J_K$, and $v_K^A \in A'_i$ and $v_K^B \in B'_j$ are adjacent.

From the above three bullets, we conclude that G'' has a minor isomorphic to $K_{f_{8.4}, f_{8.4}}$; thus $\text{tw}(G'') \geq f_{8.4}$. This proves (12).

From (12) and Theorem 1.2, it follows that G' has a subgraph isomorphic to a subdivision of $W_{r \times r}$. Let us denote this subgraph by W . We claim that:

(13) W is an induced subgraph of G' .

Suppose not; let $xy \in E(G')$ with $x, y \in W$ such that x and y are not adjacent in W . Since every vertex of W has degree at least two in W , it follows that x, y are adjacent vertices of degree at least three in G' , contrary to (11). This proves (13).

By contracting some edges of W if needed, we obtain an induced minor of G' isomorphic to $W_{r \times r}$. Since G' is an induced minor of G , we deduce that G has an induced minor isomorphic to $W_{r \times r}$. This completes the proof of Theorem 8.4. \blacksquare

Now Theorem 1.3 follows by combining Theorem 8.2, Lemma 8.3, and Theorem 8.4.

9. THE INTERRUPTED AND THE ZIGZAGGED OUTCOME ARE BOTH NECESSARY

Our goal in this last section is to prove that Theorems 2.1 and 2.3 are “best possible” in the sense that both the interrupted and the zigzagged constellations are unavoidable outcomes of those theorems.

9.1. The interrupted case. For $r \in \mathbb{N}$, we denote by T_r the full binary tree of radius r (on $2^{r+1} - 1$ vertices). It is well-known [18] that for every $r \in \mathbb{N}$, all subdivisions of T_{2r} and their line graphs have *pathwidth* at least r (see [10] for the definition of pathwidth).

Note that all constellations are K_4 -free, and all ample constellations are $K_{3,3}$ -free. Thus, to show that interrupted constellations cannot be omitted from the outcomes of Theorems 2.1 and 2.3, it is enough to prove the following (we remark that the second bullet is not necessary for our purposes here; however, it will be used in a future paper [7]):

Theorem 9.1. *Let \mathfrak{c} be an ample interrupted constellation. Then \mathfrak{c} has no induced subgraph isomorphic to any of the following.*

- An ample q -zigzagged $(3q + 6, 6\binom{q+2}{3})$ -constellation, for $q \in \mathbb{N}$.
- A subdivision of T_7 or the line graph of a subdivision of T_7 .
- A subdivision of $W_{6 \times 6}$ or the line graph of a subdivision of $W_{6 \times 6}$.

The proof of Theorem 9.1 relies on the next two lemmas:

Lemma 9.2. *Let \mathfrak{c} be an ample interrupted constellation. Then for every two anticomplete induced subgraphs X_1, X_2 of \mathfrak{c} , there exists $i \in \{1, 2\}$ such that each component of X_i intersects S_c in at most one vertex.*

Proof. Suppose for a contradiction that there are two anticomplete subsets X_1, X_2 of $V(\mathfrak{c})$ such that for every $i \in \{1, 2\}$, there is a component K_i of $G[X_i]$ with $|K_i \cap S_c| \geq 2$; in particular, $X_i \cap S_c \neq \emptyset$. Let $x_1 \prec \dots \prec x_s$ be the linear order on the vertices in S_c with respect to which \mathfrak{c} is interrupted. Let $j \in \mathbb{N}_s$ be maximum such that $x_j \in X_1 \cup X_2$. Without loss of generality, we may assume that $x_j \in X_1$ (and so $x_j \notin X_2$). Let R be a shortest path in K_2 whose ends $x_i, x_{i'}$ belong to $K_2 \cap S_c$ (note that R exists because K_2 is connected and $|K_2 \cap S_c| \geq 2$). By the choice of R , we have $R^* \subseteq K_2 \setminus S_c \subseteq L$ and by the choice of j , we have $i, i' < j$. But now $x_j \in X_1$ has a neighbor in $R^* \subseteq X_2$ (because \mathfrak{c} is interrupted), a contradiction to the assumption that X_1 and X_2 are anticomplete in \mathfrak{c} . ■

Lemma 9.3. *Let $c, q \in \mathbb{N}$ and let \mathfrak{c} be an ample $(2c + 3q, 2c\binom{c+q-1}{c})$ -constellation which q -zigzagged. Then there are anticomplete subsets X, Y of $V(\mathfrak{c})$ with $\text{tw}(X), \text{tw}(Y) \geq c$.*

Proof. Since $|S_c| = 2c + 3q$, we may choose $x_1, x_2 \in S_c$ and pairwise disjoint subsets $Q, S_1, S_2 \subseteq S_c$ such that $|Q| = q, |S_1| = |S_2| = c + q - 1$ and $x_1 \prec S_1 \prec Q \prec S_2 \prec x_2$.

For each $i \in \{1, 2\}$ and every $L \in \mathcal{L}_c$, let $R_{i,L}$ be a \mathfrak{c} -route from x_i to a vertex in Q with $|R_{i,L}|$ as small as possible. We claim that:

(14) *For every $L \in \mathcal{L}_c$, the sets S_1 and $R_{2,L}^*$ are anticomplete in \mathfrak{c} , and the sets S_2 and $R_{1,L}^*$ are anticomplete in \mathfrak{c} .*

Suppose not. Then, by symmetry, we may assume that some for some $L \in \mathcal{L}_c$, there is a vertex $u \in S_2$ with a neighbor in $R_{1,L}^*$. Let $y \in Q$ be the end of $R_{1,L}$ other than x . Since \mathfrak{c} is ample, it follows that there is a \mathfrak{c} -route R' from x_1 to u with $R'^* \subseteq R_{1,L}^* \setminus N_R(y)$. Since \mathfrak{c} is q -zigzagged, and since $x_1 \prec Q \prec u$, it follows that some vertex $z \in Q$ has a neighbor in R'^* . Consequently, there is a \mathfrak{c} -route R'' from x to $z \in Q_1$ with $R''^* \subseteq R'^* \subseteq R_{1,L}^* \setminus N_R(y)$, and so $|R''| < |R_{1,L}|$. This violates the choice of $R_{1,L}$, hence proving (14).

Let $\mathcal{L}_1, \mathcal{L}_2 \subseteq \mathcal{L}_c$ be disjoint with $|\mathcal{L}_1| = |\mathcal{L}_2| = c \binom{c+q-1}{q}$. Since \mathfrak{c} is q -zigzagged and since $|S_1| = |S_2| = c+q-1$, it follows that for every $i \in \{1, 2\}$ and every $L \in \mathcal{L}_i$, there is a c -subset $S'_{i,L}$ of S_i such that every vertex in $S'_{i,L}$ has a neighbor in $R_{i,L}^*$. Since $|\mathcal{L}_1| = |\mathcal{L}_2| = c \binom{c+q-1}{q}$, it follows that for every $i \in \{1, 2\}$, there is a c -subset S'_i of S_i and a c -subset \mathcal{L}'_i of \mathcal{L}_i such that for every $L \in \mathcal{L}'_i$, we have $S'_{i,L} = S'_i$.

Now, let

$$X = (S'_1, \{R_{1,L}^* : L \in \mathcal{L}'_1\})$$

and let

$$Y = (S'_2, \{R_{2,L}^* : L \in \mathcal{L}'_2\}).$$

By (14) and since \mathfrak{c} is a constellation, it follows that X and Y are anticomplete (c, c) -constellations in \mathfrak{c} . In particular, X and Y are anticomplete induced subgraphs of \mathfrak{c} each with an induced minor isomorphic to $K_{c,c}$. Hence, we have $\text{tw}(X), \text{tw}(Y) \geq c$. This completes the proof of Lemma 9.3. \blacksquare

Proof of Theorem 9.1. By Lemma 9.2, \mathfrak{c} has no two anticomplete induced subgraphs, each of pathwidth more than 2. On the other hand, by Lemma 9.3, for every $q \in \mathbb{N}$, every ample q -zigzagged $(3q+6, 6 \binom{q+2}{3})$ -constellation has two anticomplete induced subgraphs, each of treewidth (and so pathwidth) at least 3. Therefore, \mathfrak{c} has no induced subgraph isomorphic to an ample q -zigzagged $(3q+6, 6 \binom{q+2}{3})$ -constellation, where $q \in \mathbb{N}$.

Moreover, observe that (the line graph of) every subdivision of T_7 has two anticomplete induced subgraphs each isomorphic to (the line graph of) a subdivision of T_6 , and so each of pathwidth at least 3. Thus, \mathfrak{c} has no induced subgraph isomorphic to a subdivision of T_7 or the line graph of a subdivision of T_7 . Finally, observe that (the line graph of) every subdivision of $W_{6 \times 6}$ has two anticomplete induced subgraphs each isomorphic to (the line graph of) a subdivision of $W_{3 \times 3}$, and so each of treewidth (and so pathwidth) at least 3. Hence, \mathfrak{c} has no induced subgraph isomorphic to a subdivision of $W_{6 \times 6}$ or the line graph of a subdivision of $W_{6 \times 6}$. This completes the proof of Theorem 9.1. \blacksquare

9.2. The zigzagged case. Again, since constellations are K_4 -free and ample constellations are $K_{3,3}$ -free, the following suffices to show that zigzagged constellations cannot be omitted from the outcomes of Theorems 2.1 and 2.3:

Theorem 9.4. *Let $q \in \mathbb{N}$ and let \mathfrak{c} be an ample q -zigzagged constellation. Then \mathfrak{c} has no induced subgraph isomorphic to any of the following.*

- An ample interrupted $(2q+6)$ -constellation.
- A subdivision of T_{64q^2} or the line graph of a subdivision of T_{64q^2} .
- A subdivision of $W_{264q^2 \times 264q^2}$ or the line graph of a subdivision of $W_{264q^2 \times 264q^2}$.

The proof is in several steps. First, we show that:

Lemma 9.5. *Let $q \in \mathbb{N}$ and let \mathfrak{c} be an ample q -zigzagged constellation. Then there is no induced subgraph H of \mathfrak{c} isomorphic to a proper subdivision of $K_{1,2q+1}$ where the center and the leaves of H have degree at least 4 in \mathfrak{c} .*

Proof. Suppose not. Then there are paths P_1, \dots, P_{2q+1} of non-zero length in \mathfrak{c} , all sharing an end x , such that $(P_i \setminus \{x\} : i \in \mathbb{N}_{2q+1})$ are pairwise anticomplete in \mathfrak{c} , and for each $i \in \mathbb{N}_{2q+1}$,

the ends of P_i have degree at least 4 in \mathbf{c} . Choose P_1, \dots, P_{2q+1} with $P_1 \cup \dots \cup P_{2q+1}$ minimal subject to the above property. For every $i \in \mathbb{N}_{2q+1}$, let x_i be the end of P_i different than x ; thus, both x and x_i have degree at least 4 in \mathbf{c} . Since \mathbf{c} is ample, it follows that all vertices of degree at least 4 in \mathbf{c} belong to $S_{\mathbf{c}}$. In particular, we have $x, x_1, \dots, x_{2q+1} \in S_{\mathbf{c}}$. Moreover, from the minimality of $P_1 \cup \dots \cup P_{2q+2}$, it follows that for every $i \in \mathbb{N}_{2q+2}$, we have $P_i^* \cap S_{\mathbf{c}} = \emptyset$, and so P_i is \mathbf{c} -route from x to x_i . Let \prec be the linear order on $S_{\mathbf{c}}$ with respect to which \mathbf{c} is q -zigzagged. Since $x, x_1, \dots, x_{2q+1} \in S_{\mathbf{c}}$, it follows that there are $i_1, \dots, i_{q+1} \in \mathbb{N}_{2q+1}$ such that either $x \prec x_{i_1} \prec \dots \prec x_{i_{q+1}}$ or $x \succ x_{i_1} \succ \dots \succ x_{i_{q+1}}$. But now since \mathbf{c} is q -zigzagged, it follows that x_{i_j} has a neighbor in $P_{i_{q+1}}$ for some $j \in \mathbb{N}_q$, a contradiction to the assumption that $(P_i \setminus \{x\}) : i \in \mathbb{N}_{2q+1}$ are pairwise anticomplete in \mathbf{c} . \blacksquare

We need the following result from [13]:

Theorem 9.6 (Groenland, Joret, Nadra, Walczak [13]). *Let $r, t \in \mathbb{N}$ and let G be a graph of treewidth at most $t - 1$. Then either G has pathwidth at most $rt + 1$ or G has a subgraph isomorphic to a subdivision of T_{r+1} .*

Given a graph H , by a *leaf-extension* of H we mean a graph G obtained from H by adding a set S of pairwise non-adjacent vertices, each with exactly one neighbor in $V(H)$.

Lemma 9.7. *Let $w \in \mathbb{N}$, let H be a graph of pathwidth at most w and let G be a subdivision of a leaf-extension of H . Then $\text{pw}(G) \leq 2w^2 + 3w + 2$.*

Proof. Since $\text{tw}(H) \leq \text{pw}(H) \leq w$ and since G is a subdivision of a leaf-extension of H , it follows that $\text{tw}(G) \leq w$. Moreover, since $\text{pw}(H) \leq w$, it follows that H has no subgraph isomorphic to a subdivision of T_{2w} , and so G has no subgraph isomorphic to a subdivision of T_{2w+2} . But now by Theorem 9.6 applied to G , we have $\text{pw}(G) \leq (w + 1)(2w + 1) + 1 = 2w^2 + 3w + 2$. \blacksquare

We continue with more definitions. Let G be a graph. For $X \subseteq V(G)$, the X -*contraction* of G is the graph obtained from G by contracting all edges in G with at least one end in X (and removing all loops and the parallel edges arising from this). It follows that the X -contraction of G is an induced minor of G . By a *star-contraction* of G , we mean the X -contraction of G for some $X \subseteq V(G)$ such that the vertices in X are pairwise at distance at least 3 in G (in particular, X is a stable set in G).

Let \mathbf{c} be a constellation and let H be an induced subgraph of \mathbf{c} . We define \mathbf{c}_H to be the graph with vertex set $V(H) \cap S_{\mathbf{c}}$ such that for all distinct $x, x' \in V(H) \cap S_{\mathbf{c}}$, we have $xx' \in E(\mathbf{c}_H)$ if and only if there is a \mathbf{c} -route R from x to x' where $R \subseteq V(H)$ and $(V(H) \cap S_{\mathbf{c}}) \setminus \{x, x'\}$ is anticomplete to R in H .

We will use the following observation in the proof of Theorem 9.4. Note that 9.8(a) is immediate from the assumption that \mathbf{c} is ample, and 9.8(b) follows easily from the assumption that H has no cycle on four or more vertices; we omit the details.

Observation 9.8. *Let \mathbf{c} be an ample constellation and let H be an induced subgraph of \mathbf{c} with no cycle on four or more vertices. Let J be the $(V(H) \cap S_{\mathbf{c}})$ -contraction of H . Then the following hold.*

- (a) J is a star contraction of H .

(b) J is isomorphic to a subdivision of a leaf-extension of \mathbf{c}_H .

We will also use the next observation (again, we omit the proof as it is easy).

Observation 9.9. *Let $r \in \mathbb{N}$. Then there is an induced subgraph of (the line graph of) $W_{2^r \times 2^r}$ isomorphic to (the line graph of) a proper subdivision of T_r .*

Let us now prove Theorem 9.4:

Proof of Theorem 9.4. First, assume that \mathbf{c} has an induced subgraph \mathbf{b} which is a 1-ample interrupted $(2q + 6)$ -constellation. Let $x_1 \prec \cdots \prec x_{2q+6}$ be the linear order on $S_{\mathbf{b}}$ with respect to which \mathbf{b} is interrupted. Since \mathbf{b} is interrupted, it follows that every vertex in $S_{\mathbf{b}} \setminus \{x_1, x_2, x_3, x_4\}$ has degree at least 4 in \mathbf{b} (and so in \mathbf{c}), and there is a \mathbf{b} -route R_i from x_{2q+6} to x_i , for each $i \in \mathbb{N}_{2q+5}$, such that $(R_i \setminus \{x_{2q+6}\} : i \in \mathbb{N}_{2q+5})$ are pairwise anticomplete in \mathbf{b} (hence in \mathbf{c}). But now $H = \bigcup_{i=5}^{2q+5} P_i$ is an induced subgraph of \mathbf{c} isomorphic to a proper subdivision of $K_{1,2q+1}$ where the center and the leaves of H of degree at least 4 in \mathbf{c} , a contradiction to Lemma 9.5. Therefore, \mathbf{c} has no induced subgraph \mathbf{b} which is a 1-ample interrupted $(2q + 6)$ -constellation.

Next, we prove that \mathbf{c} has no induced subgraph isomorphic to a subdivision of T_{64q^2} or the line graph of a subdivision of T_{64q^2} . This, combined with Observations 9.9, will also imply that \mathbf{c} has no induced subgraph isomorphic to a subdivision of $W_{2^{64q^2} \times 2^{64q^2}}$ or the line graph of a subdivision of $W_{2^{64q^2} \times 2^{64q^2}}$.

Suppose for a contradiction that \mathbf{c} has an induced subgraph isomorphic to (the line graph of) a subdivision of T_{64q^2} . Then there is a (≥ 3) -subdivision T of T_{16q^2} such that \mathbf{c} has an induced subgraph H isomorphic to (the line graph of) T . Let J be the $(V(H) \cap S_{\mathbf{c}})$ -contraction of H . By Observation 9.8(a), J is a star contraction of H . From this and the fact that T is a (≥ 3) -subdivision of T_{16q^2} , it follows that J has an induced minor isomorphic to T_{16q^2} . In particular, we deduce that:

(15) J has pathwidth at least $8q^2$.

Let us now show that:

(16) *There is an enumeration x_1, \dots, x_s of the vertices of \mathbf{c}_H such that if $x_i x_j \in E(\mathbf{c}_H)$ for some $i, j \in \mathbb{N}_s$ with $i < j$, then we have $j - i \leq q$.*

Let \prec be the linear order on $S_{\mathbf{c}}$ with respect to which \mathbf{c} is q -zigzagged, and let $V(\mathbf{c}_H) = V(H) \cap S_{\mathbf{c}} = \{x_1, \dots, x_s\}$ such that $x_1 \prec \cdots \prec x_s$. Suppose that $x_i x_j \in E(\mathbf{c}_H)$ for some $i, j \in \mathbb{N}_s$ with $j > i + q$. Then, by the definition of \mathbf{c}_H , there is a \mathbf{c} -route R with $R \subseteq V(H)$ such that $\{x_{i+1}, \dots, x_{i+q}\} \subseteq (V(H) \cap S_{\mathbf{c}}) \setminus \{x_i, x_j\}$ is anticomplete to R , a contradiction to the assumption that \mathbf{c} is zigzagged. This proves (16).

From (16), we deduce that $\text{pw}(\mathbf{c}_H) \leq q$. Moreover, by Observation 9.8(b), J is isomorphic to a subdivision of a leaf-extension of \mathbf{c}_H , and so by Lemma 9.7, we have $\text{pw}(J) \leq 2q^2 + 3q + 2 < 8q^2$. But this violates (15), hence completing the proof of Theorem 9.4. \blacksquare

10. ACKNOWLEDGEMENTS

This work was partly done during the 2024 Barbados Graph Theory Workshop at the Bellairs Research Institute of McGill University, in Holetown, Barbados. We thank the organizers for the stimulating circumstances of the event. We also thank Bogdan Alecu who worked with us on results in Section 3.

REFERENCES

- [1] Pierre Aboulker, Isolde Adler, Eun Jung Kim, Ni Luh Dewi Sintuari, and Nicolas Trotignon. On the tree-width of even-hole-free graphs. *European Journal of Combinatorics*, 98:103394, 2021.
- [2] Bogdan Alecu, Maria Chudnovsky, Sepehr Hajebi, and Sophie Spirkl. Induced subgraphs and tree decompositions IX. Grid theorem for perforated graphs. Manuscript available at <https://arxiv.org/abs/2305.15615>, 2023.
- [3] Bogdan Alecu, Maria Chudnovsky, Sepehr Hajebi, and Sophie Spirkl. Induced subgraphs and tree decompositions XI. Local structure in even-hole-free graphs of large treewidth. Manuscript available at <https://arxiv.org/abs/2309.04390>, 2023.
- [4] Bogdan Alecu, Maria Chudnovsky, Sepehr Hajebi, and Sophie Spirkl. Induced subgraphs and tree decompositions XII. Grid theorem for pinched graphs. Manuscript available at <https://arxiv.org/abs/2309.12227>, 2023.
- [5] Bogdan Alecu, Maria Chudnovsky, Sepehr Hajebi, and Sophie Spirkl. Induced subgraphs and tree decompositions XIII. Basic obstructions in \mathcal{H} -free graphs for finite \mathcal{H} . Manuscript available at <https://arxiv.org/abs/2311.05066>, 2023.
- [6] Marthe Bonamy, Édouard Bonnet, Hugues Déprés, Louis Esperet, Colin Geniet, Claire Hilaire, Stéphan Thomassé, and Alexandra Wesolek. Sparse graphs with bounded induced cycle packing number have logarithmic treewidth. *J. Combin. Theory Ser. B*, 167:215–249, 2024.
- [7] Maria Chudnovsky, Sepehr Hajebi, and Sophie Spirkl. Induced subgraphs and tree decompositions XVIII. Excluding a forest. Manuscript, 2024.
- [8] Maria Chudnovsky, Meike Hatzel, Tukka Korhonen, Nicolas Trotignon, and Sebastian Wiederrecht. Unavoidable induced subgraphs in graphs with complete bipartite induced minors. Manuscript available at <https://arxiv.org/abs/2405.01879>, 2024.
- [9] James Davies. Appeared in an Oberwolfach technical report, DOI:10.4171/OWR/2022/1.
- [10] Reinhard Diestel. *Graph theory*, volume 173 of *Graduate Texts in Mathematics*. Springer, Berlin, fifth edition, 2018.
- [11] Guoli Ding, Paul Seymour, and Peter Winkler. Bounding the vertex cover number of a hypergraph. *Combinatorica*, 14(1):23–34, 1994.
- [12] Zdeněk Dvořák. Induced subdivisions and bounded expansion. *European J. Combin.*, 69:143–148, 2018.
- [13] Carla Groenland, Gwenaël Joret, Wojciech Nadara, and Bartosz Walczak. Approximating pathwidth for graphs of small treewidth. *ACM Trans. Algorithms*, 19(2), March 2023.
- [14] Sepehr Hajebi. Chordal graphs, even-hole-free graphs and sparse obstructions to bounded treewidth. Manuscript available at <https://arxiv.org/abs/2401.01299>, 2023.
- [15] Sepehr Hajebi. Induced subdivisions with pinned branch vertices. *European J. Combin.*, 124:Paper No. 104072, 2025.
- [16] Andrei Cosmin Pohoata. Unavoidable induced subgraphs of large graphs. Senior thesis, Princeton University, 2014.
- [17] Frank P. Ramsey. On a Problem of Formal Logic. *Proc. London Math. Soc. (2)*, 30(4):264–286, 1929.
- [18] Neil Robertson and P. D. Seymour. Graph minors. I. Excluding a forest. *J. Combin. Theory Ser. B*, 35(1):39–61, 1983.
- [19] Neil Robertson and Paul Seymour. Graph minors. V. Excluding a planar graph. *Journal of Combinatorial Theory, Series B*, 41(1):92–114, 1986.

- [20] Paul Seymour. Five-minute talk. BIRS Workshop on Geometric and Topological Graph Theory (2013-09-30). Recording available at <http://www.birs.ca/events/2013/5-day-workshops/13w5091/videos/watch/201309301509-5minutetalks.html> (see 28:35).
- [21] Ni Luh Dewi Sintuari and Nicolas Trotignon. (Theta, triangle)-free and (even hole, K_4)-free graphs—Part 1: Layered wheels. *Journal of Graph Theory*, 97(4):475–509, 2021.
- [22] Paul Wollan. No big clique minor but a big grid minor. Posted on *Mathoverflow*. <https://mathoverflow.net/q/145201> (version: 2013-10-18).