

EVEN PAIRS IN BERGE GRAPHS WITH NO BALANCED SKEW-PARTITIONS

TARA ABRISHAMI^{*†}, MARIA CHUDNOVSKY^{*¶}, AND YAQIAN TANG^{*¶}

ABSTRACT. Let G be a Berge graph that has no odd prism and no antihole of length at least six as an induced subgraph. We show that every such graph G with no balanced skew-partition is either complete or has an even pair.

1. INTRODUCTION

All graphs in this paper are finite and simple. Let $\chi(G)$ and $\omega(G)$ denote the chromatic number and the clique number of a graph G , respectively. A graph G is *perfect* if every induced subgraph H of G satisfies $\chi(H) = \omega(H)$. The *complement* of a graph G , denoted by \overline{G} , has the same vertex set as G , and two distinct vertices in \overline{G} are adjacent if and only if they are not adjacent in G . A *hole* in a graph G is an induced subgraph isomorphic to a cycle on at least five vertices, and an *antihole* is an induced subgraph whose complement is a hole in \overline{G} . The *length* of a hole (antihole) is equal to the number of its vertices. A graph is *Berge* if it contains no odd hole and no odd antihole as an induced subgraph. In the 1960s, Berge [1] conjectured that a graph is perfect if and only if it is *Berge*. The study of perfect graphs became a major area of research in structural graph theory after Berge's conjecture. In 2002, Chudnovsky, Robertson, Seymour, and Thomas [5] proved the conjecture, which then became known as the *Strong Perfect Graph Theorem (SPGT)*.

An *even pair* in a graph G is a pair $\{u, v\}$ of nonadjacent vertices such that every induced path from u to v in G has an even number of edges. Before the SPGT was proved, many results focused on properties of *minimal imperfect graphs*: imperfect graphs G such that every proper induced subgraph of G is perfect. In particular, Meyniel [15] proved that minimal imperfect graphs do not have an even pair. Also, the proof of the SPGT was simplified by Chudnovsky and Seymour in 2007 using even pairs [7].

A graph G is *complete* if every pair of vertices in G is adjacent. For a vertex $v \in V(G)$, we denote the set of vertices adjacent to v by $N_G(v) = N(v)$. We say a graph G' is obtained by *contracting an even pair* $\{u, v\}$ in G if:

- $V(G') = (V(G) \setminus \{u, v\}) \cup \{w\}$;
- $G' \setminus \{w\} = G \setminus \{u, v\}$; and
- $N_{G'}(w) = N_G(u) \cup N_G(v)$

We denote the graph obtained by contracting the even pair $\{u, v\}$ by $G/\{u, v\}$. A *sequence of contraction* for a graph G is a sequence of graphs G_0, \dots, G_k such that $G_0 = G$, G_k has no even pair, and for all $0 \leq i \leq k - 1$, there exists an even pair $\{u, v\}$ in G_i such that $G_{i+1} = G_i/\{u, v\}$. A graph is *even-contractible* if it has a sequence of contraction with G_k being a complete graph. Fonlupt and Uhry [10] observed that if G is Berge with an even pair $\{u, v\}$, then $G/\{u, v\}$ is also Berge and $\omega(G/\{u, v\}) = \omega(G)$. In particular, given a $\chi(G/\{u, v\})$ -coloring of $G/\{u, v\}$, one can obtain a $\chi(G)$ -coloring of G by preserving the same colors for vertices in $G \setminus \{u, v\}$ and assigning u, v the color of the additional vertex w . Also, an even pair in a Berge graph can be recognized in polynomial time using the algorithm for detecting an odd hole ([6] and [13]): Given a pair of nonadjacent vertices $\{u, v\}$, we add a new vertex p to G such that $N(p) = \{u, v\}$; if the graph $G \cup \{p\}$ contains no odd hole, then $\{u, v\}$ is an even pair. Therefore, we can find a sequence of

contraction G_0, \dots, G_k for a Berge graph G in polynomial time, and thus a $\chi(G)$ -coloring of G can be derived from a $\chi(G_k)$ -coloring of G_k by the procedure above in polynomial time. This algorithm works especially well when G is even-contractile, as the $\chi(G_k)$ -coloring of the complete graph G_k is trivial. Therefore, a natural question is to identify which Berge graph is even-contractile.

To this end, Everett and Reed [9] proposed a conjecture for characterizing even-contractile Berge graphs. A *prism* in a graph G is an induced subgraph that consists of two cliques $\{a_1, a_2, a_3\}$ and $\{b_1, b_2, b_3\}$ and three disjoint paths P_1, P_2, P_3 from a_i to b_i for each i , and with no other edge except for those in the two cliques and in the three paths. The paths P_1, P_2, P_3 are called the *rungs* of the prism. Under these conditions, a prism is *odd* (*even*) if all the three rungs have odd (even) number of edges. Note that every prism in a Berge graph is either even or odd. Everett and Reed conjectured the following:

Conjecture 1.1 ([9]). *A Berge graph with no induced subgraph isomorphic to an antihole of length at least six or an odd prism is even-contractile.*

This conjecture remains open, but several related theorems have been proved. Maffray and Trotignon [14] showed that a Berge graph that has no prism and no antihole of length at least six is even-contractile. Chudnovsky, Maffray, Seymour, and Spirkl [4] showed that if a Berge graph contains no cycle on four vertices and no odd prism of a particular type, then it is either complete or has an even pair.

The main theorem of this paper is the following:

Theorem 1.2. *Let G be a Berge graph with no induced subgraph isomorphic to an antihole of length at least six or an odd prism. If G does not admit a balanced skew-partition, then G is either complete or has an even pair.*

A *balanced skew-partition* is a type of decomposition that appears in the proof of the SPGT. In 2003, Chudnovsky [2] proved a structural decomposition theorem for *trigraphs*, which is a generalization of graphs with possible “undecided” edges called *switchable pairs*. In particular, the theorem implies that a Berge graph either belongs to some “basic” class, or has a balanced skew-partition, or a 2-join, or a 2-join in the complement. Our result is based on this decomposition theorem, and the notion of trigraph is very helpful to the proof.

The remainder of the paper is organized as follows. In Section 2, we introduce the definitions related to trigraphs and present relevant theorems that have been proved. We also define *basic* trigraphs and decompositions, namely balanced skew-partition, 2-join, and the complement of 2-join. In Section 3, we define a class \mathcal{F} of Berge trigraphs and a subclass called *favorable trigraphs* that interact well with the 2-join decomposition. In particular, we will show that almost all trigraphs in \mathcal{F} are favorable when forbidding antihole of length six and balanced skew-partition. In Section 4, we show that basic trigraphs have even pairs, and favorable basic trigraphs have even pairs in certain desirable location. In Section 5, we apply the technique of *block of decompositions* introduced in [8] to handle 2-join and its complement. This technique allows us to decompose any trigraph in \mathcal{F} with no balanced skew-partition into basic trigraphs while keeping track of even pairs. Finally, we prove a generalization of our main theorem 1.2 for trigraphs.

2. TRIGRAPHS

In this paper, we mainly adopt the notation regarding trigraphs from the work by Chudnovsky, Trotignon, Trunck, and Vušković [8]. For the sake of clarity, we restate relevant definitions and introduce new definitions that will appear in the paper.

For a set X , we denote by $\binom{X}{2}$ the set of all subsets of X of size 2. For brevity, an element $\{u, v\}$ of $\binom{X}{2}$ is also denoted by uv , or equivalently, vu . A *trigraph* T consists of a finite vertex set $V(T)$, called the *vertex set* of T , and a map $\theta : \binom{V(T)}{2} \rightarrow \{-1, 0, 1\}$, called the *adjacency*

function of T . Two distinct vertices of T are *strongly adjacent* if $\theta(uv) = 1$, *strongly antiadjacent* if $\theta(uv) = -1$, and *semiadjacent* if $\theta(uv) = 0$. We say that u and v are *adjacent* if $\theta(uv) \in \{0, 1\}$ and *antiadjacent* if $\theta(uv) \in \{0, -1\}$. If u and v are adjacent (antiadjacent), we also say that u is *adjacent* (*antiadjacent*) to v , or u is a *neighbor* (*antineighbor*) of v . Similarly, if u and v are strongly adjacent (strongly antiadjacent), we say u is a *strong neighbor* (*strong antineighbor*) of v . For $v \in V(T)$, let $N(v)$ denote the set of all vertices in $V(T) \setminus \{v\}$ that are adjacent to v , and let $N[v]$ denote $N(v) \cup \{v\}$. An *edge* (*antiedge*) is a pair of adjacent (antiadjacent) vertices. A *switchable pair* is a pair of semiadjacent vertices, and a *strong edge* (*antiedge*) is a pair of strongly adjacent (strongly antiadjacent) vertices. An edge uv (antiedge, strong edge, strong antiedge, switchable pair) is *between* two sets $A \subseteq V(T)$ and $B \subseteq V(T)$ if $u \in A$ and $v \in B$, or if $u \in B$ and $v \in A$.

Let T be a trigraph. The *complement* of T , denoted by \bar{T} , is a trigraph with $V(\bar{T}) = V(T)$ and the adjacency function $\bar{\theta} = -\theta$. Let $A \subset V(T)$ and $b \in V(T) \setminus A$. We say that b is *strongly complete* (*strongly anticomplete*) to A if b is strongly adjacent (strongly antiadjacent) to every vertex of A ; b is *complete* (*anticomplete*) to A if b is adjacent (antiadjacent) to every vertex of A . For two disjoint subsets $A \subset V(T)$ and $B \subset V(T)$, B is *strongly complete* (*strongly anticomplete*, *complete*, *anticomplete*) to A if every vertex of B is strongly complete (strongly anticomplete, complete, anticomplete) to A .

A *clique* of T is set of pairwise adjacent vertices of T , and a *strong clique* is a set of pairwise strongly adjacent vertices of T . A trigraph T is *complete* if $V(T)$ is a clique. A *stable set* of T is a set of pairwise antiadjacent vertices of T . For $X \subseteq V(T)$, the trigraph *induced by T on X* , denoted by $T|X$, has vertex set X and adjacency function $\theta|_X$, the restriction of θ to $\binom{X}{2}$. We denote by $T \setminus X$ the trigraph $T|(V(T) \setminus X)$. Isomorphism between trigraphs is defined in the natural way. For two trigraphs T and H , H is an *induced subtrigraph* of T (or T *contains H as an induced subtrigraph*) if H is isomorphic to $T|X$ for some $X \subseteq V(T)$. Since this paper mainly considers the induced subtrigraph containment relation, we say that T *contains H* if T contains H as an induced subtrigraph.

Let $\eta(T)$ denote the set of all strong edges of T , $\nu(T)$ the set of all strong antiedges of T , $\sigma(T)$ the set of all switchable pairs of T . If $\sigma(T)$ is empty, T is a *graph*. A *semirealization* of T is a trigraph T' with vertex set $V(T)$ that satisfies $\eta(T) \subseteq \eta(T')$ and $\nu(T) \subseteq \nu(T')$. A *realization* of T is any graph that is semirealization of T . For $S \subseteq \sigma(T)$, we denote by G_S^T the realization of T with edge set $\eta(T) \cup S$. The realization $G_{\sigma(T)}^T$ is called the *full realization* of T .

Let T be a trigraph. For $X \subseteq V(T)$, we say that X and $T|X$ are *connected* (*anticonnected*) if the graph $G_{\sigma(T|X)}^{T|X}$ ($G_{\emptyset}^{T|X}$) is connected. A *connected component* (or simply *component*) of X is maximal connected subset of X , and an *anticonnected component* (or simply *anticomponent*) of X is a maximal anticonnected subset of X .

A *path* P of T is a sequence of distinct vertices p_1, \dots, p_k such that either $k = 1$, or for $i, j \in \{1, \dots, k\}$, p_i is adjacent to p_j if $|i - j| = 1$ and p_i is antiadjacent to p_j if $|i - j| > 1$. We say that P is a path *from p_1 to p_k* , and the *endpoints* of P are p_1 and p_k . Under these conditions, let $V(P) = \{p_1, \dots, p_k\}$, the *interior* of P , denoted by P^* , is the induced subtrigraph of P with $V(P^*) = V(P) \setminus \{p_1, p_k\}$, and the *length* of P is $k - 1$. We say P is *even* (*odd*) if it has even (odd) length. Two paths P_1 and P_2 are *disjoint* if $V(P_1) \cap V(P_2) = \emptyset$, and they are *internally disjoint* if $V(P_1^*) \cap V(P_2^*) = \emptyset$; P_1 is a *subpath* of P_2 if P_1 is a connected induced subtrigraph of P_2 . Sometimes we denote P by $p_1 - \dots - p_k$. Notice that, as a graph is also a trigraph, our definition of a path of a graph here is equivalent to a *chordless path* of a graph in some literature.

A *cycle* in a trigraph T is an induced subtrigraph H of T with vertices h_1, \dots, h_k such that $k \geq 3$, and for $i, j \in \{1, \dots, k\}$, h_i is adjacent to h_j if $|i - j| = 1$ or $|i - j| = k - 1$; a *hole* is a cycle that further satisfies that h_i is antiadjacent to h_j if $1 < |i - j| < k - 1$. The *length* of a hole (cycle) is the number of vertices in it. Sometimes we denote H by $h_1 - \dots - h_k - h_1$. An *antipath* (*antihole*) in T is an induced subtrigraph of T whose complement is a path (hole) in \bar{T} .

A *prism* in a trigraph T is an induced subtrigraph H such that the full realization of H is a prism. A trigraph T is *Berge* if it contains no odd hole and no odd antihole. By this definition, T is Berge if and only if \overline{T} is Berge. Also, T is Berge if and only if every realization (semirealization) of T is Berge. An *even pair* in T is a strongly nonadjacent pair $uv \in \binom{V(T)}{2}$ such that every path from u to v in T is even.

2.1. Basic Trigraphs. Here, we define the classes of basic trigraphs. A trigraph T is *bipartite* if its vertex set can be partitioned into two strongly stable sets, called a *bipartition*. A trigraph T is a *line trigraph* if its full realization is the line graph of a bipartite graph and every clique of size at least 3 in T is a strong clique. A trigraph is a *doubled graph* if it has a *good partition*. A good partition is a partition (X, Y) of $V(T)$ satisfying the following:

- Every component of $T|X$ has at most two vertices, and every anticomponent of $T|Y$ has at most two vertices.
- No switchable pair of T is between X and Y .
- For every component C_x of $T|X$ and every anticomponent C_y of $T|Y$, every vertex v of $C_x \cup C_y$ is incident with at most one strong edge and at most one strong antiedge between C_x and C_y .

A trigraph is *basic* if it is either a bipartite trigraph, the complement of a bipartite trigraph, a line trigraph, the complement of a line trigraph, or a doubled trigraph. The following is Theorem 2.3 from [8]:

Theorem 2.1 ([8]). *Basic trigraphs are Berge, and are closed under taking induced subtrigraphs, semirealizations, realizations, and complementation.*

2.2. Decompositions. We now describe the decompositions for trigraphs. First, a *2-join* in a trigraph T is a partition (X_1, X_2) of $V(T)$ such that there exist disjoint sets $A_1, B_1, C_1, A_2, B_2, C_2 \subseteq V(T)$ satisfying:

- $X_1 = A_1 \cup B_1 \cup C_1$ and $X_2 = A_2 \cup B_2 \cup C_2$;
- A_1, A_2, B_1 and B_2 are non-empty;
- no switchable pair is between X_1 and X_2 ;
- every vertex of A_1 is strongly adjacent to every vertex of A_2 , and every vertex of B_1 is strongly adjacent to every vertex of B_2 ;
- there are no other strong edges between X_1 and X_2 ;
- for $i = 1, 2$ $|X_i| \geq 3$; and
- for $i = 1, 2$, if $|A_i| = |B_i| = 1$, then the full realization of $T|X_i$ is not a path of length two containing the members of A_i and B_i .

Under these conditions, we say that $(A_1, B_1, C_1, A_2, B_2, C_2)$ is a *split* of (X_1, X_2) . A 2-join is *proper* if for $i = 1, 2$, every component of $T|X_i$ meets both A_i and B_i . A *complement 2-join* of a trigraph T is a 2-join of \overline{T} . We need the following fact about 2-joins (Theorem 2.4 of [8]):

Theorem 2.2 ([8]). *Let T be a Berge trigraph and $(A_1, B_1, C_1, A_2, B_2, C_2)$ a split of a proper 2-join of T . Then all paths with one end in A_i , one end in B_i and interior in C_i , for $i = 1, 2$, have lengths of the same parity.*

Next, a partition (A, B) of $V(T)$ is a *skew-partition* if A is not connected and B is not anticonnected. A skew-partition (A, B) is *balanced* if there is no odd path of length greater than one with ends in B and interior in A , and there is no odd antipath of length greater than one with ends in A and interior in B . Given a balanced skew-partition (A, B) , the 4-tuple (A_1, A_2, B_1, B_2) is a *split of (A, B)* if A_1, A_2, B_1 , and B_2 are disjoint non-empty sets, $A_1 \cup A_2 = A$, $B_1 \cup B_2 = B$, A_1 is strongly anticomplete to A_2 , and B_1 is strongly complete to B_2 . Note that there exists at least one split for every balanced skew-partition.

When (A, B) is a skew-partition of a trigraph T , we say that B is a *star cutset* of T if at least one anticomponent of B has size one. The following is Theorem 5.9 from [2].

Theorem 2.3 ([2]). *If a Berge trigraph admits a star cutset, then it admits a balanced skew-partition.*

We will often use the following corollary:

Theorem 2.4 ([2]). *If T is a Berge trigraph with no balanced skew-partition, then T does not admit a star cutset.*

3. DECOMPOSING TRIGRAPHS

3.1. Decomposing Trigraphs from \mathcal{F} . In order to handle 2-join partitions and their complements in Section 5, we define a class of trigraphs that will be useful.

Let T be a trigraph. Denote by $\Sigma(T)$ the graph with vertex set $V(T)$ and edge set $\sigma(T)$ (the switchable pairs of T). The connected components of $\Sigma(T)$ are called the *switchable components* of T . Let \mathcal{F} be the class of Berge trigraphs T such that the following hold:

- (1) T has at most one switchable component, and the switchable component D of T has at most two edges.
- (2) If D contains exactly one edge xy , then $N(x) \cap N(y) = \emptyset$ in the trigraph T . In this case, we say it is a *small* switchable component.
- (3) Next, assume that D has two edges. Let $v \in V(T)$ be the vertex of degree two in $\Sigma(T)$, denote its neighbors by x and y . Then v is strongly anticomplete to $V(T) \setminus \{v, x, y\}$ in T , x is strongly antiadjacent to y in T , and $N(x) \cap N(y) = \{v\}$ in T . In this case, we say that the switchable component is *light*.

Our class \mathcal{F} of trigraph is a subclass of the class of the same name studied in [8], and we make use of several of their results.

Theorem 3.1 ([8]). *Let T be a trigraph from \mathcal{F} with no balanced skew-partition, and let $(A_1, B_1, C_1, A_2, B_2, C_2)$ be a split of a 2-join (X_1, X_2) in T . Then the following hold:*

- (i) (X_1, X_2) is a proper 2-join;
- (ii) if $C_i = \emptyset$, then $|A_i| \geq 2$ and $|B_i| \geq 2$, $i = 1, 2$;
- (iii) $|X_i| \geq 4$, $i = 1, 2$.

Theorem 3.2 ([8]). *Every trigraph in \mathcal{F} is either basic, or admits a proper 2-join, or admits a proper 2-join in the complement.*

3.2. Favorable Trigraphs. Let T be a trigraph in \mathcal{F} . We say a pair uv of vertices of T is *disjoint from its switchable component* if D is the switchable component of T and $V(D) \cap \{u, v\}$ is empty. In particular, if the switchable component D of T is empty, every pair of vertices is disjoint from its switchable component. A trigraph $T \in \mathcal{F}$ is *favorable* if it satisfies the following conditions:

- (1) $|V(T)| \geq 5$;
- (2) T has at least one pair of strongly nonadjacent vertices uv disjoint from D ; and
- (3) if D is small and $V(D) = \{x, y\}$, then at least one of $T \setminus (D \cup N(x))$ or $T \setminus (D \cup N(y))$ is not a clique.

A trigraph is *unfavorable* if it is not favorable. By this definition, if T is complete, then T is unfavorable; if T is a graph with at least five vertices and is not complete, then T is favorable as it has empty switchable component. Notice that condition (2) and (3) of being a favorable trigraph are also necessary conditions for trigraphs to have even pairs disjoint from the switchable component.

Next, we will show that, with a few exceptions, a trigraph T in \mathcal{F} with no balanced skew-partition and no antihole is favorable. Further, we prove in section 4 that a basic favorable trigraph has an

even pair disjoint from its switchable component. Both results are essential for handling 2-joins in section 5.

Theorem 3.3. *Let T be a trigraph in \mathcal{F} with no balanced skew-partition and no antihole of length six. If T is unfavorable, then either T is complete or $|V(T)| \leq 5$.*

Proof. We may assume that $|V(T)| > 5$. Let D be the switchable component of T , and let $T' = T \setminus V(D)$ be the induced subtrigraph of T . If D is small, we denote the pair by x and y ; if D is light, we denote the vertex of degree two in $\Sigma(T)$ by v and its neighbors by x and y . Therefore, we can partition $V(T')$ into three sets: $T_1 = T' \setminus (N(x) \cup N(y))$, $T_2 = T' \setminus N(x)$, $T_3 = T' \setminus N(y)$.

First, suppose that D is a light switchable component. Since D is unfavorable, it follows that $V(T) \setminus V(D)$ is a clique. If both T_2 and T_3 are nonempty, then x - a - b - y - v - x with $a \in T_2$ and $b \in T_3$ is a hole of length five, contradicting that T is Berge, so we may assume up to symmetry that $T_3 = \emptyset$. Since $|V(T)| \geq 6$, it follows that $T_1 \cup T_2$ contains three distinct vertices including s and t . If $T_2 = \emptyset$, then $\{s, t\}$ is a star cutset, contradicting 2.4. So we may assume $t \in T_2$. Now, $V(T) \setminus \{v, y, s\}$ is a star cutset, again contradicting 2.4.

Therefore, D is a small switchable component. Now, if $T_1 \neq \emptyset$, T_2 is strongly complete to T_3 since T contains no hole of length five. In this case, $T_2 \cup T_3$ is a star cutset, contradicting 2.4. Thus, $T_1 = \emptyset$. By the definition of unfavorable, since T is not complete, it follows that both T_2 and T_3 are cliques. As T has at least five vertices, at least one of T_2 or T_3 has more than two vertices. Without loss of generality, suppose T_2 contains two distinct vertices. Let s be the vertex in T_2 such that $|N(s) \cap T_3|$ is the maximum, and let t be a vertex in T_2 distinct from s . By 2.4, we may assume $N(s) \cup \{s\} \setminus \{t\}$ is not a star cutset. It follows that there exists a vertex $p \in T_3 \setminus N(s)$ adjacent to t . By maximality of $|N(s) \cap T_3|$, there exists $q \in N(s) \cap T_3$ such that q is not connected to t . Now, $T \setminus \{x, y, s, t, p, q\}$ is an antihole of length of six, a contradiction. This completes the proof. ■

4. EVEN PAIRS IN BASIC TRIGRAPHS

The goal of this section is to prove the following theorem by analyzing each class of basic trigraph:

Theorem 4.1. *Let T be a basic trigraph in \mathcal{F} with no odd prism and no antihole. Then the following statements hold:*

- (1) T is either complete or has an even pair.
- (2) If T is favorable, then T has an even pair disjoint from its switchable component.

4.1. Bipartite Trigraph. Let T be a bipartite trigraph with bipartition (A, B) , where A and B are strongly stable sets. We have the following observation.

Theorem 4.2. *Let T be a bipartite trigraph in \mathcal{F} . Then the following statements hold:*

- (1) T is either complete or has an even pair.
- (2) If T is favorable, then T has an even pair disjoint from its switchable component.

Proof. By the definition of bipartite trigraph, it holds that T is complete or has an even pair, so the first statement holds. For the second statement, suppose that T is favorable and has a nonempty switchable component D . If either $A' = A \setminus V(D)$ or $B' = B \setminus V(D)$ contains at least two vertices, then any two vertices $a_1, a_2 \in A'$ (or $b_1, b_2 \in B'$) form an even pair disjoint from the switchable component, so we may assume that $|A'| = |B'| = 1$. Since T is favorable, it follows that $|V(T)| \geq 5$. Thus $|V(D)| \geq 3$, and so T has a light switchable component, and $|V(T)| = 5$. Assume up to symmetry that $A = \{v, a\}$ and $B = \{x, y, b\}$, where v is the vertex of degree two in D , and x and y are neighbors of v in D . Since T is favorable, it follows that ab is a strong antiedge and b is strongly anticomplete to $V(T) \setminus \{b\}$. Now, a and b are in disjoint connected components of T , so ab is an even pair of T disjoint from the switchable component. ■

4.2. Line Trigraph. Let T be a line trigraph, and let H be the bipartite graph such that its line graph, denoted by $L(H)$, is the full realization of T . Let (A, B) be a bipartition of H . A pair (a_1b_1, a_2b_2) of disjoint edges in H with $a_1, a_2 \in A$ and $b_1, b_2 \in B$ is a *good pair* of H if both of the followings are satisfied:

- Every path P_1 with endpoints a_1 and a_2 satisfies $V(P_1) \cap \{b_1, b_2\} \neq \emptyset$; and
- every path P_2 with endpoints b_1 and b_2 satisfies $V(P_2) \cap \{a_1, a_2\} \neq \emptyset$.

We prove that a good pair in H corresponds to an even pair in T . This is analogous to a result by Hougardy in [12].

Proposition 4.3. *Let H be a bipartite graph, let (a_1b_1, a_2b_2) be a good pair of H , and let u and v be the vertices in $L(H)$ that represent a_1b_1 and a_2b_2 , respectively. Let T be a trigraph such that $L(H)$ is the full realization of T . Then, uv is an even pair in T .*

Proof. First, note that uv is a strong antiedge in T , as a_1b_1 and a_2b_2 are disjoint in H . Suppose that there is an odd path P from u to v in T . Then, P corresponds to an even path Q in H such that $L(Q) = P$. Therefore, up to symmetry, we may assume that $Q = a_1-b_1-\dots-b_2-a_2$, and let Q' be the subpath of Q with endpoints b_1 and b_2 . Thus, $V(Q') \cap \{a_1, a_2\} = \emptyset$. However, this contradicts that (a_1b_1, a_2b_2) is a good pair. ■

Next, we show that forbidding odd prisms guarantees even pairs in line trigraphs. Let H be a bipartite graph. (Note that the following theorems consider all subgraphs of H , which are not necessarily induced subgraphs.) A path Q of H is a *chord path* if its endpoints are contained in the vertex set of a cycle C in H and $V(Q^*) \cap V(C) = \emptyset$. An *even theta* is a graph composed of three internally disjoint even paths with the same endpoints. A *path along the cycle C* is an induced subgraph of C that is a path in $H|C$. A graph is *series-parallel* if and only if it has no subgraph isomorphic to a K_4 -minor.

Proposition 4.4. *Let T be a line trigraph with no odd prism, and let H be a bipartite graph such that $L(H)$ is the full realization of T . Then, H has no subgraph isomorphic to an even theta, and H is series-parallel.*

Proof. Since the line graph of an even theta is an odd prism, it follows that H contains no even theta as a subgraph. As H is bipartite, all cycles in H have even length. It follows that a chord path P of a cycle C in H must have odd length, and the endpoints of P in C divide C into two odd paths along the cycle. Suppose the contrary that H is not series-parallel and thus has a subgraph isomorphic to a K_4 -minor. Since K_4 has maximum degree three, it follows that H has a subgraph J isomorphic to a K_4 -subdivision. Let a, b, c, d be the vertices of degree three of J , and let $P_1, P_2, P_3, P_4, P_5, P_6$ denote the paths with endpoints $(a, b), (b, c), (c, d), (d, a), (b, d),$ and (a, c) , respectively, in J . Notice that each P_i is a chord path, so they are all odd. Now, $P_1 \cup P_4 \cup P_5$ is an odd cycle, contradicting that H is bipartite. ■

Finally, we prove the main result of this subsection.

Theorem 4.5. *Let T be a line trigraph in \mathcal{F} with no odd prism. The following statements hold:*

- (1) T is either complete or has an even pair.
- (2) If T is favorable, then T has an even pair disjoint from its switchable component.

Proof. Let H be a bipartite graph such that $L(H)$ is the full realization of T . We may assume that H is connected and that T is not complete. If T has a nonempty switchable component D , let J be the subgraph of H such that $T|V(L(J)) = D$. In particular, J is a path $p_1-\dots-p_k$ of length either two or three. Thus, following the notation for a path, we call p_1 and p_k the endpoints of J and denote $V(J) \setminus \{p_1, p_k\}$ by $V(J^*)$. Also, note that any vertex $v \in V(J^*)$ has degree at most two in H : Otherwise, the line graph induced by the edges adjacent to v is a clique K of size at least three, and $T|V(K)$ contains a switchable pair, which contradicts the definition of a line trigraph.

By 4.3, to prove the first statement, it suffices to find a good pair in H . Also, to prove the second statement, it suffices to find a good pair in $H \setminus V(J^*)$. Thus, in the following discussion, the proof is completed when the corresponding good pair is found.

Case 1: H is a tree. Since T is not complete, it follows that H is not a star, so H has a path $a_1-b_1-a_2-b_2$ of length three. Now, (a_1b_1, a_2b_2) is a good pair. This proves the first statement for this case.

Next, suppose that T is favorable and has a nonempty switchable component D . Let x and y be the endpoints of J , and let H_x and H_y be the components of $H \setminus V(J^*)$ containing x and y correspondingly. It suffices to show that $H_x \cup H_y$ contains a good pair. If either H_x or H_y contains a path $a_i-b_i-a_j-b_j$ of length three, then (a_ib_i, a_jb_j) is a good pair. Thus, we may assume both H_x and H_y are either empty or isomorphic to a star. If D is small, then T contradicts the third condition of being favorable. So we may assume D is light. By the second condition of being favorable, both H_x and H_y are nonempty. In particular, H_x contains an edge xx' , and H_y contains an edge yy' . Now, (xx', yy') is a good pair. This completes the proof of the second statement for this case.

Case 2: H has a cycle of length at least six. Let $C = a_1-b_1-\cdots-a_k-b_k-a_1$ where $k \geq 6$ be a cycle (not necessarily induced) of maximum length in H . If C has no chord path, then every pair of disjoint edges (a_ib_i, a_jb_j) is a good pair. In particular, as $|E(J)| \leq 3$, there is a good pair in $C \setminus V(J^*)$. Thus, we may assume that C has a chord path P . By 4.4, P is odd and has ends a_i and b_j for $1 \leq i < j \leq k$. Let Q_1 and Q_2 be the two disjoint paths along the cycle C with endpoints a_i and b_j . We may assume by symmetry that $E(J) \cap E(Q_1) = \emptyset$ as any $v \in V(J^*)$ has degree two. Now, to prove both statements for this case, it suffices to show that Q_1 contains a good pair.

Let S_1 be a minimal subpath of Q_1 such that the endpoints of S_1 are joined by a chord path of C , and let this chord path be P' . Thus, S_1 has odd length. If S_1 has length one, then $P' \cup (C \setminus S_1)$ is a longer cycle, a contradiction. So $S_1 = a_t-b_t-\cdots-a_s-b_s$ has length at least three. Further, if there is a chord path P'' of C with exactly one endpoint in $V(S_1^*)$, then $C \cup P' \cup P''$ forms a K_4 minor, contradicting 4.4. Therefore, there is no path in $H \setminus \{a_s, a_t\}$ with endpoints b_s and b_t , and there is no path in $H \setminus \{b_s, b_t\}$ with endpoints a_s and a_t . So (a_tb_t, a_sb_s) is a good pair of H contained in Q_1 . This completes the proof.

Case 3: All the cycles in H have length four. Let $C = a_1-b_1-a_2-b_2-a_1$ be a cycle of length four in H . By 4.4, there is no chord path of C with endpoints a_1 and a_2 (or b_1 and b_2). Also, if there is a chord path with endpoints a_i and b_j with $i, j \in \{1, 2\}$, then G contains a cycle of length greater than four, a contradiction. Thus, there is no path in $H \setminus \{a_1, a_2\}$ with endpoints b_1 and b_2 , and there is no path in $H \setminus \{b_1, b_2\}$ with endpoints a_1 and a_2 . So (a_1b_1, a_2b_2) is a good pair of H . In particular, this proves that in this case every cycle C in H contains a good pair, and thus the first statement follows.

Now, suppose T is favorable with nonempty switchable component D . We may assume that $E(J) \cap E(C) \neq \emptyset$, and $H \setminus V(J^*)$ contains no cycle. As any vertex $v \in V(J^*)$ has degree two, we have $E(J) \subseteq E(C)$ and $V(J) \subseteq V(C)$. Thus, we may assume that $H \setminus V(C)$ is a tree. If $D = \{x, y, v\}$ (where v has degree two in the switchable component) is a light switchable component, then $N(x) \cap N(y) \neq \emptyset$ as the endpoints of J are adjacent, contrary to the fact that $T \in \mathcal{F}$. Therefore, we may assume that D is small and $J = a_1-b_1-a_2$. As T is favorable, $T \setminus V(D)$ is not a clique, which means that there is an edge a_tb_t in $H \setminus V(J^*)$ such that $b_t \neq b_2$. Also, since $b_1 \in V(J^*)$ has degree two in H , we have $b_t \neq b_1$. If $a_t = a_1$, then $\{a_tb_t, a_2b_2\}$ is a good pair in $H \setminus V(J^*)$. Thus, by symmetry, we may assume that $\{a_t, b_t\} \cap V(C) = \emptyset$, and every edge between C and $H \setminus V(C)$ has b_2 as a vertex. In this case, as C has no chord path and $H \setminus V(C)$ is a tree, $\{a_tb_t, a_2b_2\}$ is a good pair in $H \setminus V(J^*)$. This completes the proof. \blacksquare

4.3. Complement of a Bipartite Trigraph and Complement of a Line Trigraph. A *diamond* in a trigraph T is an induced subtrigraph H such that the full realization of H is K_4 minus an edge. A *claw* in a trigraph T is an induced subtrigraph H such that the full realization of H

is the complete bipartite graph $K_{1,3}$. We will need the following characterization of line trigraph, which is a generalization of the main theorem of [11].

Proposition 4.6. *Let T be a line trigraph. Then, T has no induced subtrigraph isomorphic to a diamond or a claw.*

Proof. Suppose that T contains a diamond or a claw, then the full realization of T contains a diamond or a claw as an induced subgraph. By definition, the full realization of T is a line graph. This contradicts to the main theorem of [11], which states that a line graph of a bipartite graph is (claw,diamond)-free. ■

Basic trigraphs which are the complement of a bipartite trigraph and the complement of a line trigraph share the following key property.

Proposition 4.7. *Let T be the complement of a bipartite trigraph or the complement of a line trigraph. Then, a path P of odd length in T has length at most three.*

Proof. First, if T is the complement of a bipartite trigraph, then for all $X \subseteq V(T)$ with $|X| \geq 3$, there exists an edge with both ends in X . Therefore, the path of maximal length in T has length three, so the result follows. Now, we may suppose that T is the complement of a line trigraph, and P is a path of T of length at least five. In this case, $\overline{T|V(P)}$ contains a diamond, which contradicts 4.6. This completes the proof. ■

Proposition 4.8. *Let T be a trigraph in \mathcal{F} such that T is either the complement of a bipartite trigraph or the complement of a line trigraph. If T is favorable, then either T is a graph, or T has an even pair disjoint from the switchable component.*

Proof. Recall that by the definitions, every clique of size at least three is a strong clique in line trigraphs and bipartite trigraphs. If T has a light switchable component $D = \{x, y, v\}$, then, $\overline{T|D}$ is a clique of size three with two switchable pairs, contradicting that T is the complement of a bipartite trigraph or the complement of a line trigraph. Thus, we may assume that T has a small switchable component $D = \{x, y\}$. Suppose that T is the complement of a bipartite trigraph with bipartition (A, B) . By definition, $T|A$ and $T|B$ are strong cliques, so we may assume $x \in A$ and $y \in B$ up to symmetry. In this case, $T \setminus (D \cup N(x)) \subseteq B$ and $T \setminus (D \cup N(y)) \subseteq A$ are both cliques, contradicting that T is favorable.

Now, we may assume that T is the complement of a line trigraph with a small switchable component. If there is a vertex v contained in $T \setminus (N(x) \cup N(y))$, then $\overline{T|\{x, y, v\}}$ is a clique of size three but not a strong clique, contradicting the definition of line trigraph. So $T \setminus (N(x) \cup N(y)) = \emptyset$. If there are two vertices $s, t \in N(x)$ (or $s, t \in N(y)$) such that st is an edge in T , then $\overline{T|\{x, s, t, y\}}$ is a claw, contradicting 4.6. Therefore, $T|N(x)$ and $T|N(y)$ are stable sets. Since T is favorable, we may assume up to symmetry that $|N(x)| \geq 2$. Let s and t be two vertices in $N(x)$, and we claim that $\{s, t\}$ is an even pair: Suppose not, then there is an odd path $s-v_1-v_2-t$ of length three by 4.7. As $T|N(x)$ is a stable set, $\{v_1, v_2\} \subseteq T \setminus (N(x) \cup \{x, y\}) = N(y)$. So v_1v_2 is an edge in $T|N(y)$, contradicting that $T|N(y)$ is a stable set. This completes the proof. ■

The proof of following proposition is inspired by the main idea of [14].

Theorem 4.9. *Let T be a trigraph in \mathcal{F} with no antihole such that T is either the complement of a bipartite trigraph or the complement of a line trigraph. The following statements hold:*

- (1) T is either complete or has an even pair.
- (2) If T is favorable, then T has an even pair disjoint from its switchable component.

Proof. By 4.8, it suffices to prove the first statement. Let T be the vertex-minimal counterexample. We may assume that T is not complete. Let M be a maximal anticonnected set in T such that there are at least two nonadjacent vertices in $V(T) \setminus M$ that are complete to M . Notice that M is

nonempty: since T is not complete, it holds that T contains at least one path of length at least two. Let $C(M)$ be the set of all vertices that are complete to M . By 2.1, each class of basic trigraphs is closed under taking induced subtrigraphs. Thus, since T is minimal, it follows that $C(M)$ has an even pair $\{a, b\}$ as $C(M)$ is not complete by our construction.

Suppose the contrary that $\{a, b\}$ is not an even pair in T . Thus, by 4.7, there is a path $P = a-c-d-b$ of length three in T . Since $\{a, b\}$ is complete to M , it follows that $V(P) \cap M = \emptyset$. First, suppose $\{c, d\} \subseteq V(T) \setminus (M \cup C(M))$. Since both c and d are not in $C(M)$, it follows that c and d each has at least one strong antineighbor in M . So there exists an antipath Q with ends c and d and $Q^* \in M$. Then, $c-Q-d-a-b-c$ is an antihole of length at least five in T , a contradiction. Thus, we may assume up to symmetry that $c \in C(M)$. Since $\{a, b\}$ is an even pair in $C(M)$, it follows that $P \not\subseteq C(M)$, and so $d \in V(T) \setminus (M \cup C(M))$. Now, $M \cup \{d\}$ is also an anticonnected set in T , and $\{c, b\}$ is a pair of nonadjacent vertices complete to $M \cup \{d\}$. This contradicts that M is maximal. Therefore, $\{a, b\}$ is an even pair in T . This completes the proof. \blacksquare

4.4. Doubled Graph. We first state a proposition regarding even pairs in doubled graphs.

Proposition 4.10. *Let T be a doubled graph with good partition (X, Y) . Then the following two statements hold:*

- (1) *Let $C_y = \{y\}$ be an anticomponent of size one in Y . If y has an antineighbor $x \in X$, then xy is an even pair. In particular, if X has an edge x_1x_2 , then T has an even pair.*
- (2) *If $T|Y$ has a strong antiedge y_1y_2 and $T|X$ has no edge, then y_1y_2 is an even pair.*

Proof.

- (1) In this case, y is complete to $N(x)$, so all paths between x and y have length 2. If X contains an edge x_1x_2 , then either x_1 or x_2 is an antineighbor of y , so one of x_1y or x_2y is an even pair.
- (2) In this case, $N(y_1) \cap X$ and $N(y_2) \cap X$ are two disjoint stable sets that partition X . So there is no path from y_1 and y_2 whose interior is contained in X . Since $\{y_1, y_2\}$ is complete to other vertices in Y , all paths from y_1 to y_2 have length 2. \blacksquare

Theorem 4.11. *Let T be a doubled graph in \mathcal{F} with no antihole of length six. The following statements hold:*

- (1) *T is either complete, or has an even pair.*
- (2) *If T is favorable, then T has an even pair disjoint from its switchable component.*

Proof. We may assume that T is not complete and is connected, as any two vertices taken from distinct connected components form an even pair. Let (X, Y) be a good partition of T . Note that by the definition of doubled graph, every switchable component of T is either an edge of $T|X$ or an edge of $T|Y$. If T has a switchable component D , it must be small: otherwise, $T|D$ is both a component and an anticomponent of size 3, contradicting that T is a doubled graph.

Case 1: $T|X$ has a component $C_1 = \{x_1, x_2\}$ of size two. If $T|Y$ is empty, then $T = C_1$ and T is a clique, so we may assume that $T|Y$ is nonempty. If $T|Y$ has two distinct anticomponents C_2, C_3 of size two, $C_1 \cup C_2 \cup C_3$ is an antihole of length six, a contradiction. Therefore, $T|Y$ has at most one anticomponent of size two.

First, suppose $T|Y$ has an anticomponent $C_4 = \{v\}$ of size one. By symmetry, we may assume that v is strongly adjacent to x_1 and strongly antiadjacent to x_2 . By (1) of 4.10, it follows that $\{v, x_2\}$ is an even pair of T , and this proves the first statement for this subcase. Next, assume that T has a small switchable component D . If $\{x_1, x_2\}$ is not the switchable component of T , then $\{v, x_2\}$ is an even pair disjoint from its switchable component, so assume that $\{x_1, x_2\}$ is the switchable component. By the structure of a doubled graph, Y is partitioned by $Y_1 = N(x_1) \setminus \{x_2\}$ and $Y_2 = N(x_2) \setminus \{x_1\}$, and both Y_1 and Y_2 are cliques. Since T is favorable, there must exist a

vertex x_3 in $T|X \setminus \{x_1, x_2\}$ such that x_3 has an antineighbor $y_1 \in T|Y$. If $\{y_1\}$ is an anticomponent of size one in $T|Y$, $\{x_3, y_1\}$ is an even pair disjoint from the switchable component by (1) of 4.10. Therefore, we may assume that y_1 is in the anticomponent $C_5 = \{y_1, y_2\}$ of size two in $T|Y$. As C_5 is the only anticomponent of size two in Y , we may also assume that $X \setminus \{x_1, x_2\}$ is complete to $Y \setminus \{y_1, y_2\}$. By the structure of a doubled graph, $X \setminus \{x_1, x_2\}$ is a nonempty stable set. Now, $\{x_3, y_1\}$ is an even pair disjoint from the switchable component $\{x_1, x_2\}$: y_1 is complete to $N(x) \setminus \{y_2\}$, so a path from x_3 to y_1 either has length two, or is exactly $x_3-y_2-x_2-x_1-y_1$, which has length four. This proves the second statement for this subcase.

Therefore, we may assume that $T|Y$ contains an antiedge y_1y_2 and $Y = \{y_1, y_2\}$. By symmetry, we may assume that x_i is strongly adjacent to y_i for $i = 1, 2$. Notice that all paths from y_1 to y_2 have length three. Every path P from y_1 to x_2 goes through either x_1 or y_2 . If $x_1 \in P$, then $P = y_1-x_1-x_2$ has length two; if $y_2 \in P$, then $P = y_1-\cdots-y_2-x_2$ has length four. Therefore, y_1x_2 is an even pair, and x_1y_2 is also an even pair by symmetry. Thus, this proves the first statement for this subcase. Next, assume that T is favorable. We may also suppose that either $\{x_1, x_2\}$ or $\{y_1, y_2\}$ is the switchable component. As $|V(T)| \geq 5$, there exists a vertex $x_3 \in X \setminus \{x_1, x_2\}$. By symmetry, we may assume x_3 is strongly adjacent to y_1 and strongly antiadjacent to y_2 . Suppose x_3 has a neighbor $x_4 \in X \setminus \{x_1, x_2\}$. Then, x_3y_2 and x_4y_1 are even pairs by the same argument above. Also, $\{x_1, x_3\}$ and $\{x_2, x_4\}$ are even pairs. So at least one of them is disjoint from the switchable component, which is either $\{x_1, x_2\}$ or $\{y_1, y_2\}$. Thus, we may suppose x_3 has no neighbor in X . Then, both $\{x_3, y_2\}$ and $\{x_1, x_3\}$ are even pairs: x_1 is complete to $N(x_3) = \{y_1\}$, and every path from x_3 to y_2 that contains y_1 has length four. So at least one even pair is disjoint from the switchable component. This completes the proof of both statements for Case 1.

Case 2: $T|X$ has no edge. It follows that the switchable component of T is contained in $T|Y$. By (2) of 4.10, we may assume that $T|Y$ is a clique. Suppose that there is no switchable pair in T . Then Y is a strong clique, and thus X is complete to Y . Since T is not complete, there exist nonadjacent vertices $x_1, x_2 \in X$. Now, $\{x_1, x_2\}$ is an even pair of T . Next, assume that there exists a switchable pair y_1y_2 in T . As $T \in \mathcal{F}$, $N(y_1) \cap N(y_2) = \emptyset$, and thus $Y = \{y_1, y_2\}$. Since T is not complete, there exists a vertex $x_1 \in X \cap N(y_1)$. Now, $\{x_1, y_2\}$ is an even pair. This proves the first statement for Case 2.

Suppose T is favorable with switchable pair y_1y_2 . Then, there exists $x_3, x_4 \in X$ such that either $\{x_3, x_4\} \subseteq N(y_1)$ or $\{x_3, x_4\} \subseteq N(y_2)$. In either cases, $\{x_3, x_4\}$ is an even pair because $N(x_3) = N(x_4)$. This completes the proof. \blacksquare

4.5. Proof of Theorem 4.1. By 4.2, 4.5, 4.9, and 4.11, we have checked that the statements hold for each class of basic trigraphs. So 4.1 follows.

5. EVEN PAIRS IN NON-BASIC TRIGRAPHS

5.1. Block of Decomposition. To handle 2-join partitions and their complements, we need the following definitions and a theorem regarding trigraphs with no balanced skew-partition from [8]. A set $X \subseteq V(T)$ is a *fragment* of a trigraph T if $(X, V(T) \setminus X)$ is a proper 2-join of T . A proper 2-join is *even* or *odd* according to the parity of the paths described in 2.2. The *block of decomposition* T_X with respect to a fragment X is defined as follows. Let $X_1 = X$, $X_2 = V(T) \setminus X$, and $(A_1, B_1, C_1, A_2, B_2, C_2)$ be a split of the proper 2-join (X_1, X_2) .

- **Case 1: (X_1, X_2) is odd.** We build the block of decomposition $T_X = T_{X_1}$ as follows: starting with $T|X_1$, we add two vertices a and b , called *marker vertices*, such that a is strongly complete to A_1 , b is strongly complete to B_1 , ab is a switchable pair, and there are no other edges between $\{a, b\}$ and X_1 . Note that $\{a, b\}$ is a small switchable component of T_X , and we call it the *marker component* of T_X .
- **Case 2: (X_1, X_2) is even.** We build the block of decomposition $T_X = T_{X_1}$ as follows: starting with $T|X_1$, we add three vertices a, b, c , called *marker vertices*, such that a is

strongly complete to A_1 , b is strongly complete to B_1 , ac and bc are switchable pairs, and there are no other edges between $\{a, b, c\}$ and X_1 . Note that $\{a, b, c\}$ is a light switchable component of T_X , and we call it the *marker component* of T_X .

In both cases, we say that a and b are the *ends* of the marker component. Again, as our class \mathcal{F} is a subclass of the class of the same name studied in [8], we make use of the following result.

Theorem 5.1 ([8]). *If X is a fragment of a trigraph T in \mathcal{F} with no balanced skew-partition, then the block of decomposition T_X is Berge and has no balanced skew-partition.*

Theorem 5.2. *Let X be a fragment of a trigraph T in \mathcal{F} with no balanced skew-partition, no odd prism, and no antihole. Then the block of decomposition T_X is Berge, and T_X has no balanced skew-partition, no odd prism, and no antihole of length at least five.*

Proof. By 5.1, it suffices to show T_X has no odd prism and no antihole of length at least six. Let M denote the marker component of T_X . Suppose the contrary that T_X has an odd prism Q . If $Q \cap M = \emptyset$, then Q is an odd prism in T , a contradiction. Therefore, we may assume up to symmetry between the ends of the marker component a and b that $a \in V(Q) \cap V(M)$. Suppose that $N(a) \cap V(M) \not\subseteq V(Q)$. Let $y \in A_2$ and let $Q' = (Q \setminus \{a\}) \cup \{y\}$. If $V(Q') \subseteq V(T)$, then Q' is an odd prism of T , a contradiction. Therefore, $M = \{a, v, b\}$, a is not adjacent to b , and $b \in V(Q')$. Let $z \in B_2$ and let $Q'' = (Q' \setminus \{b\}) \cup \{z\}$. Now, Q'' is an odd prism of T , a contradiction. Therefore, $v \in N(a) \cap V(M) \subseteq V(Q)$. As any vertex in Q has degree at least two and $N(v) = \{a, b\}$, $b \in V(Q)$. Thus, $V(M) \subseteq V(Q)$, and M is contained in a rung of the prism Q . Let $(A_1, C_1, B_1, A_2, C_2, B_2)$ be a split of $(X, V(T) \setminus X)$. Since $(X, V(T) \setminus X)$ is a proper 2-join, it follows that there is a path P of T with ends in A_2 and B_2 and interior in C_2 such that P has the same parity as M . Now, $(Q \setminus M) \cup P$ is an odd prism of T , a contradiction. Therefore, T_X does not contain an odd prism.

Next, suppose that T_X contains an antihole, and let $H = v_1 \cdots v_k v_1$ be the shortest antihole in T_X . Since T_X is Berge and an antihole of length six is an odd prism, we may assume that $k \geq 7$. When T_X has the marker component $\{a, b, c\}$, it follows that $c \notin V(H)$ because c is strongly anticomplete to $T_X|X$. If $|V(H) \cap V(M)| = 1$, we may assume by symmetry that $a \in V(H) \cap V(M)$. Now, we may replace a by a vertex $a' \in A_2$, and $T|(V(H)) \setminus \{a\} \cup \{a'\}$ is an antihole of the same length in T , a contradiction. Therefore, we may assume that $V(H) \cap V(M) = \{a, b\}$, and let $a = v_i$ and $b = v_j$ where $i < j$. First, suppose ab is an antiedge of the antihole such that $i - j = 1$ or $i - j = k - 1$. Because $\{a, b\}$ is strongly anticomplete to C_1 , it follows that no vertex of the antihole is contained in C_1 . Also, at most one vertex of H is in A_1 and at most one vertex of H is in B_1 , since a is strongly anticomplete to B_1 and b is strongly anticomplete to A_1 . Therefore, $k \leq 4$, a contradiction. Hence, we may suppose that $1 < i - j < k - 1$. However, as $k \geq 7$, at least one of $v_i v_{i+1} \cdots v_j v_i$ or $v_j v_{j+1} \cdots v_i v_j$ is an antihole and has length less than k , which is a contradiction that H is the shortest antihole in T_X . Therefore, T_X does not contain an antihole. This completes the proof. \blacksquare

Theorem 5.3. *Let X be a fragment of a trigraph T in \mathcal{F} . If the block of decomposition T_X has an even pair uv disjoint from its marker component, then uv is also an even pair in T .*

Proof. Let $(A_1, B_1, C_1, A_2, B_2, C_2)$ be a split of a proper two join (X_1, X_2) with $X = X_1$. Suppose that there is a path P from u to v in T of odd length. If $P \subseteq T|X_1$, then P is a path of T_X , a contradiction, so $P \cap (T|X_2)$ is not empty. First, suppose $P \cap (T|X_2)$ is a path Q of X_2 from A_2 to B_2 . Let a and b be the ends of the marker component of T_X . By construction of the marker component, it follows that the path from a to b in the marker component of T_X has the same parity as Q . Let Q' be the path from a to b in the marker component of T_X . Now, $P' = (P \setminus Q) \cup Q'$ induces a path of T_X of the same parity as P , a contradiction.

Therefore, if $Q = P \cap (T|X_2)$ is a path, we may assume up to symmetry that the endpoints of $P \cap (T|X_2)$ are contained in A_2 . However, as A_1 is strongly complete to A_2 , any vertex a_1 in A_1

induces a cycle with Q , contradicting that P is a path with endpoints in X_1 . Thus, no edge of P is contained in $T|X_2$, and either $P \cap (T|X_2) = \{a_2, b_2\}$, or $P \cap (T|X_2) = \{a_2\}$, or $P \cap (T|X_2) = \{b_2\}$, where $a_2 \in A_2$ and $b_2 \in B_2$. Let $P' \subseteq T_X$ be the path obtained by replacing a_2 with the marker vertex a if $P \cap A_2 \neq \emptyset$, and replacing b_2 with the marker vertex b if $P \cap B_2 \neq \emptyset$. Now, P' is a path of T_X of the same parity as P , a contradiction. \blacksquare

5.2. Handling 2-joins and their complements. First, we show that it remains to consider trigraphs that admit proper 2-joins.

Theorem 5.4. *Let T be a trigraph in \mathcal{F} with no balanced skew-partition and no antihole. Then, either T is basic, or T admits a proper 2-join.*

Proof. By Lemma 3.2, we may assume that \bar{T} admits a proper 2-join (X_1, X_2) with split $(A_1, C_1, B_1, A_2, C_2, B_2)$. By the definition of balanced skew-partition, a trigraph admits a balanced skew-partition if and only if its complement admits a balanced skew-partition. Thus, by Lemma 2.3, \bar{T} has no star cutset.

Suppose that $C_1 \neq \emptyset$. Let D_1 be an anticomponent of C_1 . Then D_1 is strongly complete to $X_2 \cup (C_1 \setminus D_1)$. Let A'_1 be the set of vertices in A_1 that are not strongly complete to D_1 , and define B'_1 similarly. If $B'_1 = \emptyset$, then for every $a_2 \in A_2$, $A_1 \cup \{a_2\}$ is a star cutset in \bar{T} , a contradiction. Thus $A'_1 \neq \emptyset$, and $B'_2 \neq \emptyset$. If some $a_1 \in A'_1$ has a neighbor $b_1 \in B'_1$, then there is an antipath of length at least two from a_1 to b_1 with interior in D_1 , that can be completed via X_2 to an antihole in T , a contradiction. It follows that A'_1 is strongly anticomplete to B'_1 . Now $(A'_1 \cup B'_1, X_2 \cup (X_1 \setminus (A'_1 \cup B'_1)))$ is a skew-partition of T . We will obtain a contradiction by showing that this is a balanced skew-partition. Suppose there is an odd path P with ends $u, v \in X_2 \cup (X_1 \setminus (A'_1 \cup B'_1))$ and interior in $A'_1 \cup B'_1$. We may assume that $V(P^*) \subseteq A'_1$. Since $B_2 \cup C_2$ is strongly complete to A'_1 , and A_2 is strongly anticomplete to A'_1 , it follows that $u, v \in X_1$. Since $u-P^*-v-a_2-u$ is not an odd hole in T for any $a_2 \in A_2$, it follows that $A_1 \cap \{u, v\} \neq \emptyset$. Since D_1 is complete to $A_1 \setminus A'_1$, we deduce that $v \notin D_1$, and so D_1 is strongly complete to $\{u, v\}$. Now P is an odd path with ends strongly complete to D_1 and with no vertex strongly complete to D_1 in its interior, and every $a_2 \in A_2$ is strongly complete to D_1 and strongly anticomplete to P , contrary to Theorem 5.1 of [3]. Next suppose that there is an odd antipath Q with ends $u, v \in A'_1 \cup B'_1$ and interior in $X_2 \cup (X_1 \setminus (A'_1 \cup B'_1))$. Since u is adjacent to v , we may assume that $u, v \in A'_1$. Since $u-Q^*-v-a_2-u$ is not an antihole in T for any $a_2 \in A_2$, it follows that $V(Q^*) \not\subseteq D_1$. Since $V(Q^*)$ is anticonnected, there is an anticomponent D_2 of $X_2 \cup (X_1 \setminus (A'_1 \cup B'_1))$ such that $V(Q^*) \subseteq D_2$, and $D_2 \neq D_1$. Since $u, v \in A'_1$, there is an antipath Q' with ends u, v and interior in D_1 . But now $u-Q^*-v-Q'^*-u$ is an antihole in T , a contradiction.

This proves that $C_1 = \emptyset$, and $C_2 = \emptyset$ by symmetry. Now, $(A_1, C_1, B_1, B_2, C_2, A_2)$ is a split of a proper 2-join of T . This completes the proof. \blacksquare

Theorem 5.5. *Let T be a trigraph in \mathcal{F} with no balanced skew partition, no odd prism, and no antihole. If T admits a proper 2-join, then T is either complete or has an even pair disjoint from its switchable component.*

Proof. Let T be the vertex-minimal counterexample to the claim. Let $(A_1, B_1, C_1, A_2, B_2, C_2)$ be a split of a proper 2-join of T with $X_1 = A_1 \cup B_1 \cup C_1$ and $X_2 = A_2 \cup B_2 \cup C_2$. If T has a switchable component D , we may assume up to symmetry that $V(D) \subseteq X_2$, as no switchable pair meets both X_1 and X_2 . Then, the only switchable component of T_{X_1} is its marker component. By 5.2, $T_{X_1} \in \mathcal{F}$, and T_{X_1} admits no balanced skew-partition, no antihole, and no odd prism.

Next, we show that T_{X_1} is not complete: if C_1 is not empty, then a vertex $y \in C_1$ is not adjacent to any vertex of the marker component of T_{X_1} ; if C_1 is empty, then by 3.1, $|A_1| \geq 2$ and $T_{X_1}|A_1$ is strongly anticomplete to b . Now, we show that T_{X_1} has an even pair disjoint from its switchable component. By 5.4, we may assume that T_{X_1} is basic as T is the vertex-minimal counterexample. By 3.1, $|X_i| \geq 4$ for $i = 1, 2$, and the marker component of T_{X_1} has either two or three vertices

(depending on the parity of the 2-join (X_1, X_2)). In either case, we have $|T_{X_1}| \geq 6$. By 3.3, T_{X_1} is favorable as it is not complete. Therefore, T_{X_1} has an even pair $\{u, v\}$ disjoint from its marker component by 4.1. However, by 5.3, $\{u, v\}$ is also an even pair in T , a contradiction. ■

5.3. Proof of the main theorem. Now, we are ready to prove 1.2. We restate it here for the sake of clarity.

Theorem 5.6 (1.2). *If G is a Berge graph with no odd prism and no antihole, and G does not admit a balanced skew-partition, then G is either complete or has an even pair.*

Proof. First, $G \in \mathcal{F}$ as G is Berge and has no switchable component. So the result follows from 5.4, 5.5, and 4.1 ■

REFERENCES

- [1] C. Berge. Färbung von Graphen, deren sämtliche bzw. deren ungerade Kreise starr sind. *Wissenschaftliche Zeitschrift der Martin-Luther-Universität Halle-Wittenberg, Mathematisch-Naturwissenschaftliche Reihe*, 10, 1961.
- [2] M. Chudnovsky. *Berge trigraphs and their applications*. PhD thesis, Princeton University, 2003.
- [3] M. Chudnovsky. Berge trigraphs. *Journal of Graph Theory*, 53:1–82, 2006.
- [4] M. Chudnovsky, F. Maffray, P. Seymour, and S. Spirkl. Even pairs and prism corners in square-free berge graphs. *Journal of Combinatorial Theory, Series B*, 131:12–39, 2018.
- [5] M. Chudnovsky, N. Robertson, P. Seymour, and R. Thomas. The strong perfect graph theorem. *Annals of Mathematics*, 164(1):51–229, 2006.
- [6] M. Chudnovsky, A. Scott, P. Seymour, and S. Spirkl. Detecting an odd hole. *J. ACM*, 67(1), jan 2020.
- [7] M. Chudnovsky and P. Seymour. Even pairs in berge graphs. *Journal of Combinatorial Theory, Series B*, 99(2):370–377, 2009.
- [8] M. Chudnovsky, N. Trotignon, T. Trunck, and K. Vušković. Coloring perfect graphs with no balanced skew-partitions. *Journal of Combinatorial Theory, Series B*, 115:26–65, 2015.
- [9] H. Everett, C. de Figueiredo, C. L. Sales, F. Maffray, and O. Porto. Even pairs. In J. L. Ramírez-Alfonsín and B. A. Reed, editors, *Perfect Graphs*, pages 67–92. Wiley Interscience, 2001.
- [10] J. Fonlupt and J. Uhry. Transformations which preserve perfectness and h-perfectness of graphs. *Annals of Discrete Mathematics*, 66:83–95, 1982.
- [11] F. Harary and C. Holzman. Line graphs of bipartite graphs. *Rev. Soc. Mat. Chile*, 1(1):19–22, 1974.
- [12] S. Hougardy. Even and odd pairs in linegraphs of bipartite graphs. *European Journal of Combinatorics*, 16(1):17–21, 1995.
- [13] K.-Y. Lai, H.-I. Lu, and M. Thorup. Three-in-a-tree in near linear time. In *Proceedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing*, STOC 2020, page 1279–1292, New York, NY, USA, 2020. Association for Computing Machinery.
- [14] F. Maffray and N. Trotignon. A class of perfectly contractile graphs. *Journal of Combinatorial Theory, Series B*, 96(1):1–19, 2006.
- [15] H. Meyniel. A new property of critical imperfect graphs and some consequences. *European Journal of Combinatorics*, 8:313–316, 1987.

*PRINCETON UNIVERSITY, PRINCETON, NJ, USA

[†] SUPPORTED BY NSF-EPSRC GRANT DMS-2120644.

[¶] SUPPORTED BY NSF-EPSRC GRANT DMS-2120644 AND BY AFOSR GRANT FA9550-22-1-0083.

[‡] SUPPORTED BY AFOSR GRANT FA9550-22-1-0083.