

Perfect Matchings in Planar Cubic Graphs

Maria Chudnovsky¹
Columbia University, New York, NY 10027

Paul Seymour²
Princeton University, Princeton, NJ 08544

January 24, 2008; revised July 21, 2009

¹This research was conducted while the author served as a Clay Mathematics Institute Research Fellow.

²Supported by ONR grant N00014-01-1-0608 and NSF grant DMS-0070912.

Abstract

A well-known conjecture of Lovász and Plummer from the mid-1970's, still open, asserts that for every cubic graph G with no cutedge, the number of perfect matchings in G is exponential in $|V(G)|$. In this paper we prove the conjecture for planar graphs; we prove that if G is a planar cubic graph with no cutedge, then G has at least

$$2^{|V(G)|/655978752}$$

perfect matchings.

1 Introduction

In 1998, Schrijver [4] proved the “Schrijver-Valiant conjecture”, a lower bound on the number of perfect matchings in a k -regular bipartite graph. A consequence of this is that every cubic bipartite graph has exponentially many perfect matchings.

But what about non-bipartite cubic graphs? They need not have any perfect matchings at all, so let us confine ourselves to cubic graphs without cutedges. In that case, Lovász and Plummer [3] conjectured in the mid-1970’s that again such a graph G must have exponentially many perfect matchings. This has proved to be a challenging question, and is still open; and the best lower bound currently known is $|V(G)|/2 + 1$ (except for one graph with twelve vertices and only six perfect matchings), proved recently by Kral, Sereni and Stiebitz [2]. In this paper we prove the conjecture for planar cubic graphs, the following:

1.1 *Let G be a planar cubic graph with no cutedge. Then G has at least*

$$2^{\lfloor |V(G)|/655978752 \rfloor}$$

perfect matchings.

Planarity provides two advantages. First, we have the four-colour theorem, which tells us that planar cubic graphs without cutedges are 3-edge-colourable, and thus provides us with a way to produce triples of perfect matchings that cover all the edges. (However, it is *not* true that these graphs necessarily have exponentially many 3-edge-colourings — unlike bipartite cubic graphs, which do — so we need to use the four-colour theorem in an indirect way.) The second advantage of planarity is that we have regions, a source of cycles of bounded length whose deletion does not reduce the connectivity of the graph very much; and these will be key to the proof.

The proof breaks into two parts; first we prove the statement for cyclically 4-edge-connected planar cubic graphs, and then we deduce it for general (planar) cubic graphs.

The idea in the cyclically 4-edge-connected case is, we first find linearly many even length cycles, disjoint and carefully chosen (ideally we would like them far apart, and each bounding either one region or the union of two regions). This set of cycles has exponentially many subsets; let X be one of its subsets. For each cycle in X , delete its even edges and replace its odd edges by pairs of parallel edges. (Let us call this “flipping” the cycle.) We arrange that for each X the graph we produce will be a planar cubic graph with no cutedges, and will therefore be 3-edge-colourable, because of the four-colour theorem. Consequently there will be a triple of perfect matchings of the original graph, such that we can reconstruct X from this triple. This shows that the original graph had exponentially many triples of perfect matchings, and therefore exponentially many perfect matchings. The problem here is that we cannot necessarily find such a large set of even cycles sufficiently far apart for them all to be flippable independently without producing cutedges, and we sometimes have to make do with something a little less (we make use of the fact that there are two ways to flip a cycle, depending on which we designate as its “even” edges, and we only need one of the two ways to work).

The second half of the proof, when the graph is not cyclically 4-edge-connected, is quite non-trivial, rather surprisingly so, and it is the proof for this case that makes the constant in our main result so large. We find it necessary to retain the stronger statement proved in the cyclically 4-edge-connected case, that we have linearly many disjoint cycles that can all be independently “flipped”, rather than just that there are exponentially many perfect matchings.

2 Good looks, bracelets, and perfect matchings

Let us be more precise. Graphs in this paper are all finite and loopless, but not necessarily simple (they may have parallel edges). For a graph G , if $X \subseteq V(G)$ we define $\delta_G(X) = \delta(X)$ to be the set of edges of G with one end in X and one end in $V(G) \setminus X$; and if $v \in V(G)$ we write $\delta(v)$ for $\delta(\{v\})$. A *cut* in G is a set of edges D such that $D = \delta(X)$ for some $X \subseteq V(G)$. If $w : E(G) \rightarrow \{-1, 0, 1\}$ and $Y \subseteq E(G)$, we define $w(Y)$ to be the sum of $w(e)$ for all $e \in Y$. If G is cubic and $w : E(G) \rightarrow \{-1, 0, 1\}$ satisfies $w(\delta(v)) = 0$ for each $v \in V(G)$, we call w a *look* of G . A look w is said to be *good* if for every cut D , $w(D) \neq 1 - |D|$. (In other words, if we delete from G all the edges e with $w(e) = -1$, and f is a cutedge of the graph we produce, then $w(f) = 1$.)

We observe first that:

2.1 *Let G be a planar cubic graph with k good looks. Then G has at least $k^{\frac{1}{3}}$ perfect matchings.*

Proof. Let w be a good look of G . Let H be obtained from G by deleting all edges e with $w(e) = -1$, and adding an edge parallel to every edge e with $w(e) = 1$. Then H is also planar and cubic, and since w is a good look of G it follows that H has no cutedge. By the four-colour theorem, there are three perfect matchings in H such that every edge of H is in exactly one of them. Consequently there are three perfect matchings in G , say F_1, F_2, F_3 , such that every edge e is in $1 + w(e)$ of them. In particular, w can be reconstructed from a knowledge of F_1, F_2, F_3 ; and so no two different good looks produce the same triple (F_1, F_2, F_3) . We deduce that there are at least k distinct triples of perfect matchings, and the theorem follows. This proves 2.1. ■

Let C be a cycle of G with even length, and let $w_0 : E(C) \rightarrow \{1, -1\}$, such that the edges of C are mapped alternately to 1 and to -1 . We call w_0 a *bracelet* on C . Any such map w_0 that arises in this way from some cycle C is called a *bracelet* of G , and C is its *supporting cycle*. Define $w : E(G) \rightarrow \{-1, 0, 1\}$ by $w(e) = w_0(e)$ if $e \in E(C)$, and $w(e) = 0$ otherwise. Then w is a look, and we call w the *look* of the bracelet w_0 . Any function $w : E(G) \rightarrow \{-1, 0, 1\}$ that arises in this way from some bracelet w_0 we call a *bracelet look* of G . (It is convenient for us to distinguish between a bracelet and its look, especially when we come to the later parts of the proof, because sometimes the same bracelet will occur in several different graphs.)

Again, let G be cubic. A *jewel-box* for G is a set \mathcal{B} of bracelets of G , satisfying:

- every two members of \mathcal{B} have disjoint supporting cycles
- for every subset $W \subseteq \mathcal{B}$, the sum of the looks of the members of W is a good look.

We define $\beta(G)$ to be the cardinality of the largest jewel-box in G . Our main result is:

2.2 *For every planar cubic graph G with no cutedge, $\beta(G) \geq |V(G)|/218659584$.*

This implies that G has at least $2^{|V(G)|/218659584}$ good looks, and hence at least $2^{|V(G)|/655978752}$ perfect matchings, by 2.1; so 1.1 follows from 2.2.

3 The cyclically 4-edge-connected case

Let us say a cubic graph G is *cyclically 4-edge-connected* or C_4C if it is 3-connected (and hence is simple and has at least four vertices) and for every set $X \subseteq V(G)$ with $|X|, |V(G) \setminus X| > 1$ there are

at least four edges between X and $V(G) \setminus X$. Our goal in this section is to prove 2.2 (with a better constant) for C4C planar cubic graphs. We shall show the following:

3.1 For every C4C planar cubic graph G , $\beta(G) \geq |V(G)|/30976$.

We first need a lemma. (To clarify – the *degree* of a vertex is the number of edges incident with it; a *cycle* has no repeated vertices; and the *length* of a path or cycle is the number of edges in it.)

3.2 Let G be a simple planar graph, let $A \subseteq V(G)$ be stable, let $d > 0$ be an integer, and let each member of A have degree at most d in G . Then there exist $X \subseteq A$ and $Y \subseteq V(G) \setminus A$, such that

- $|X| \geq |A|/(64d + 8)$
- each member of X is adjacent to at most two members of Y
- every path in G of length at most three between two members of X has a vertex in Y .

Proof. Let Y be the set of vertices in $V(G) \setminus A$ with at least ten neighbours in A . Let A_1 be the set of vertices in A with at most two neighbours in Y , and let $A_2 = A \setminus A_1$.

$$(1) |A_1| \geq \frac{1}{2}|A|.$$

For we may assume that $A_2 \neq \emptyset$, and therefore $Y \neq \emptyset$. Consequently $|A \cup Y| \geq 11 \geq 3$. Let H_1 be the bipartite subgraph of G with $V(H_1) = A \cup Y$ and edge set all edges of G between A and Y . Since H_1 is planar, simple, and bipartite, and has at least three vertices, it follows (from Euler's formula; this is elementary and well known) that $|E(H_1)| \leq 2|V(H_1)| - 4$. But $|E(H_1)| \geq 10|Y|$, since every vertex in Y has at least 10 neighbours in A , and so $10|Y| \leq 2(|A| + |Y|) - 4$. Consequently $|Y| \leq \frac{1}{4}|A|$. Now let H_2 be the subgraph of H_1 induced on $A_2 \cup Y$. Then $|V(H_2)| \geq 4$ (since $A_2 \neq \emptyset$), and therefore $|E(H_2)| \leq 2(|A_2| + |Y|) - 4$ as before; but $|E(H_2)| \geq 3|A_2|$ since every member of A_2 has at least three neighbours in Y , and so $3|A_2| \leq 2(|A_2| + |Y|) - 4$, and in particular, $|A_2| \leq 2|Y|$. Since $|Y| \leq \frac{1}{4}|A|$ as we already saw, it follows that $|A_2| \leq \frac{1}{2}|A|$, and therefore $|A_1| \geq \frac{1}{2}|A|$. This proves (1).

Let H_3 be the graph with vertex set A_1 , in which distinct u, v are adjacent if there is a path in $G \setminus Y$ between u, v of length two. (Since A is stable, it follows that no internal vertex of such a path is in A .) Since each vertex of A has degree at most d in G , and each vertex of $V(G) \setminus (A \cup Y)$ has at most nine neighbours in A , it follows that for each $v \in A$ there are at most $8d$ paths in $G \setminus Y$ of length two with one end v , and hence v has degree at most $8d$ in H_3 . Consequently H_3 is colourable with $8d + 1$ colours; and therefore there is a subset A_3 of A_1 such that A_3 is stable in H_3 and $|A_3| \geq |A_1|/(8d + 1)$. But by (1), $|A_1| \geq \frac{1}{2}|A|$, and so $|A_3| \geq |A|/(16d + 2)$. Let H_4 be the graph with vertex set A_3 , in which distinct $u, v \in A_3$ are adjacent if there is a path in G between u, v of length 3 with no vertex in Y . Since no two members of A_3 have a common neighbour in $G \setminus Y$, this graph is a subgraph of the graph obtained from $G \setminus Y$ by contracting every edge with an end in A_3 . In particular, H_4 is planar, and since it is simple, it is 4-colourable, and so there is a subset $A_4 \subseteq A_3$ with $|A_4| \geq |A_3|/4$ that is stable in H_4 . Consequently, in G every path between two members of A_4 has a vertex in Y . But

$$|A_4| \geq \frac{1}{4}|A_3| \geq |A|/(64d + 8),$$

and so setting $X = A_4$ satisfies the theorem. This proves 3.2. ■

Now we prove the main result of this section.

Proof of 3.1. Let G be a C4C planar cubic graph with n vertices. Take a drawing of G in a 2-sphere Σ . Let us say a *domino* of G is a closed disc $\Delta \subseteq \Sigma$ with boundary a cycle of G of even length, such that Δ includes at most two regions of G , either exactly one region of G of even length, or two regions of odd length.

(1) *There exist at least $n/32$ dominos of G , pairwise disjoint, and each with boundary of length at most 15.*

For let G have R regions; thus by Euler's formula, $R = n/2 + 2$. Since the dual graph G^* of G is a 4-connected planar triangulation, Whitney's theorem [5] implies that the dual graph is Hamiltonian, and therefore we can number the regions of G as r_1, \dots, r_R , where for $1 \leq i \leq R$, the boundaries of r_i, r_{i+1} share an edge (reading subscripts modulo R). We choose the numbering of the regions such that r_R is the region of greatest length (the "length" of a region is the number of edges incident with it). Let $k = \lfloor \frac{1}{2}R \rfloor$. Then the average length of r_1, \dots, r_{2k} is at most that of r_1, \dots, r_R , and consequently less than six. Now for $1 \leq i \leq k$, the closure of one of $r_{2i-1}, r_{2i}, r_{2i-1} \cup r_{2i}$ is a domino Δ_i say, and its length is at most the sum of the lengths of r_{2i-1} and r_{2i} minus two. Thus the average length of $\Delta_1, \dots, \Delta_k$ is less than 10. We claim that at least half of them have length at most 15. For let a of them have length at most 15, and b length at least 16, and let L be the sum of all their lengths. Then $10(a+b) > L$; but since they all have length at least four, $L \geq 4a + 16b$. Consequently, $10(a+b) > 4a + 16b$, and so $a > b$. Thus at least half of $\Delta_1, \dots, \Delta_k$ have length at most 15. If we say that two of these dominos are adjacent when their boundaries share an edge, this defines a loopless planar graph, which is therefore 4-colourable; and consequently we can choose a quarter of our $\frac{1}{2}k$ dominos pairwise disjoint. Since $k \geq \frac{1}{2}(R-1) \geq \frac{1}{4}n$, this proves (1).

Let A be a set of disjoint dominos as in (1), with $|A| \geq n/32$. Let R be the set of all regions of G not included in any member of A . Let H be the graph with $V(H) = A \cup R$, in which $\Delta \in A$ and $r \in R$ are adjacent if the boundaries of Δ and r share an edge, and distinct $r_1, r_2 \in R$ are adjacent if their boundaries share an edge. Thus H is simple, and planar. By 3.2 with $d = 15$, there exist $X \subseteq A$ and $Y \subseteq R$, such that

- $|X| \geq |A|/968 \geq n/30976$
- each member of X is adjacent to at most two members of Y
- every path in H of length at most three between two members of X has a vertex in Y .

Let $X = \{\Delta_1, \dots, \Delta_k\}$ say, where $k \geq n/30976$. For $1 \leq i \leq k$, let C_i be the cycle that forms the boundary of Δ_i . There are two bracelets on C_i , and we choose one, w_i say, as follows. There are at most two members of Y adjacent to Δ_i in H . If there is at most one, let w_i be an arbitrary bracelet on C_i . Suppose there are exactly two, say r_1, r_2 . Thus the boundaries of r_1, r_2 both share at least one edge with the boundary of Δ_i . If possible, choose a bracelet w_i on C_i such that for some edge e of G , and for some $j \in \{1, 2\}$, $w_i(e) = 1$, and e is the unique edge in common between the boundaries of Δ_i and r_j . If this is not possible, and so both r_1, r_2 are incident with two edges that belong to the boundary of Δ_i , let w_i be an arbitrary bracelet on C_i .

We claim that the set $\{w_1, \dots, w_k\}$ is a jewel-box. Let $W \subseteq \{1, \dots, k\}$, and let w be the sum of the looks of all w_i ($i \in W$); we need to show that w is a good look. It is certainly a look; let us check that it is good, that is, $w(D) \neq 1 - |D|$ for every cut D of G . Suppose not; and choose a cut D of G with $|D|$ minimal such that $w(D) = 1 - |D|$. Consequently $w(f) = 0$ for some edge $f \in D$, and $w(e) = -1$ for all other edges of D . If some proper subset D_1 of D is also a cut, then so is $D \setminus D_1$, and so we may assume that $f \in D_1$; but then $w(D_1) = 1 - |D_1|$ contrary to the minimality of D . Thus no proper subset of D is a cut, and so D is a bond of G (that is, a minimal nonempty cut) and in particular, there is a cycle C of the dual graph G^* of G with $E(C) = D$ (we identify $E(G^*)$ with $E(G)$ in the natural way). We recall that R is the set of regions of G not included in any member of A . Let S be the set of all other regions of G ; thus, R, S form a partition of $V(G^*)$. For $i = -1, 0, 1$, let F_i be the set of edges e of G with $w(e) = i$; thus F_{-1}, F_0, F_1 form a partition of $E(G) = E(G^*)$. We have seen that every edge of C belongs to F_{-1} except for one in F_0 .

(2) *If $s \in V(C) \cap S$, and r_1, r_2 are its neighbours in C , then not both $r_1, r_2 \in Y$.*

For let e_i be the edge $r_i s$ of G^* . Suppose that $r_1, r_2 \in Y$ and hence $r_1, r_2 \in R$. Since $s \in S$, there is a unique $\Delta \in A$ such that s is included in Δ . Since $e_1, e_2 \in E(C)$, at least one of them belongs to F_{-1} ; and so $\Delta \in X$, say $\Delta = \Delta_j$ where $1 \leq j \leq k$, and $j \in W$. Consequently e_1, e_2 both belong to the supporting cycle of w_j , and so $w_j(e_1) = w_j(e_2) = -1$. From the choice of w_j it follows that r_1, r_2 are both incident with two edges of the boundary of Δ_j in G ; and since G is 3-connected, it follows that Δ_j is not the closure of a region of G , and so Δ_j is the closure of the union of s and some other region s' of G , and their boundaries share an edge $v_1 v_2$ of G that is drawn in the interior of Δ_j in the drawing of G . In particular, s has odd length. Moreover, for $i = 1, 2$, the boundary of r_i meets that of s and that of s' ; and since G is C4C it follows that r_i is incident with exactly one of v_1, v_2 in the drawing of G . Thus we may assume that r_1 is incident with v_1 and r_2 with v_2 . Consequently for $i = 1, 2$, e_i is incident with v_i in G ; and yet $w_j(e_1) = w_j(e_2) = -1$, contradicting that s has odd length. This proves (2).

(3) *Let e be the unique edge of C in F_0 . Then in G^* , either both ends of e are in R or both are in S .*

For suppose that $e = rs$ in G^* , where $r \in R$ and $s \in S$. Let r' be the second neighbour of s in C . Since $r's \in F_{-1}$, it follows that $r' \in R$, and s is included in some Δ_i where $i \in W$. But then every edge of the boundary of Δ_i belongs to $F_1 \cup F_{-1}$, contradicting that $e \in F_0$. This proves (3).

(4) *Let $s, s' \in S$ both be incident with some edge e of G , so $e = ss'$ in G^* . If both $s, s' \in V(C)$ then they are adjacent in C .*

Let P_1, P_2 be the two paths of C between s, s' . In one of them, say P_1 , every edge belongs to F_{-1} . Since in the drawing of G , the edge e is drawn in the interior of a member of A , it follows that $e \in F_0$; and so the cycle C_1 of G^* made by adding e to P_1 satisfies $w(E(C_1)) = 1 - |E(C_1)|$. By the minimality of $|D|$, it follows that the length of C_1 is at least that of C , and so P_2 has only one edge, and hence s, s' are adjacent in C . This proves (4).

Now every edge of F_{-1} has (in G^*) an end in R and an end in S , and consequently every edge of C except one is between R and S . Since G is 3-connected, it follows that G^* is simple, and so C has length at least three. By (3), C has odd length, say $2t + 1$. Suppose that two consecutive vertices of C belong to S ; then we can number the vertices of C in order as

$$s_0-r_1-s_1-r_2-s_2-\cdots-r_t-s_t-s_0,$$

where $r_1, \dots, r_t \in R$ and $s_0, s_1, \dots, s_t \in S$. For $1 \leq i \leq t$ there exists $\Delta \in A$ with $s_i \subseteq \Delta$, and since the edge $r_i s_i$ of G^* belongs to F_{-1} , it follows that $\Delta = \Delta_j$ for some $j \in W$. Moreover, by (4) all these dominos are distinct, and so we may assume that $s_i \subseteq \Delta_i$ and $i \in W$ for $1 \leq i \leq t$. Suppose that $t \geq 2$. Then for $i = 1, 2$, the vertex r_i of H is adjacent in H to the vertices Δ_i, Δ_{i+1} of H , and so $r_i \in Y$ from the choice of X, Y ; but this contradicts (2). It follows that $t = 1$, and so r_1 is incident with an edge of the boundary of s_0 , and an edge of the boundary of s_1 . Since G is C4C, it follows that these two edges have a common end in G , and therefore do not both belong to F_{-1} , a contradiction. This proves that no two consecutive vertices of C belong to S .

Consequently there are two consecutive vertices of C that belong to R , and we can number the vertices of C in order as

$$r_0-s_1-r_1-s_2-\cdots-s_t-r_t-r_0,$$

where $r_0, r_1, \dots, r_t \in R$ and $s_1, \dots, s_t \in S$. As before we may assume that $s_i \subseteq \Delta_i$ and $i \in W$ for $1 \leq i \leq t$. Suppose that $t \geq 3$; then for $i = 1, 2$, the vertex r_i of H is adjacent in H to the vertices Δ_i, Δ_{i+1} of H , and so $r_i \in Y$ from the choice of X, Y ; but this contradicts (2). So $t \leq 2$. Suppose that $t = 2$; then since there is a path of H with vertices Δ_1, r_2, Δ_2 , it follows that $r_2 \in Y$ from the choice of X, Y ; and since also there is a path of H with vertices $\Delta_2, r_0, r_1, \Delta_1$ in order, it follows from the choice of X, Y that at least one of $r_0, r_1 \in Y$, contrary to (2). Thus $t = 1$. But then r_0, r_1, s_1 are regions that pairwise are incident with a common edge, and since G is C4C, it follows that for some vertex v of G , these three edges are all incident with v , and yet two of the edges belong to F_{-1} , a contradiction. This proves that $\{w_1, \dots, w_k\}$ is a jewel-box, and therefore completes the proof of 3.1. ■

4 The 3-connected case.

In this section we extend 3.1 (changing the constant 30976 to something larger) to planar cubic graphs that are 3-connected but not necessarily C4C. It would seem natural to do this by induction on the size of the graph, but we were not able to do so. Instead, we need to work with a set of three-edge cuts that decompose the graph into C4C pieces, and we begin by describing this decomposition.

Let G be a graph. A *cut-decomposition* of G is a pair (T, ϕ) such that:

- T is a tree with $E(T) \neq \emptyset$,
- $\phi : V(G) \rightarrow V(T)$ is a map, and
- for each $t \in V(T)$ with degree one or two in T , there exists $v \in V(G)$ with $\phi(v) = t$.

If $t \in V(T)$, $\phi^{-1}(t)$ denotes the set of $v \in V(G)$ with $\phi(v) = t$. Similarly, if $Y \subseteq V(T)$, we denote $\{v \in V(G) : \phi(v) \in Y\}$ by $\phi^{-1}(Y)$; and if S is a subgraph of T we write $\phi^{-1}(S)$ for $\phi^{-1}(V(S))$. For

each edge e of T , let T_1, T_2 be the two components of $T \setminus e$, and for $i = 1, 2$ let $X_i = \phi^{-1}(T_i)$. Thus X_1, X_2 is a partition of $V(G)$, and therefore $\delta(X_1) = \delta(X_2)$ is a cut; we denote this cut by $\phi^{-1}(e)$. If $|\phi^{-1}(e)| = k$ for each $e \in E(T)$ we call (T, ϕ) a k -cut-decomposition of G . We will only be concerned with 2- and 3-cut-decompositions of cubic graphs.

Let (T, ϕ) be a 3-cut-decomposition of G , and let T_0 be a subtree of T . Let T_1, \dots, T_s be the components of $T \setminus V(T_0)$, and for $1 \leq i \leq s$ let e_i be the unique edge of T with an end in $V(T_0)$ and an end in $V(T_i)$. For $0 \leq i \leq s$, let $X_i = \phi^{-1}(T_i)$. Thus X_0, X_1, \dots, X_s are pairwise disjoint subsets of $V(G)$ with union $V(G)$. Let G' be the graph obtained from G by, for $1 \leq i \leq s$, deleting all edges of $G|X_i$ and identifying all the vertices in X_i . (If $G|X_i$ is connected, this is the same as contracting all edges of $G|X_i$.) Thus the graph G' has $|X_0| + s$ vertices, and all the vertices of G' not in X_0 have degree 3. We call G' the 3-hub of G at T_0 (with respect to (T, ϕ)). If $t_0 \in V(T)$, by the “3-hub of G at t_0 ” we mean the 3-hub of G at T_0 , where T_0 is the subtree of T with vertex set $\{t_0\}$. If \mathcal{C} is a class of graphs and (T, ϕ) is a 3-cut-decomposition of G , and for each $t \in V(T)$ the 3-hub of G at t belongs to \mathcal{C} , we say that (T, ϕ) is a 3-cut-decomposition of G over \mathcal{C} .

Let \mathcal{C}_4 be the class of C4C planar cubic graphs. We begin with:

4.1 *Every 3-connected planar cubic graph G that is not C4C admits a 3-cut-decomposition over \mathcal{C}_4 .*

Proof. We proceed by induction on $|V(G)|$. Since G is not C4C, there is a partition X_1, X_2 of $V(G)$ such that $|\delta(X_1)| = 3$ and $|X_1|, |X_2| > 1$. Since G is 3-connected, it follows that $G|X_1, G|X_2$ are connected. Let G_1 be obtained from G by contracting the edges of $G|X_2$, and let x_1 be the vertex of G_1 formed by identifying the vertices of X_2 . Define G_2, x_2 similarly. Then G_1, G_2 are 3-connected, planar, cubic, and have fewer vertices than G . For $i = 1, 2$, if G_i is C4C let T_i be a tree with one vertex t_i and define $\phi_i(v) = t_i$ for each $v \in V(G_i)$; if G_i is not C4C, let (T_i, ϕ_i) be a 3-cut-decomposition of G_i over \mathcal{C}_4 (this exists from the inductive hypothesis), and let $t_i = \phi_i(x_i)$. Note that for $i = 1, 2$, if t_i has degree zero in T_i , then $|\phi_i^{-1}(t_i)| \geq 4$ since $\phi_i^{-1}(t_i) = V(G_i)$ and G_i is 3-connected; while if t_i has degree one in T_i then $|\phi_i^{-1}(t_i)| \geq 3$ since the 3-hub of G_i at t_i with respect to (T_i, ϕ_i) is C4C. Thus in either case $|\phi_i^{-1}(t_i)| \geq 3$. Let T be the tree obtained from the disjoint union of T_1 and T_2 by making t_1, t_2 adjacent; and for $v \in V(G)$, define $\phi(v) = \phi_i(v)$ where $v \in X_i$. Consequently, for $i = 1, 2$, if t_i has degree at most two in T then $|\phi^{-1}(t_i)| \geq 2$. It is easy to check that (T, ϕ) is a 3-cut-decomposition of G over \mathcal{C}_4 . This proves 4.1. \blacksquare

4.2 *Let (T, ϕ) be a 3-cut-decomposition of a 3-connected cubic graph G , let T_1, \dots, T_k be pairwise vertex-disjoint subtrees of T , and for $1 \leq i \leq k$ let H_i be the 3-hub of G at T_i . For $1 \leq i \leq k$ let \mathcal{B}_i be a set of bracelets of G , such that*

- \mathcal{B}_i is a jewel-box of H_i , and
- if C is the supporting cycle of a member of \mathcal{B}_i , then $V(C) \subseteq \phi^{-1}(T_i)$.

Then $\mathcal{B}_1 \cup \dots \cup \mathcal{B}_k$ is a jewel-box in G .

Proof. Let $w, w' \in \mathcal{B}_1 \cup \dots \cup \mathcal{B}_k$ be distinct, on cycles C, C' of G respectively. We must show that C, C' are vertex-disjoint. Choose $i, j \in \{1, \dots, k\}$ such that $w \in \mathcal{B}_i$ and $w' \in \mathcal{B}_j$. Thus $V(C) \subseteq \phi^{-1}(T_i)$ and $V(C') \subseteq \phi^{-1}(T_j)$. If $i \neq j$ then T_i, T_j are disjoint and therefore C, C' are disjoint; while if $i = j$ then w, w' both belong to the same jewel-box in H_i , and therefore have disjoint supporting cycles.

Second, let $W \subseteq \mathcal{B}_1 \cup \dots \cup \mathcal{B}_k$, and let w be the sum of the looks of all members of W ; and for $i = -1, 0, 1$ let F_i be the set of edges e of G such that $w(e) = i$. We must show that w is a good look. Suppose not; then there is a cut D of G and an edge $f \in D$ such that $f \in F_0$ and $D \setminus \{f\} \subseteq F_{-1}$. Choose such a cut D with $|D|$ minimum. Then as in the proof of 3.1, D is a bond of G , that is, $D = \delta(X)$ for some $X \subseteq V(G)$ with $G|X, G|(V(G) \setminus X)$ both connected.

(1) Let $e \in E(T)$, let S_1, S_2 be the two components of $T \setminus e$, and for $i = 1, 2$ let $Y_i = \phi^{-1}(V(S_i))$. Then one of $X \cap Y_1, X \cap Y_2, Y_1 \setminus X, Y_2 \setminus X$ is empty.

Suppose all four of these sets are nonempty. Since $\delta(Y_1) \subseteq F_0$, and $\delta(X) \setminus \{f\} \subseteq F_{-1}$, it follows that $\delta(Y_1) \cap \delta(X) \subseteq \{f\}$. There are two cases depending whether $f \in \delta(Y_1)$ or not. Suppose first that $f \notin \delta(Y_1)$. Then $\delta(Y_1) \cap \delta(X) = \emptyset$, and from the symmetry between Y_1 and Y_2 we may assume that both ends of f belong to Y_2 . Since $G|X$ is connected and $X \cap Y_1, X \cap Y_2$ are both nonempty, some edge of $\delta(Y_1)$ has both ends in X , and similarly some edge of $\delta(Y_1)$ has both in $V(G) \setminus X$. Since $|\delta(Y_1)| = 3$ and every edge of $\delta(Y_1)$ either has both ends in X or both in $V(G) \setminus X$, we may assume (replacing X by its complement if necessary) that exactly one edge f' of $\delta(Y_1)$ has both ends in X . But then $f' \in F_0$, and f' belongs to the cut $\delta(X \cap Y_1)$, and every other edge of this cut belongs to $\delta(X)$ and therefore to F_{-1} . From the minimality of $|D|$, it follows that $|\delta(X \cap Y_1)| \geq |\delta(X)|$, and so there is at most one edge of G between $X \cap Y_2$ and $Y_2 \setminus X$. But then $|\delta(X \cap Y_2)| \leq 2$, a contradiction since G is 3-connected.

This proves that $f \in \delta(Y_1)$. From the symmetry between Y_1, Y_2 we may assume that f is between $X \cap Y_2$ and $Y_1 \setminus X$. Since $G|X$ is connected, there is at least one edge between $X \cap Y_1$ and $X \cap Y_2$. Suppose that there is only one. Since G is 3-connected, there are at least two edges between $X \cap Y_1$ and $Y_1 \setminus X$, and an edge between $X \cap Y_2$ and $Y_2 \setminus X$; and so $\delta(X \cap Y_1)$ contradicts the minimality of $|D|$. Thus there are at least two edges between $X \cap Y_1$ and $X \cap Y_2$. Since $|\delta(Y_1)| = 3$, there are no edges between $Y_1 \setminus X$ and $Y_2 \setminus X$, contradicting that $G|(V(G) \setminus X)$ is connected. This proves (1).

Since G is 3-connected and therefore $|D| \geq 3$, we may choose $e \in D$ with $e \neq f$. Thus $\phi(e) = -1$, and so e is an edge of one of C_1, \dots, C_k , say C_1 . Let $Y_0 = \phi^{-1}(T_1)$. Let $e = xy$; then since $V(C_1) \subseteq Y_0$ it follows that $x, y \in Y_0$, and since $e \in \delta(X)$, we deduce that $X \cap Y_0, Y_0 \setminus X$ are both nonempty. Let S_1, \dots, S_s be the components of $T \setminus V(T_1)$, and for $1 \leq i \leq s$ let e_i be the unique edge of T with an end in $V(T_1)$ and an end in $V(S_i)$. For $1 \leq i \leq s$, let $Y_i = \phi^{-1}(S_i)$. Thus Y_0, Y_1, \dots, Y_s are pairwise disjoint subsets of $V(G)$ with union $V(G)$. The 3-hub H_1 of G at T_1 is therefore obtained by contracting all edges of $G|Y_i$ for $1 \leq i \leq s$. Since D is not a cut of H_1 (because \mathcal{B}_1 is a jewel-box of H_1) it follows that for some $i \in \{1, \dots, s\}$, both $X \cap Y_i, Y_i \setminus X$ are nonempty. But since both $Y_0 \cap X, Y_0 \setminus X$ are nonempty, this contradicts (1) applied to the edge e_i . This proves 4.2. \blacksquare

Here is a convenient lemma:

4.3 Let G be a cubic graph, and let F_1, F_2, F_3 be pairwise disjoint perfect matchings of G . Let C_1, \dots, C_k be the cycles of G with edge-set included in $F_1 \cup F_2$, and for $1 \leq i \leq k$ let w_i be the bracelet on C_i such that $w_i(e) = -1$ if $e \in F_1 \cap E(C_i)$ and $w_i(e) = 1$ if $e \in F_2 \cap E(C_i)$. Then $\{w_1, \dots, w_k\}$ is a jewel-box in G , and so $\beta(G) \geq k$. Moreover, if $e \in E(G)$, then $\beta(G) > 0$, and there is a bracelet w such that e is not in the supporting cycle of w and $\{w\}$ is a jewel-box.

Proof. We must show that for every subset $X \subseteq F_1 \cup F_2$ that is a union of edge-sets of cycles, if we delete from G every edge in $F_1 \cap X$, and add a new edge parallel to every edge in $F_2 \cap X$, then

the graph G' that we construct has no cutedge. But F_2, F_3 are disjoint perfect matchings of G' , and the remaining edges of G' form a third, and so G' is 3-edge-colourable and therefore has no cutedge. Consequently $\{w_1, \dots, w_k\}$ is a jewel-box, and so $\beta(G) \geq k$. For the final claim, let $e \in E(G)$; from the symmetry between F_1, F_2, F_3 we may assume that $e \in F_3$, and since $F_1, F_2 \neq \emptyset$ there is a cycle within $F_1 \cup F_2$, so the final claim follows from the first. This proves 4.3. \blacksquare

A 3-edge-colouring of G is a map $\kappa : E(G) \rightarrow \{1, 2, 3\}$ such that $\kappa(e) \neq \kappa(f)$ for every two distinct edges $e, f \in E(G)$ with a common end; and two 3-edge-colourings κ, κ' are *equivalent* if there is a permutation π of $\{1, 2, 3\}$ such that $\kappa'(e) = \pi(\kappa(e))$ for each $e \in E(G)$. If κ is a 3-edge-colouring of G and $X \subseteq E(G)$, we denote the restriction of κ to X by $\kappa|X$; and if $i \in \{1, 2, 3\}$, we denote the set of $e \in E(G)$ with $\kappa(e) = i$ by $\kappa^{-1}(i)$. A cubic graph G is *uniquely 3-edge-colourable* (U3C) if there is a unique set $\{F_1, F_2, F_3\}$ of perfect matchings of G with union $E(G)$; that is, if it is 3-edge-colourable and all its 3-edge-colourings are equivalent.

Finding a 3-edge-colouring of a cubic graph G equipped with a 3-cut-decomposition is just a matter of finding a 3-edge-colouring of the 3-hub of G at each vertex of the tree. In particular, if (T, ϕ) is a 3-cut-decomposition of a cubic graph G , and H is the 3-hub of G at some $t \in V(T)$, and κ is a 3-edge-colouring of G , then $\kappa|E(H)$ is a 3-edge-colouring of H ; while if G admits a 3-edge-colouring, then every 3-edge-colouring of H can be extended to a 3-edge-colouring of G .

If (T, ϕ) is a 3-cut-decomposition of a cubic graph G with $|V(T)| \geq 3$, and $e = t_1 t_2$ is an edge of T , let T' be the tree obtained from T by contracting e (forming a vertex t' say), and for $v \in V(G)$, define $\phi'(v) = \phi(v)$ if $\phi(v) \neq t_1, t_2$, and $\phi'(v) = t'$ if $\phi(v) = t_1$ or t_2 . Then (T', ϕ') is also a 3-cut-decomposition of G , and we say it is obtained from (T, ϕ) by *contracting* e . Note that if the 3-hub of G at one of t_1, t_2 (with respect to (T, ϕ)) is not U3C, then the 3-hub of G at t' (with respect to (T', ϕ')) is not U3C, since if the second 3-hub is 3-edge-colourable then every 3-edge-colouring of the first 3-hub extends to a 3-edge-colouring of the second.

4.4 *Let (T, ϕ) be a 3-cut-decomposition of a 3-connected 3-edge-colourable cubic graph G . Then there are at most $6\beta(G)$ vertices $t \in V(T)$ such that the 3-hub of G at t is not U3C.*

Proof. Let there be n_1 vertices $t \in V(T)$ such that the 3-hub of G at t is not U3C; we need to show that $n_1 \leq 6\beta(G)$. Since $\beta(G) \geq 1$ by 4.3, we may assume that $n_1 \geq 7$. By contracting edges of T appropriately, we may therefore assume that for every vertex $t \in V(T)$, the 3-hub of G at t is not U3C. Consequently we may choose 3-edge-colourings κ, κ' of G , such that for each $t \in V(T)$ with 3-hub H say, $\kappa|E(H)$ and $\kappa'|E(H)$ are not equivalent. By permuting the elements of $\{1, 2, 3\}$ in one of κ, κ' , we may assume that $\kappa|\phi^{-1}(e) = \kappa'|\phi^{-1}(e)$ for at least one-sixth of all edges $e \in E(T)$. By contracting all other edges of T , we obtain a 3-cut-decomposition (S, ψ) of G , such that

- $|E(S)| \geq (n_1 - 1)/6$,
- for each $t \in V(S)$ with 3-hub H say, $\kappa|E(H) \neq \kappa'|E(H)$,
- $\kappa|\psi^{-1}(e) = \kappa'|\psi^{-1}(e)$ for all $e \in E(S)$.

Let $V(S) = \{s_1, \dots, s_m\}$ say, where $m \geq (n_1 - 1)/6 + 1$, and for $1 \leq i \leq m$ let H_i be the 3-hub of G at s_i with respect to (S, ψ) . Fix i with $1 \leq i \leq m$. For $j = 1, 2, 3$, let F_j be the set of edges e of H_i with $\kappa(e) = j$, and let F'_j be the set of edges e of H with $\kappa'(e) = j$. Since $\kappa|E(H_i) \neq \kappa'|E(H_i)$, there exists $j \in \{1, 2, 3\}$ such that $F_j \neq F'_j$. Choose a cycle C_i with $E(C_i) \subseteq (F_j \setminus F'_j) \cup (F'_j \setminus F_j)$,

and let w_i be some bracelet on C_i . Note that $\psi(v) = s_i$ for every $v \in V(C_i)$, since every edge of C belongs to just one of F_j, F'_j , and yet $\kappa|\psi^{-1}(e) = \kappa'|\psi^{-1}(e)$ for every edge e of S . We claim that the look of w_i is a good look in H_i . To see this, let J be obtained from H_i by deleting every edge e with $w_i(e) = -1$, and adding a new edge parallel to every edge e with $w(e) = 1$; we must show that J has no cutedge. Since every edge of C_i belongs to exactly one of F_j, F'_j , we may assume that the edges e of C_i with $w_i(e) = -1$ belong to F_j , and that $j = 1$ say. But then F_2, F_3 are disjoint perfect matchings of J , and the remaining edges of J form a third perfect matching, and so J is 3-edge-colourable, and therefore has no cutedge. This proves that the look of w_i is a good look in H_i , and so $\{w_i\}$ is a jewel-box in H_i .

These jewel-boxes satisfy the hypotheses of 4.2, and so by 4.2, it follows that the set $\{w_1, \dots, w_m\}$ is a jewel-box in G . Since

$$\beta(G) \geq m \geq (n_1 - 1)/6 + 1 \geq n_1/6,$$

this proves 4.4. ■

We need the following theorem of Fowler [1]:

4.5 *The only planar cubic graph that is both C_4C and U_3C is the graph K_4 .*

Let u_1, u_2, u_3 be pairwise adjacent vertices of a cubic graph G' , and for $1 \leq i \leq 3$ let u_i have a neighbour $v_i \notin \{u_1, u_2, u_3\}$. Let G be obtained from G' by contracting the three edges u_1u_2, u_2u_3, u_3u_1 , forming a vertex v say. We say that G' is obtained from G by “replacing v by a triangle”. Let w be a bracelet in G , with supporting cycle C . If $v \notin V(C)$ then w is a bracelet in G' . If $v \in V(C)$, and say the edges vv_1, vv_2 belong to $E(C)$, let C' be the cycle of G' consisting of the path $C \setminus v$ and the path $v_1u_1u_3u_2v_2$, and let w' be the bracelet on C' that equals w on the edges of $C \setminus v$. In this case we call w' the “natural rerouting of w ”.

4.6 *Let $G, v, G', u_1, u_2, u_3, v_1, v_2, v_3$ be as above, and let $\mathcal{B} = \{w_1, \dots, w_k\}$ be a jewel-box in G . For $1 \leq i \leq k$, if v is in the supporting cycle of w_i let w'_i be the natural rerouting of w_i , and otherwise let $w'_i = w_i$. Let $\mathcal{B}' = \{w'_1, \dots, w'_k\}$. Then \mathcal{B}' is a jewel-box in G' .*

Proof. Certainly the members of \mathcal{B}' have disjoint supporting cycles, because v belongs to the supporting cycle of at most one member of \mathcal{B} . Let $W' \subseteq \mathcal{B}'$, and let w' be the sum of the looks of members of W' ; we must show that w' is a good look in G' . Suppose not, and let D' be a cut of G' with $w'(D') = 1 - |D'|$. Choose $X' \subseteq V(G')$ with $D' = \delta_{G'}(X')$. By replacing X' by its complement if necessary, we may assume that at most one of $u_1, u_2, u_3 \in X'$. For $-1 \leq i \leq 1$, let F_i be the set of edges $e \in E(G)$ with $w(e) = i$, and let F'_i be the set of edges $e \in E(G')$ with $w'(e) = i$. Thus every edge of D' belongs to F'_{-1} except for exactly one in F'_0 . Let W be the set of all w_i such that $w'_i \in W'$, and let w be the sum of the looks in G of the members of W . Thus w is a good look in G .

Suppose first that none of $u_1, u_2, u_3 \in X'$; then D' is also a cut of G , and $w(D') = w'(D') = 1 - |D'|$, contradicting that w is a good look in G . So we may assume that $u_1 \in X'$ and $u_2, u_3 \notin X'$. In particular, $u_1u_2, u_1u_3 \in D'$, and since $|D' \setminus F'_{-1}| = 1$, we may assume that $u_1u_2 \in F'_{-1}$. Consequently there exists some member of W' , say w'_1 , such that u_1u_2 is an edge of the supporting cycle of w'_1 , and $w'_1(u_1u_2) = -1$; and so $w_1 \in W$, and v belongs to the supporting cycle of w_1 . Now $u_1u_3 \notin F'_{-1}$ from the definition of w' , and yet no edge of D' is in F'_1 ; so $u_1u_3 \in F'_0$. It follows that $u_2u_3 \in F'_1$, since u_2u_3 is an edge of the supporting cycle of w'_1 ; and so $u_2v_2 \in F_0 \cap F'_0$, and $u_3v_3 \in F_{-1} \cap F'_{-1}$. Let $X = (X' \setminus \{u_1\}) \cup \{v\}$, and let D be the cut $\delta_G(X)$ of G ; then every edge of D belongs to F_{-1}

except for exactly one, namely $u_2v_2 \in F_0$, contradicting that w is a good look in G . This proves 4.6. ■

A *leaf* of a tree means a vertex with degree one. The *inner tree* of a tree T with at least three vertices is the tree obtained by deleting all leaves of T . An *twig* of a tree T is a leaf of its inner tree. Thus, for every twig t of T , t has at least two neighbours in T , and all of its neighbours are leaves of T except exactly one.

4.7 *Let G be a 3-connected 3-edge-colourable cubic graph, and let (T, ϕ) be a 3-cut-decomposition of G over \mathcal{C}_4 . Then T has at most $7\beta(G)$ twigs.*

Proof. Let P be the set of vertices t of T such that the 3-hub at t is not U3C.

(1) *Let t_0 be a twig and t_1, \dots, t_k all the leaves adjacent to t_0 . Let S be the subtree of T with vertex set $\{t_0, t_1, \dots, t_k\}$, and suppose that $V(S) \cap P = \emptyset$. Let H be the 3-hub of G at S . Then there is a cycle C of G , satisfying $\phi(v) \in V(S)$ for every vertex $v \in V(C)$, and a bracelet w on C , such that $\{w\}$ is a jewel-box of H .*

For $0 \leq i \leq k$ let H_i be the 3-hub of G at t_i . Since H_i is both C4C and U3C, it follows from 4.5 that H_i is isomorphic to K_4 for $0 \leq i, \dots, k$. In particular, since H_0 is isomorphic to K_4 , it follows that $1 \leq k \leq 3$. Let H_0 have vertex set $\{v_1, \dots, v_4\}$, where v_4 is the vertex formed by identifying all vertices v of G with $\phi(v) \notin V(S)$, and for $1 \leq i \leq k$ v_i is the vertex formed by identifying the members of $\phi^{-1}(t_i)$. Thus $\{v_{k+1}, \dots, v_3\} = \phi^{-1}(t_0)$. Consequently H is obtained from K_4 by replacing v_1, \dots, v_k by triangles. Since $k \geq 1$, it follows that H is obtained from a ‘‘prism’’ (the complement of a six-vertex cycle), say J , with one vertex called v_4 , by replacing one or two other vertices by triangles. In view of 4.6, it suffices to check that there is a cycle C of the prism J with $v_4 \notin V(C)$ such that $\{w\}$ is a jewel-box in J for some bracelet w on C . We leave this to the reader (use the cycle of length four). This proves (1).

Let there be m twigs in T . The at least $m - |P|$ of them satisfy the hypotheses of (1), and so by 4.2, $\beta(G) \geq m - |P|$. But $|P| \leq 6\beta(G)$ by 4.4, and so $m \leq 7\beta(G)$. This proves 4.7. ■

4.8 *Let G be a 3-connected 3-edge-colourable cubic graph, and let (T, ϕ) be a 3-cut-decomposition of G over \mathcal{C}_4 . Then the inner tree of T (if it exists) has at most $84\beta(G)$ vertices.*

Proof. Let P be the set of vertices t of T such that the 3-hub at t is not U3C. Let T' be the inner tree of T (we may assume that this exists). Let $t_1 \cdots t_4$ be a four-vertex path of T' such that t_1, \dots, t_4 all have degree two in T' ; and let S be the subtree of T induced on the union of $\{t_1, \dots, t_4\}$ and the set of leaves of T adjacent to one of t_1, \dots, t_4 . We call S a *limb* of T .

(1) *Let S be a limb with t_1, \dots, t_4 as above, and suppose that $P \cap V(S) = \emptyset$. Let H be the 3-hub of G at S . Then there is a cycle C of G with $V(C) \subseteq \phi_{-1}(S)$, and a bracelet w on C such that $\{w\}$ is a jewel-box of H .*

For since $P \cap V(S) = \emptyset$, it follows from 4.5 that the 3-hub of G at t is isomorphic to K_4 for every vertex $t \in V(S)$; and so t_1, \dots, t_4 each have degree two, three or four in T . It follows that the

3-hub of G at the subtree with vertex set $\{t_1, \dots, t_4\}$ can be constructed as follows: start with K_4 and call one of its vertices x ; choose another vertex of the K_4 and replace it by a triangle; choose a vertex of this triangle and replace it by a triangle; choose a vertex of the most recent triangle and replace it by a triangle; and choose a vertex of this latest triangle and call it y . Let us call this graph J . Then H can be obtained from J by replacing any of its vertices by triangles, except x, y . We must check that there is a cycle of H not containing x, y , and a bracelet w on H , such that $\{w\}$ is a jewel-box of H . By 4.6, it is enough to prove the claim for J rather for H . To check the claim for J , note that any cycle of length four in J not using x, y will do, so we can assume there is no such cycle, and then the possibilities for J are greatly restricted (there are only three). We leave the rest to the reader. This proves (1).

Let $|V(T')| = m$ (and we may assume that $m > 1$), and let T' have m_1 leaves, and m_2 vertices of degree two. Thus T' has $m - m_1 - m_2$ vertices of degree at least three. Since every tree has at least as many leaves as it does vertices of degree at least three, it follows that $m - m_1 - m_2 \leq m_1$, and so $m \leq 2m_1 + m_2$. A *branch* of a tree is a maximal subpath (with at least one edge) such that all its internal vertices have degree two in the tree; the number of branches of a tree is one less than the number of vertices of degree different from two. Thus T' has $m - m_2 - 1$ branches. Let its branches be B_1, \dots, B_k say, where $k = m - m_2 - 1$, and let B_i have $b_i + 2$ vertices for $1 \leq i \leq k$. Thus $b_1 + \dots + b_k = m_2$. Each B_i has b_i internal vertices, and so there are $\lfloor b_i/4 \rfloor \geq (b_i - 3)/4$ disjoint 4-vertex paths within the interior of B_i . Consequently there are at least

$$\sum_{i=1, \dots, k} (b_i - 3)/4 \geq m_2/4 - 3k/4$$

disjoint limbs in T , and at most $|P|$ of them contain members of P , so at least $m_2/4 - 3k/4 - |P|$ of them satisfy the hypotheses of (1). By 4.2, it follows that $m_2/4 - 3k/4 - |P| \leq \beta(G)$. Since $k \leq m - m_2$, it follows that $4m_2 - 3m - 4|P| \leq 4\beta(G)$. We already saw that $m \leq 2m_1 + m_2$ (that is, $m - 8m_1 \leq 4m_2 - 3m$), and therefore $m \leq 8m_1 + 4|P| + 4\beta(G)$. But $m_1 \leq 7\beta(G)$ by 4.7, and $|P| \leq 6\beta(G)$ by 4.4, and it follows that $m \leq 84\beta(G)$. This proves 4.8. \blacksquare

4.9 *Let H be a C4C planar cubic graph, let $X \subseteq V(H)$, and let G be obtained from H by replacing some vertices not in X by triangles. Then there is a jewel-box \mathcal{B} in G of cardinality at least $|V(G)|/92928 - |X|$, such that the supporting cycle of each member of \mathcal{B} contains no vertex in X .*

Proof. By 3.1, there is a jewel-box in H with cardinality at least $|V(H)|/30976 \geq |V(G)|/92928$. The supporting cycles of at most $|X|$ members of this jewel-box contain a vertex of X , and so there is a jewel-box in H with cardinality at least $|V(G)|/92928 - |X|$ such that none of the supporting cycles of its members contain members of X . By 4.6, the same holds in G . This proves 4.9. \blacksquare

4.10 *For every 3-connected planar cubic graph G , $\beta(G) \geq |V(G)|/16819968$.*

Proof. If G is C4C this follows from 3.1, so we assume that G is not C4C. By 4.1, there is a 3-cut-decomposition (T, ϕ) of G over \mathcal{C}_4 . Let the vertices of T be t_1, \dots, t_k . Let P be the set of vertices t of T such that the 3-hub of G at t is not U3C. Let L be the set of all leaves of T that do not belong to P , and let $S = T \setminus L$. If S is null then $|L| = 2$ and both members of L have

U3C 3-hubs, and so by 4.5 $|V(G)| \leq 6$ and the theorem holds. We may therefore assume that S is nonnull. Let $V(S) = \{t_1, \dots, t_s\}$ say, where $s \leq k$; and so $L = \{t_{s+1}, \dots, t_k\}$. For $1 \leq i \leq s$, let T_i be the subtree of T with vertex set consisting of t_i and all members of L adjacent to t_i . Thus T_1, \dots, T_s are pairwise disjoint subtrees of T , and every vertex of T belongs to exactly one of them. For $1 \leq i \leq s$, let H_i be the 3-hub of G at T_i . Then $V(H_i)$ is the disjoint union of two sets X_i, Y_i say, where $Y_i = \phi^{-1}(T_i)$. Now H_i is obtained from the 3-hub of G at t_i (which is C4C) by replacing some vertices not in X_i by triangles; and so by 4.9 there is a jewel-box \mathcal{B}_i in H_i of cardinality at least $|V(H_i)|/92928 - |X_i| \geq |Y_i|/92928 - |X_i|$, such that the supporting cycle of each member of \mathcal{B}_i contains no vertex in X_i . By 4.2, it follows that the union of these jewel-boxes is a jewel-box in G , and so

$$\sum_{1 \leq i \leq s} |Y_i|/92928 - |X_i| \leq \beta(G).$$

But $|Y_1| + \dots + |Y_s| = |V(G)|$, and $|X_1| + \dots + |X_s| = 2|E(S)| \leq 2|V(S)|$, so we deduce that $|V(G)|/92928 \leq 2|V(S)| + \beta(G)$. Now every vertex of S either belongs to the inner tree of T or to P , and G is 3-edge-colourable by the four-colour theorem; so by 4.8 and 4.4, $|V(S)| \leq 90\beta(G)$. Thus $|V(G)|/92928 \leq 181\beta(G)$, that is, $\beta(G) \geq |V(G)|/16819968$. This proves 4.10. \blacksquare

5 The non-3-connected case.

Now we extend 4.10 to all planar cubic graphs without cutedges. We begin with:

5.1 *Let G be a 3-edge-colourable 2-connected cubic graph. If (T, ϕ) is a 2-cut-decomposition of G then $|V(T)| \leq 6\beta(G)$.*

Proof. Let F_1, F_2, F_3 be pairwise disjoint perfect matchings of G .

(1) *Let (T, ϕ) be a 2-cut-decomposition of G such that*

- $\phi^{-1}(t) \neq \emptyset$ for each $t \in V(T)$
- $\phi^{-1}(f) \subseteq F_3$ for every edge $f \in E(T)$.

Then $|E(T)| \leq \beta(G)$.

Let $t \in V(T)$. Since $\phi^{-1}(f) \subseteq F_3$ for every edge $f \in E(T)$ incident with t , it follows that every edge $e = uv \in F_1 \cup F_2$, if $\phi(u) = t$ then $\phi(v) = t$. Moreover, since $\phi^{-1}(t) \neq \emptyset$, it follows that there is an edge $e = uv \in F_1 \cup F_2$ with $\phi(u) = t$. Consequently there is a cycle C with $E(C) \subseteq F_1 \cup F_2$ such that $V(C) \subseteq \phi^{-1}(t)$. Since this holds for each $t \in V(T)$, we deduce that there are at least $|V(T)|$ cycles within $F_1 \cup F_2$, and so $|E(T)| \leq \beta(G)$ by 4.3. This proves (1).

(2) *For every 2-cut-decomposition (T, ϕ) of G , if $\phi^{-1}(t) \neq \emptyset$ for each $t \in V(T)$ then $|E(T)| \leq 3\beta(G)$.*

For let $f \in E(T)$, and let e_1, e_2 be the two edges in $\phi^{-1}(f)$. Since $\{e_1, e_2\}$ is a cut of even cardinality of a cubic graph, it follows that every perfect matching contains an even number of members of this cut, and in particular one of F_1, F_2, F_3 includes both of e_1, e_2 . From the symmetry between

F_1, F_2, F_3 , we may assume that $\phi^{-1}(f) \subseteq F_3$ for at least one-third of all edges $f \in E(T)$, and so by contracting all other edges $f \in E(T)$, we obtain a 2-cut-decomposition (S, ϕ) with $|E(S)| \geq |E(T)|/3$, satisfying the hypotheses of (1). Consequently $|E(S)| \leq \beta(G)$, and so $|E(T)| \leq 3\beta(G)$. This proves (2).

Now we deduce the theorem. Let (T, ϕ) be a 2-cut-decomposition of G , and let n_1, n_2 and n_3 be the number of vertices of T of degree 1, 2 and at least 3 respectively. From the definition of a cut-decomposition, $\phi^{-1}(t) \neq \emptyset$ for each $t \in V(T)$ with degree one or two; so by contracting edges of T appropriately, we deduce that there is a 2-cut-decomposition (S, ϕ) of G with $|V(S)| \geq n_1 + n_2$, such that $\phi^{-1}(s) \neq \emptyset$ for each $s \in V(S)$. It follows from (2) that $|E(S)| \leq 3\beta(G)$, and so $n_1 + n_2 - 1 \leq 3\beta(G)$. Since T is a tree, it follows that $n_3 \leq n_1 - 2$, and so

$$|V(T)| = n_1 + n_2 + n_3 \leq 2n_1 + n_2 - 2 \leq 2(n_1 + n_2 - 1) \leq 6\beta(G).$$

This proves 5.1. ■

We would like to define a notion analogous to “3-hub” for a 2-cut-decomposition, but we have to be more careful. Let (T, ϕ) be a 2-cut-decomposition of G , and let $t \in V(T)$. We say that t is *solid* if for every edge $f \in E(T)$ incident with t , and every $e \in \phi^{-1}(f)$, e is incident in G with some vertex of $\phi^{-1}(t)$. Let $t \in V(T)$ be solid; let $X_0 = \phi^{-1}(t)$, and let f_1, \dots, f_k be the edges of T incident with t , where f_i is incident with a vertex of T_i for $1 \leq i \leq k$. Let G' be obtained from $G|X_0$ by adding a new edge $x_i y_i$ for $1 \leq i \leq k$, where x_i, y_i are the two vertices in X_0 incident in G with edges in $\phi^{-1}(f_i)$. We call G' the *2-hub* of G at t . We need an analogue of 4.2, as follows.

5.2 *Let (T, ϕ) be a 2-cut-decomposition of a 2-connected cubic graph G , let $t_1, \dots, t_k \in V(T)$ be solid, and for $1 \leq i \leq k$, let H_i be the 2-hub of G at t_i . For $1 \leq i \leq k$ let \mathcal{B}_i be a set of bracelets of G , such that*

- \mathcal{B}_i is a jewel-box of H_i , and
- if C is the supporting cycle of a member of \mathcal{B}_i , then $V(C) \subseteq \phi^{-1}(t)$.

Then $\mathcal{B}_1 \cup \dots \cup \mathcal{B}_k$ is a jewel-box in G .

Proof. Suppose not. Let w be the sum of the looks of a subset of $\mathcal{B}_1 \cup \dots \cup \mathcal{B}_k$, and for $i = -1, 0, 1$ let F_i be the set of all $e \in E(G)$ with $w(e) = i$; and suppose there is a cut D of G such that every edge of D belongs to F_{-1} except for one, f say, that belongs to F_0 . Choose $D = \delta(X)$ and $f \in D$ with $|D|$ minimum, and, subject to that, with $|X|$ minimum; then, as in the proof of 4.2, $G|X, G|(V(G) \setminus X)$ are both connected.

(1) *Let $e \in T$, let S_1, S_2 be the two components of $T \setminus e$, and for $i = 1, 2$ let $Y_i = \phi^{-1}(V(S_i))$. Then one of $X \cap Y_1, X \cap Y_2, Y_1 \setminus X, Y_2 \setminus X$ is empty.*

Suppose all four of these sets are nonempty. Let $\delta(Y_1) = \{a_1 a_2, b_1 b_2\}$, where $a_i, b_i \in Y_i$ for $i = 1, 2$. Since $G|X, G|(V(G) \setminus X)$ are both connected, at least one of the edges $a_1 a_2, b_1 b_2$ has both ends in X , and at least one has both ends in $V(G) \setminus X$; so we may assume that $a_1, a_2 \in X$ and $b_1, b_2 \in V(G) \setminus X$. In particular, neither of $a_1 a_2, b_1 b_2$ belongs to D ; so from the symmetry between Y_1, Y_2 , we may assume that both ends of f belong to Y_2 . Now $\delta(Y_1) \subseteq F_0$, from the choice of $\mathcal{B}_1, \dots, \mathcal{B}_k$, and so

$a_1a_2, b_1b_2 \in F_0$. Thus every edge of $\delta(X \cap Y_1)$ belongs to F_{-1} except for exactly one in F_0 , namely a_1a_2 . From the minimality of $|D|$, it follows that $|\delta(X \cap Y_1)| \geq |\delta(X)|$, and so $|\delta(X \cap Y_2)| = 2$, and $|\delta(X \cap Y_1)| = |\delta(X)|$, and yet $|X \cap Y_1| < |X|$, contrary to the minimality of $|X|$. This proves (1).

Since G is 2-connected and therefore $|D| \geq 2$, we may choose $e \in D$ with $e \neq f$. Thus $\phi(e) = -1$, and so e is an edge of the supporting cycle of some member of $\mathcal{B}_1 \cup \dots \cup \mathcal{B}_k$; say $e \in E(C_1)$ where C_1 is the supporting cycle of some $w_1 \in \mathcal{B}_1$. Let $Y_0 = \phi^{-1}(t_1)$. Let $e = xy$; then since $V(C_1) \subseteq Y_0$ it follows that $x, y \in Y_0$, and since $e \in \delta(X)$, we deduce that $X \cap Y_0, Y_0 \setminus X$ are both nonempty. Let S_1, \dots, S_s be the components of $T \setminus \{t_1\}$, and for $1 \leq i \leq s$ let e_i be the unique edge of T incident with t_1 and an end in $V(S_i)$. For $1 \leq i \leq s$, let $Y_i = \phi^{-1}(S_i)$. Thus Y_0, Y_1, \dots, Y_s are pairwise disjoint subsets of $V(G)$ with union $V(G)$. The 2-hub H_1 of G at t_1 is therefore obtained by contracting all edges of $G|Y_i$ and one of the two edges of $\phi^{-1}(e_i)$, for $1 \leq i \leq s$. Since D is not a cut of H_1 (because \mathcal{B}_1 is a jewel-box of H_1) it follows that for some $i \in \{1, \dots, s\}$, both $X \cap Y_i, Y_i \setminus X$ are nonempty. But since both $Y_0 \cap X, Y_0 \setminus X$ are nonempty, this contradicts (1) applied to the edge e_i . This proves 5.2. \blacksquare

Now we prove the main theorem of the paper, 2.2, which we restate:

5.3 *Let G be a planar cubic graph with no cutedge. Then $\beta(G) \geq |V(G)|/218659584$.*

Proof. We proceed by induction on $|V(G)|$. If G is not connected, the result follows from the inductive hypothesis applied to the components of G (for the union of jewel-boxes in different components is a jewel-box in G , as is easily seen). Thus we may assume that G is connected and hence 2-connected, since it has no cutedge. If G is 3-connected the result follows from 4.10, so we may assume that G is not 3-connected. Hence there is a 2-cut-decomposition (T, ϕ) of G ; choose such a decomposition with $|V(T)|$ maximum. (This is possible by 5.1.)

(1) *Let $t \in V(T)$ such that $\phi^{-1}(t) \neq \emptyset$. Then t is solid, and the 2-hub of G at t is 3-edge-connected.*

For let f_1, \dots, f_k be the edges of T incident with t , and let T_1, \dots, T_k be the components of $T \setminus \{t\}$, where f_i is incident with a vertex t_i of T_i for $1 \leq i \leq k$. Let $X_0 = \phi^{-1}(t)$, and for $1 \leq i \leq k$ let $X_i = \phi^{-1}(T_i)$. To prove that t is solid, let $e \in \phi^{-1}(f_1)$ say, where $e = uv$. Since $\phi^{-1}(f_1) = \delta(X_1)$, exactly one of $u, v \in X_1$, say u . We must show that $v \in X_0$. For suppose not; then v is in one of X_2, \dots, X_k , say X_2 . Let T' be the tree obtained from T by deleting the edges f_1, f_2 and adding a new vertex s adjacent to t_1, t_2, t . Then (T', ϕ) is a 2-cut-decomposition of G (to see this, note that s has degree at least three in G , and every other vertex of T has the same degree in T and in T' , except for t , and $\phi^{-1}(t) \neq \emptyset$.) But this contradicts the maximality of $|V(T)|$. This proves that t is solid.

Let H be the 2-hub of G at t , and for $1 \leq i \leq k$ let x_i, y_i be the two vertices in X_0 incident in G with an edge of $\delta(X_i)$; thus x_iy_i is an edge e_i say of H . Suppose that H is not 3-edge-connected; then there is a partition Y_1, Y_2 of $\phi^{-1}(t)$ into two nonempty subsets, such that there are exactly two edges of H between Y_1, Y_2 . Let T' be the tree obtained from T by deleting t and adding two new vertices s_1s_2 , where s_1, s_2 are adjacent, and for $1 \leq i \leq k$ s_1 is adjacent to t_i if and only if $x_i, y_i \in Y_1$, and otherwise s_2 is adjacent to t_i . (In particular, if e_i joins a vertex of Y_1 to a vertex of Y_2 then s_2 is adjacent to t_i .) Define $\phi' : V(G) \rightarrow V(T')$ by: $\phi'(v) = s_i$ if $v \in Y_i$ for $i = 1, 2$, and $\phi'(v) = \phi(v)$ if $\phi(v) \neq t$. We claim that (T', ϕ') is 2-cut-decomposition of G . Note that $\phi'^{-1}(s_i) \neq \emptyset$ for $i = 1, 2$, so it remains to check that there are exactly two edges of G in $\phi'^{-1}(s_1s_2)$. But since t is solid, this

set consists of all edges uv of G with $u \in Y_1$ and $v \in Y_2$ (which therefore belongs to $\delta_H(X_1)$) and all edges uv of G such that $u \in Y_1$, $v \in X_i$ and $e_i \in \delta_H(X_1)$. Since $|\delta_H(X_1)| = 2$, it follows that there are exactly two edges in $\phi'^{-1}(s_1s_2)$. Thus (T', ϕ') is 2-cut-decomposition of G , contrary to the maximality of $|V(T)|$. This proves (1).

Let t_1, \dots, t_k be the vertices t of T such that $\phi^{-1}(t) \neq \emptyset$, and for $1 \leq i \leq k$ let H_i be the 2-hub of G at t_i . Let Y_i be the set of edges $e = uv$ of G such that $u, v \in \phi^{-1}(t_i)$, and let $X_i = E(H_i) \setminus Y_i$. Thus $|X_i|$ is the degree of t_i in T . Let $n_i = |\phi^{-1}(t_i)|$. We claim there is a set \mathcal{B}_i of bracelets of G , such that

- \mathcal{B}_i is a jewel-box of H_i , and
- if C is the supporting cycle of a member of \mathcal{B}_i , then $V(C) \subseteq \phi^{-1}(t)$
- $|\mathcal{B}_i| \geq n_i/16819968 - |X_i|$.

For if $n_i \leq 2$ and $X_i \neq \emptyset$, we may take $\mathcal{B}_i = \emptyset$, and if $n_i \leq 2$ and $X_i = \emptyset$ then $|V(G)| \leq 2$ and the result follows from 4.3. Thus we may assume that $n_i \geq 3$, and so H_i is 3-connected, since it is cubic and 3-edge-connected by (1). By 4.10 there is a jewel-box in H_i of cardinality at least $n_i/16819968$, and at most $|X_i|$ of its members have a supporting cycle that contains a member of X_i . Removing these members gives the jewel-box \mathcal{B}_i as claimed. By 5.2, the union of these jewel-boxes is a jewel-box in G , and so

$$\sum_{1 \leq i \leq k} (n_i/16819968 - |X_i|) \leq \beta(G).$$

But $n_1 + \dots + n_k = |V(G)|$, and

$$|X_1| + \dots + |X_k| \leq 2|E(T)| \leq 12\beta(G)$$

by 5.1, so $|V(G)|/16819968 - 12\beta(G) \leq \beta(G)$, that is, $\beta(G) \geq |V(G)|/218659584$. This proves 5.3. ■

References

- [1] T. Fowler, *Unique Coloring of Planar Graphs*, Ph.D. thesis, Georgia Institute of Technology, 1998.
- [2] D. Kral, J.-S. Sereni and M. Stiebitz, “A new lower bound on the number of perfect matchings in cubic graphs”, submitted for publication (manuscript 2008).
- [3] L. Lovász and M. Plummer, *Matching Theory*, *Annals of Discrete Math.* 29, North Holland, Amsterdam, 1986.
- [4] A. Schrijver, “Counting 1-factors in regular bipartite graphs”, *J. Combinatorial Theory, Ser. B* 72 (1998), 122-135.
- [5] H. Whitney, “A theorem on graphs”, *Ann. Math.* 32 (1931), 378-390.