

Edge-Disjoint Cycles in Regular Directed Graphs

Noga Alon* Colin McDiarmid† Michael Molloy‡

February 22, 2002

Abstract

We prove that any k -regular directed graph with no parallel edges contains a collection of at least $\Omega(k^2)$ edge-disjoint cycles, conjecture that in fact any such graph contains a collection of at least $\binom{k+1}{2}$ disjoint cycles, and note that this holds for $k \leq 3$.

In this paper we consider the maximum size of a collection of edge-disjoint cycles in a directed graph. We pose the following conjecture:

Conjecture 1: *If G is a k -regular directed graph with no parallel edges, then G contains a collection of $\binom{k+1}{2}$ edge-disjoint cycles.*

We prove two weaker results:

Theorem 1: *If G is a k -regular directed graph with no parallel edges, then G contains a collection of at least $5k/2 - 2$ edge-disjoint cycles.*

Theorem 2: *If G is a k -regular directed graph with no parallel edges, then G contains a collection of at least ϵk^2 edge-disjoint cycles, where $\epsilon = \frac{3}{2^{19}}$.*

*AT & T Bell Labs, Murray Hill, NJ 07974, USA and Department of Mathematics, Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv, Israel. Research supported in part by a United States Israel BSF Grant.

†Faculty of Mathematical Sciences, University of Oxford.

‡Department of Mathematics, Carnegie Mellon University and Department of Computer Science, University of Toronto. Research supported in part by an NSERC fellowship.

Note that Theorem 1 implies that Conjecture 1 is true for $k \leq 3$. The proof of Theorem 2 is probabilistic, and we make no attempt to compute the best value of ϵ , as it is clear that our methods will not yield ϵ near $\frac{1}{2}$.

Before proving Theorems 1 and 2, we note that the bound in Conjecture 1, if true, is tight. To see this, consider the directed graph $C_n^k, n \geq 2k + 1$, which has vertex set $\{0, \dots, n - 1\}$, and edge set $\{(i, i + j) : 0 \leq i \leq n - 1, 1 \leq j \leq k\}$, where the addition is taken mod n . It is easy to see that any cycle in C_n^k must contain one of the $\binom{k+1}{2}$ edges from $\{n - k, \dots, n - 1\}$ to $\{0, \dots, k - 1\}$. (Note that C_n^k does in fact contain $\binom{k+1}{2}$ edge-disjoint cycles, as there are k edge-disjoint cycles through $n - 1$, whose deletion, along with the deletion of $n - 1$, yields a graph isomorphic to C_{n-1}^{k-1} .) Also, the complete directed graph on $k + 1$ vertices has $2\binom{k+1}{2}$ edges, and hence provides another example.

As usual, we use $\delta^+(v)$ and $\delta^-(v)$ to denote the outdegree and indegree respectively, of a vertex v . The maximum (resp. minimum) degree of a graph is the maximum (resp. minimum) of the indegrees and outdegrees of its vertices. We say a graph G is *Eulerian* if $\delta^-(v) = \delta^+(v)$ for all $v \in V(G)$. The degree of a vertex v in an Eulerian graph is $\delta^-(v)$, and an Eulerian graph is *k-regular* if each vertex degree is k . Note that here we do not require an Eulerian graph to be connected.

1 Related Work

When considering Conjecture 1, we are reminded of a few other conjectures and theorems.

Behzad, Chartrand and Wall [3] have conjectured that every k -regular digraph on n vertices has a cycle of length at most $\lceil \frac{n}{k} \rceil$. Caccetta and Häggkvist

[4] made the stronger conjecture that this is true for every digraph with minimum outdegree k . The best results in this direction are due to Chvátal and Szemerédi [5] who showed that every digraph with minimum outdegree k on n vertices has a cycle of length at most $\min(\frac{2n}{k+1}, \lceil \frac{n}{k} \rceil + 2500)$. Note that Conjecture 1, if true, implies that any k -regular digraph on n vertices has a cycle of length at most $\frac{2n}{k+1}$ (as proved in [5].)

Bermond and Thomassen [2] conjectured that any digraph with minimum outdegree k has at least $k/2$ vertex-disjoint cycles. Thomassen [11] proved that such a digraph has at least r vertex disjoint cycles if $k \geq (r + 1)!$. In Section 3, we improve this to a linear bound for graphs with minimum degree k and maximum degree $2k$.

Let us say that a directed graph G has the *cycle-packing property* if the maximum size of a collection of edge-disjoint cycles equals the minimum size of a set of edges whose removal leaves an acyclic graph. The following proposition shows that Conjecture 1 is true for any such graph.

Proposition: *If G is a directed graph with no parallel edges and minimum outdegree k , and $S \subseteq E(G)$ meets every cycle in G , then $|S| \geq \binom{k+1}{2}$.*

Proof: Since $G - S$ is acyclic there is an ordering v_1, v_2, \dots, v_n of its vertices so that for every directed edge $v_i v_j$ of $G - S$, $i < j$. It follows that v_{n-j} has at least $k - j$ outedges in S , for all $0 \leq j < k$, implying the desired result. \square

Lucchesi and Younger [7] showed in 1978 that any planar digraph has the cycle-packing property. This result has recently been extended for Eulerian flat digraphs. A graph is *flat* if it can be embedded in R^3 so that each cycle bounds a disc disjoint from the rest of the graph; and a digraph is *flat* if the underlying undirected graph is. Examples of flat graphs include the *apex* graphs, that is

graphs G such that $G \setminus v$ is planar for some vertex v . Seymour [10] shows that any Eulerian flat digraph has the cycle-packing property.

Unfortunately, these results have limited application for us here. If each outdegree in G is at least k and G has no 2-cycles, then G is not planar if $k > 2$, and G is not flat if $k > 3$.

Younger has conjectured that for any $r \geq 1$ there exist (least) integers $f(r)$ (resp. $g(r)$) such that every digraph has either a set of r edge- (resp. vertex-) disjoint cycles or a set of $f(r)$ (resp. $g(r)$) edges (resp. vertices) which meets every cycle. Soares pointed out that if $f(r), g(r)$ exist then they must be equal. McCuaig [8] showed $f(2) = 3$. For $r > 2$, $f(r)$ is not known to exist, but Alon and Seymour (see [9]) observed that if it exists then $f(r) = \Omega(r \log r)$, whereas Seymour proved in [9] that if a digraph does not have a "fractional" packing of directed cycles of value greater than k then one can delete $O(k \log k \log \log k)$ of its edges and obtain an acyclic digraph.

2 A Linear Bound

Proof of Theorem 1: We shall, in fact, show more strongly that if G is an Eulerian directed graph with no parallel edges and with minimum degree k , then G contains a collection of $5k/2 - 2$ edge-disjoint cycles. Let x be any vertex of degree k . Clearly we can find k edge-disjoint directed cycles through x . Choose a set of k such cycles $\mathcal{C} = C_1, \dots, C_k$ such that the sum of the lengths is minimum.

Claim: The union of all the arcs not incident with x in \mathcal{C} gives an acyclic graph H , and so generates a partial order $\mathcal{P}(\mathcal{C})$. Further, if $u < v$ under \mathcal{P} then

G has no $u \rightarrow v$ edge outside of \mathcal{C} .

Proof of Claim: Let D be the graph formed from all the arcs in \mathcal{C} . D is Eulerian, and x has degree k in D . Further, any such graph yields a collection of k edge-disjoint directed cycles through x . Thus by our choice of \mathcal{C} to minimize the number of arcs in the graph D , H must be acyclic, and similarly the second sentence in the claim follows. \square

Note that any two vertices, other than x , each of which lies in more than $\frac{k}{2}$ members of \mathcal{C} , are comparable, as they must lie on a common cycle. Hence, there must be some minimal element x_2 of \mathcal{P} which is less than all such vertices.

Let G_2 be the graph formed by deleting the edges of \mathcal{C} from G . As G has no multiple edges, x_2 lies in exactly one member of \mathcal{C} , and thus has degree at least $k - 1$ in G_2 . Therefore we can choose a set of $k - 1$ edge-disjoint cycles in G_2 , each passing through x_2 . Again, choose a set whose total length is minimum, and let \mathcal{P}_2 be the partial order that it induces. Remove its edges, leaving G_3 .

Let x_3 be any minimal element in \mathcal{P}_2 , and note that there is an edge from x_2 to x_3 . By our Claim, G_2 has no edges from x_2 to any vertex which lies in more than $\frac{k}{2}$ members of \mathcal{C} , and so x_3 has degree at least $\frac{k}{2}$ in G_2 . Also, x_3 lies in exactly one of the second set of cycles, and so x_3 has degree at least $\frac{k}{2} - 1$ in G_3 . Thus we can find $\frac{k}{2} - 1$ edge-disjoint cycles in G_3 , proving the theorem. \square

3 A Quadratic Bound

Note that if it were true that any Eulerian directed graph with minimum degree k has a collection of k vertex-disjoint cycles, then Conjecture 1 would follow. Unfortunately, this is not the case, for example with C_n^k where n is not a multiple

of k or with the complete directed graph on $k + 1 > 2$ vertices. However, we can prove a weaker result which is enough to give us the quadratic bound of Theorem 2, and which may be interesting in its own right:

Lemma 1: *If G is a directed graph with no parallel edges, and with minimum degree at least $k \geq 1$ and maximum degree at most $2k$, then the vertices of G may be coloured with at least $k/2^{16}$ colours (each used) in such a way that for each colour, the corresponding induced subgraph H has all vertex indegrees and outdegrees in an interval $[a, 4a]$ where $a \geq 1$.*

Before presenting the proof of this lemma, we will see that it implies Theorem 2:

Proof of Theorem 2: We set $G_k = G$, and recursively define G_j for $\lceil \frac{k}{2} \rceil \leq j \leq k$. For each j , we can apply Lemma 1 to find a collection \mathcal{C}_j of at least $j/2^{16}$ vertex disjoint cycles in G_j , and then define G_{j-1} as the graph obtained by deleting the edges of \mathcal{C}_j from G_j . Now $\mathcal{C} = \cup_{j=\lceil \frac{k}{2} \rceil}^k \mathcal{C}_j$ is a collection of edge-disjoint cycles in G , where:

$$\begin{aligned} |\mathcal{C}| &\geq \sum_{j=\lceil \frac{k}{2} \rceil}^k \frac{j}{2^{16}} \\ &\geq \frac{3}{2^{19}} k^2 \end{aligned}$$

□

Note that if the conjecture of Bermond and Thomassen discussed in Section 1 holds, then the value of ϵ in Theorem 2 can be raised to $1/4$.

The proof of Lemma 1 applies a method similar to the one used in [1] and makes use of the Lovász Local Lemma, which we state here in its symmetric case.

The Lovász Local Lemma [6]: *Let A_1, A_2, \dots, A_n be events in an arbitrary probability space, such that $\Pr(A_i) \leq p$ for each $1 \leq i \leq n$. Suppose that each event A_i is mutually independent of a set of all other events but at most d . If $ep(d+1) \leq 1$, then $\Pr(\bigcap_{i=1}^n \overline{A_i}) > 0$.*

We use this to prove:

Lemma 2: *Suppose that H is a directed graph with no parallel edges, and with minimum degree x and maximum degree y , where $x \geq 1000$, and $y \leq 4x$. Then the vertices of H can be coloured red and blue such that for any vertex $v \in V(H)$, the number of red outneighbours of v lies in the interval $[\delta_H^+(v)/2 - \delta_H^+(v)^{2/3}, \delta_H^+(v)/2 + \delta_H^+(v)^{2/3}]$ (and thus so also does the number of blue ones), and similarly the number of red inneighbours of v lies in the interval $[\delta_H^-(v)/2 - \delta_H^-(v)^{2/3}, \delta_H^-(v)/2 + \delta_H^-(v)^{2/3}]$.*

Proof: Colour each vertex of H either red or blue, making each choice independently and uniformly at random. For each v , let A_v^+ be the event that the number of red outneighbours of v does *not* lie in the interval $[\delta_H^+(v)/2 - \delta_H^+(v)^{2/3}, \delta_H^+(v)/2 + \delta_H^+(v)^{2/3}]$, and define A_v^- similarly.

For each v ,

$$\begin{aligned} \Pr(A_v^+) &\leq 2e^{-2\delta_H^+(v)^{1/3}} \\ \Pr(A_v^-) &\leq 2e^{-2\delta_H^-(v)^{1/3}}. \end{aligned}$$

Each of these probabilities is bounded above by $2e^{-2x^{1/3}}$.

Furthermore, for each v , A_v^- is mutually independent of all but at most $\sum_{u \in N^-(x)} (\delta^+(u) + \delta^-(u) - 1)$ other events, and A_v^+ is mutually independent of all but at most $\sum_{u \in N^+(x)} (\delta^+(u) + \delta^-(u) - 1)$ other events. Both sums are at

most $32x^2 - 1$.

Now, for $x \geq 1000$, $64x^2 e^{1-2x^{1/3}} < 1$. Therefore, by the Lovász Local Lemma, $\Pr(\cap_{v \in V(H)} (\overline{A_i^+} \cap \overline{A_i^-})) > 0$, and so there must be at least one satisfactory 2-colouring. \square

We now use Lemma 2 to prove Lemma 1:

Proof of Lemma 1: Let $c = 15$ and note that $2^{1-c/3}(2^{1/3}-1)^{-1} \leq \ln \frac{4}{3} \leq \frac{1}{3}$, and hence $1 - 2^{1-\frac{c}{3}}(2^{\frac{1}{3}} - 1)^{-1} \geq \frac{2}{3}$, and $\frac{2}{3}2^c \geq 1000$.

We shall see that, if $r = \lfloor \log_2 k \rfloor - c$, then G can be coloured as required with $2^r \geq 2^{-(c+1)}k$ colours. Clearly, we can assume $k \geq 2^{16}$, and so $r \geq 1$.

Let $f(x) = \frac{1}{2}x - x^{2/3}$, for $x \geq 1$. Let $z \geq k$, let $x_0 = z$, and $x_{i+1} = f(x_i)$ for $i = 1, 2, \dots$ while $x_i \geq 1$. Clearly the x_i are decreasing and $x_i \leq 2^{-i}z$ while x_i is defined. Let $1 \leq j \leq r$ be such that $x_{j-1} \geq 1$, so that x_j is defined. Then

$$\begin{aligned}
x_j &= 2^{-1}x_{j-1} - x_{j-1}^{\frac{2}{3}} \\
&= 2^{-j}z - \sum_{i=1}^j 2^{-i+1}x_{j-i}^{\frac{2}{3}} \\
&\geq 2^{-j}z - \sum_{i=1}^j 2^{-i+1}(2^{-(j-i)}z)^{\frac{2}{3}} \\
&= 2^{-j}z - 2(2^{-j}z)^{\frac{2}{3}} \left(\sum_{i=1}^j 2^{-i/3} \right) \\
&\geq 2^{-j}z - 2(2^{-j}z)^{\frac{2}{3}}(2^{1/3} - 1)^{-1} \\
&= (2^{-j}z)(1 - 2(2^{-j}z)^{-\frac{1}{3}}(2^{1/3} - 1)^{-1}) \\
&\geq (2^{-j}z)(1 - (2^{1-\frac{c}{3}})(2^{1/3} - 1)^{-1}) \\
&\geq \frac{2}{3}(2^{-j}z) \\
&\geq \frac{2}{3}2^c 2^{r-j}
\end{aligned}$$

$$\geq 1000 2^{r-j}.$$

Thus each x_j for $j = 1, \dots, r$ is defined, and satisfies $x_j \geq \frac{2}{3}(2^{-j}z) \geq 1000 2^{r-j}$.

Now let $g(x) = \frac{1}{2}x + x^{2/3}$, for $x \geq 0$. Let $y_0 = z$, and $y_{i+1} = g(y_i)$ for $i = 1, 2, \dots$. Clearly we have $y_i \geq 2^{-i}z$. Let $1 \leq j \leq r$. Then

$$\begin{aligned} y_j &= 2^{-1}y_{j-1}(1 + 2y_{j-1}^{-1/3}) \\ &= (2^{-j}z) \prod_{i=0}^{j-1} (1 + 2y_i^{-1/3}) \\ &\leq (2^{-j}z) \exp\left(\sum_{i=0}^{j-1} 2y_i^{-1/3}\right) \\ &\leq (2^{-j}z) \exp\left(2z^{-1/3} \sum_{i=0}^{j-1} 2^{i/3}\right) \\ &\leq (2^{-j}z) \exp\left(2(2^{-j}z)^{-1/3}(2^{1/3} - 1)^{-1}\right) \\ &\leq (2^{-j}z) \exp\left(2^{1-c/3}(2^{1/3} - 1)^{-1}\right) \\ &\leq \frac{4}{3}(2^{-j}z). \end{aligned}$$

Thus each y_j for $j = 1, \dots, r$ satisfies $y_j \leq \frac{4}{3}(2^{-j}z)$.

Let us use Lemma 2 to 2-colour G , then to 2-colour each of the subgraphs induced by the colour classes, and so on. Suppose that we have performed this for j levels where $0 \leq j \leq r$, and let H be one of the 2^j corresponding induced subgraphs of G . Let u and v be any two vertices of H . By the above we see that $d_H^+(u) \geq \frac{2}{3}2^{-j}d_G^+(u) \geq \frac{2}{3}2^{-j}k \geq 1000 2^{r-j}$, and $d_H^+(v) \leq \frac{4}{3}2^{-j}d_G^+(v) \leq \frac{4}{3}2^{-j}(2k)$; and there is a similar result for indegrees. Thus if H has minimum degree x and maximum degree y then $y \leq 4x$ and $x \geq 1000 2^{r-j}$. Hence, if $j < r$ we may continue to apply Lemma 2 to colour for one more level, and the lemma follows (with $a = \frac{2}{3}2^{-r}k$). \square

Remark: Using a more straightforward application of the Lovász Local Lemma (to repeatedly find vertex disjoint cycles as before, but without the iterated splitting procedure), one can also get a nearly quadratic bound with a more reasonable constant:

Theorem 3: *If G is a k -regular directed graph with no parallel edges and with $k \geq 2$, then G contains a collection of at least $k^2/8 \ln k$ edge-disjoint cycles.*

Acknowledgement We would like to thank Adrian Bondy, Xinhua Luo and Dan Younger for fruitful discussions, and Bill McCuaig and Paul Seymour for pointing out further references.

References

- [1] N. Alon, *The strong chromatic number of a graph*, Random Structures and Algorithms 3 (1992), 1-7.
- [2] J. C. Bermond and C. Thomassen, *Cycles in digraphs - a survey*, Journal of Graph Theory 5 (1981), 1 - 43.
- [3] M. Behzad, G. Chartrand and C. E. Wall, *On minimal regular digraphs with given girth*, Fund. Math. 69 (1970), 227 - 231.
- [4] L. Cacetta and R. Häggvist, *On minimal digraphs with given girth*, Proc. 9th Southeastern Conf. on Comb., Graph Theory and Computing (F. Hoffman et. al. Eds), Boca Raton, Utilitas Mathematica, Winnipeg (1978), 181-187.
- [5] V. Chvátal and E. Szemerédi, *Short cycles in directed graphs*. J. Comb. Th., Ser. B 35 (1983), 323 - 327.

- [6] P. Erdős and L. Lovász, *Problems and results on 3-chromatic hypergraphs and some related questions*, in: “Infinite and Finite Sets” (A. Hajnal et. al. Eds), Colloq. Math. Soc. J. Bolyai 11, North Holland, Amsterdam, 1975, 609-627.
- [7] C. L. Lucchesi and D. H. Younger, *A minimax theorem for directed graphs*, J. London Math. Soc. 17 (1978), 369 - 374.
- [8] W. McCuaig, *Intercyclic Digraphs*, in: “Graph Structure Theory” (N. Robertson and P. D. Seymour Eds), AMS Contemporary Math. 147 (1993), 203 - 245.
- [9] P. D. Seymour, *Packing directed circuits fractionally*, *Combinatorica*, to appear.
- [10] P. D. Seymour, *Packing circuits in Eulerian digraphs*, manuscript, 1994.
- [11] C. Thomassen, *Disjoint cycles in digraphs*, *Combinatorica* 3 (1983), 393 - 396.