

# ESTIMATES FOR NONSINGULAR MIXED CHARACTER SUMS

NICHOLAS M. KATZ

## 1. INTRODUCTION AND STATEMENT OF THE MAIN RESULT

Let  $k$  be a finite field,  $p$  its characteristic,  $q$  its cardinality,

$$\psi : (k, +) \rightarrow \mathbb{Z}[\zeta_p]^\times \subset \mathbb{C}^\times$$

a nontrivial additive character of  $k$ , and

$$\chi : k^\times \rightarrow \mathbb{Z}[\zeta_{q-1}]^\times \subset \mathbb{C}^\times$$

a nontrivial multiplicative character of  $k$ . We extend  $\chi$  to  $k$  by defining  $\chi(0) = 0$ .

We wish to consider character sums over  $\mathbb{A}^n$ ,  $n \geq 1$ , of the following form. We are given a polynomial  $f(x) := f(x_1, \dots, x_n)$  in  $k[x_1, \dots, X_n]$  of degree  $d \geq 1$ , and we are given a second polynomial  $g(X) := g(x_1, \dots, x_n)$  in  $k[x_1, \dots, x_n]$  of degree  $e \geq 1$ . We are interested in understanding when the sum

$$\sum_{x \in k^n} \psi(f(x))\chi(g(x))$$

has “square root” cancellation, i.e., when we can exhibit an explicit constant  $C = C(n, d, e)$  and prove the estimate

$$\left| \sum_{x \in k^n} \psi(f(x))\chi(g(x)) \right| \leq C(\#k)^{n/2}.$$

In this paper, we will exhibit one particularly nice class of pairs  $(f, g)$  for which such estimates hold. The general problem of understanding for which pairs  $(f, g)$  one has, or should have, such estimates is far from being understood.

Let us first recall the notion of a “Deligne polynomial”. A polynomial  $f = f(x_1, \dots, x_n)$  in  $n \geq 1$  variables over  $k$  of degree  $d \geq 1$  is called a Deligne polynomial if its degree  $d$  is prime to  $p$  and if its highest degree term,  $f_d$ , is a homogeneous form of degree  $d$  in  $n$  variables which is nonzero, and whose vanishing, if  $n \geq 2$ , defines a smooth hypersurface in the projective space  $\mathbb{P}^{n-1}$ .

For  $f = f(x_1, \dots, x_n)$  a Deligne polynomial of degree  $d$ , one has Deligne's fundamental estimate [De-Weil I, 8.4]

$$\left| \sum_{x \in k^n} \psi(f(x)) \right| \leq (d-1)^n (\#k)^{n/2}.$$

If  $g = g(x_1, \dots, x_n)$  is a Deligne polynomial of degree  $e$ , such that  $g = 0$  defines a smooth hypersurface in  $\mathbb{A}^n$ , then one has the analogous estimate [Ka-ENSMCS, Thm. 1]

$$\left| \sum_{x \in k^n} \chi(g(x)) \right| \leq (e-1)^n (\#k)^{n/2}.$$

Our main result is that if  $f$  and  $g$  above are suitably transverse, then we have a good estimate for the mixed sum. To state the estimate, we define the constant

$$\begin{aligned} C(n, d, e) &:= (-1)^n \times \text{coef. of } L^n \text{ in } \frac{(1+L)^{n+1}}{(1+L)(1+dL)(1+eL)} \\ &= \text{the value at } (x, y) = (d, e) \text{ of } \frac{x(x-1)^n - y(y-1)^n}{x-y} \\ &= \sum_{a+b=n} (d-1)^a (e-1)^b + \sum_{a+b=n-1} (d-1)^a (e-1)^b. \end{aligned}$$

Recall also that, given an integer  $w$ , a number  $\alpha \in \mathbb{C}$  is said to be pure of weight  $w$  (relative to  $q$ ) if it and all its  $\text{Aut}(\mathbb{C}/\mathbb{Q})$ -conjugates have absolute value  $q^{w/2}$ . Such an  $\alpha$  is necessarily algebraic over  $\mathbb{Q}$ . A polynomial  $P(T) \in 1 + T\mathbb{C}[T]$  is said to be pure of weight  $w$  if all its reciprocal roots are pure of weight  $w$ ; it is said to be mixed of weight  $\leq w$  if each of its reciprocal roots  $\alpha$  is pure of some integer weight  $w_\alpha \leq w$ .

**Theorem 1.1.** *Suppose that  $f = f(x_1, \dots, x_n)$  and  $g = g(x_1, \dots, x_n)$  are Deligne polynomials over  $k$  of degrees  $d$  and  $e$  respectively. If  $n \geq 2$ , suppose in addition that the smooth hypersurfaces in  $\mathbb{P}^{n-1}$  defined by  $f_d = 0$  and by  $g_e = 0$  are transverse, i.e., their intersection is smooth of codimension 2 in  $\mathbb{P}^{n-1}$ . Then we have the following results.*

(1) *We have the estimate*

$$\left| \sum_{x \in k^n} \psi(f(x)) \chi(g(x)) \right| \leq C(n, d, e) (\#k)^{n/2}.$$

*The associated  $L$  function is a polynomial  $P(T)$  (for  $n$  odd) or a reciprocal polynomial  $1/P(T)$  (for  $n$  even) of degree  $\leq C(n, d, e)$ , which is mixed of weight  $\leq n$ .*

(2) *If  $P(T)$  has degree  $= C(n, d, e)$ , then  $P(T)$  is pure of weight  $n$ .*

- (3) If  $g = 0$  defines a nonsingular hypersurface in  $\mathbb{A}^n$ , then  $P(T)$  has degree  $= C(n, d, e)$ , and is pure of weight  $n$ .

We are indebted to Steve Sperber for the observation that the ideas which go into proving this theorem lead in a straightforward way to a theorem dealing with the following more general situation. Instead of  $(f, g)$ , we give ourselves an integer  $r \geq 1$ , and  $r + 1$  Deligne polynomials  $(f, g_1, \dots, g_r)$  in  $n$  variables over  $k$ , of degrees  $(d, e_1, \dots, e_r)$ . If  $n \geq 2$ , we assume that the  $r + 1$  smooth hypersurfaces in  $\mathbb{P}^{n-1}$  defined by the vanishing of their highest degree forms are transverse, in the sense that for any integer  $j$  with  $r + 1 \geq j \geq 1$ , the intersection of any  $j$  of them is smooth of codimension  $j$  in  $\mathbb{P}^{n-1}$  if  $j \leq n - 1$ , and is empty if  $j \geq n$ . Then we get the following result.

**Theorem 1.2.** *For any  $r$ -tuple of nontrivial multiplicative characters  $(\chi_1, \dots, \chi_r)$ , we have the bound*

$$\left| \sum_{x \in k^n} \psi(f(x)) \prod_i \chi_i(g_i(x)) \right| \leq C(n, d, e_1, \dots, e_r) (\#k)^{n/2},$$

where  $C(n, d, e_1, \dots, e_r)$  is defined as

$$C(n, d, e_1, \dots, e_r) := (-1)^n \times \text{coef. of } L^n \text{ in } \frac{(1 + L)^{n+1}}{(1 + L)(1 + dL) \prod_i (1 + e_i L)}.$$

We will discuss the proof of this more general result in the appendix.

## 2. STATEMENT OF A SECOND VERSION OF THE MAIN RESULT

In this section, we give a generalization in the spirit of [Ka-SE, 5.1.1] and [Ka-ENSMCS, Thm.'s 3,4]. Let  $X/k$  be a projective, smooth, and geometrically connected  $k$ -scheme of dimension  $n \geq 1$ , given with a projective embedding  $X \hookrightarrow \mathbb{P}_k^N := \mathbb{P}$ . We fix integers  $d \geq 1$  and  $e \geq 1$ , both prime to  $p$ . We are given a linear form

$$Z \in H^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(1)),$$

a degree  $d$  form

$$F \in H^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(d)),$$

and a degree  $e$  form

$$G \in H^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(e)),$$

all on the ambient projective space  $\mathbb{P}$ . Assume that the following four transversality hypotheses hold.

- (1)  $X \cap Z$  is lisse of codimension 1 in  $X$ .
- (2)  $X \cap Z \cap F$  is lisse of codimension 1 in  $X \cap Z$  ( $=$  empty, if  $n = 1$ ).

- (3)  $X \cap Z \cap G$  is lisse of codimension 1 in  $X \cap Z$  ( $:=$  empty, if  $n = 1$ ).
- (4)  $X \cap Z \cap F \cap G$  is lisse of codimension 2 in  $X \cap Z$  ( $:=$  empty, if  $n \leq 2$ ).

To this data, we attach the smooth affine  $k$ -scheme

$$V := X - X \cap Z = X[1/Z],$$

and the functions

$$f := F/Z^d : V \rightarrow \mathbb{A}_k^1$$

and

$$g := G/Z^e : V \rightarrow \mathbb{A}_k^1.$$

We denote by  $c(X)$  the total Chern class of  $X$ , and by  $L$  the class of  $\mathcal{O}_X(1)$ . We define the constant  $C(X, d, e)$  by

$$C(X, d, e) := (-1)^n \int_X \frac{c(X)}{(1+L)(1+dL)(1+eL)}.$$

Thus when  $X$  is  $\mathbb{P}^n$  with the identity embedding of itself into  $\mathbb{P} = \mathbb{P}^n$ ,  $C(X, d, e)$  is the constant  $C(n, d, e)$  of the first section. When  $X$  is a complete intersection in  $\mathbb{P}^{n+r}$  of multidegree  $(a_1, \dots, a_r)$ , then

$$\begin{aligned} C(X, d, e) &:= (-1)^n \int_{\mathbb{P}^{n+r}} \frac{a_1 \dots a_r L^r (1+L)^{n+r+1}}{(1+L)(1+dL)(1+eL) \prod_i (1+a_i L)} \\ &= \text{coef. of } L^n \text{ in } \frac{a_1 \dots a_r (1+L)^{n+r+1}}{(1+L)(1+dL)(1+eL) \prod_i (1+a_i L)} \end{aligned}$$

**Theorem 2.1.** *Suppose that  $(X, Z, F, G)$  are as above. Then we have the following results.*

- (1) *We have the estimate*

$$\left| \sum_{x \in V(k)} \psi(f(x)) \chi(g(x)) \right| \leq C(X, d, e) (\#k)^{n/2}.$$

*The associated  $L$  function is a polynomial  $P(T)$  (for  $n$  odd) or a reciprocal polynomial  $1/P(T)$  (for  $n$  even) of degree  $\leq C(X, d, e)$ , which is mixed of weight  $\leq n$ .*

- (2) *If  $P(T)$  has degree  $= C(X, d, e)$ , then  $P(T)$  is pure of weight  $n$ .*
- (3) *If  $X \cap G$  is smooth of codimension 1 in  $X$ , or equivalently if  $g = 0$  is smooth of codimension 1 in  $V$ , then  $P(T)$  has degree  $= C(X, d, e)$ , and is pure of weight  $n$ .*

Thus when  $X$  is  $\mathbb{P}^n$  with the identity embedding of itself into  $\mathbb{P} = \mathbb{P}^n$ , this theorem is just Theorem 1.1.

## 3. PROOF OF THEOREM 2.1; THE STRATEGY

As customary in such questions, we choose a prime number  $\ell \neq p$  and choose an embedding of  $\mathbb{Q}(\zeta_p, \zeta_{q-1})$  into  $\overline{\mathbb{Q}}_\ell$ , so that we can view all our characters, both additive and multiplicative, as having values in  $\overline{\mathbb{Q}}_\ell^\times$ , and so that we can apply  $\ell$ -adic cohomology.

On the smooth, geometrically connected, affine variety  $V[1/g]$  of dimension  $n$ , we have the lisse, rank one, Artin-Schreier sheaf  $\mathcal{L}_{\psi(f)}$ , the lisse, rank one, Kummer sheaf  $\mathcal{L}_{\chi(g)}$ , and their lisse, rank one, tensor product  $\mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)}$ , cf. [De-ST, 1.4.2, 1.4.3]. Each of these lisse sheaves is pure of weight 0. By the Lefschetz Trace Formula [Gr-Rat], we have

$$\begin{aligned} & \sum_{x \in V(k)} \psi(f(x))\chi(g(x)) \\ &= \sum_i (-1)^i \text{Trace}(Frob_k | H_c^i(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)})). \end{aligned}$$

By Deligne's Weil II result [De-Weil II, 3.3.4], the (reversed characteristic polynomial of  $Frob_k$  on the) cohomology group  $H_c^i$  above is mixed of weight  $\leq i$ . By the dual of the Lefschetz affine theorem,  $H_c^i$  vanishes for  $i < n$ , cf. [SGA 4 Tome 3, Exposé XVIII, Thm. 3.2.5 and Exposé XIV, Cor. 3.2].

Let us admit temporarily the following theorem, and explain how it implies Theorem 2.1.

**Theorem 3.1.** *Suppose that  $(X, Z, F, G)$  are as in Theorem 2.1. Then we have the following results.*

- (1)  $H_c^i := H_c^i(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)})$  vanishes for  $i \neq n$ .
- (2) If  $X \cap G$  is smooth of codimension 1 in  $X$ , or equivalently if  $g = 0$  is smooth of codimension 1 in  $V$ , then we have the following results.
  - (2a)  $H_c^n$  has dimension  $C(X, d, e)$ , and is pure of weight  $n$ .
  - (2b) The "forget supports" map is an isomorphism

$$H_c^n(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)}) \cong H^n(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)}).$$

Using this, we prove Theorem 2.1 as follows. Over the affine space

$$\mathbb{A} := H^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(1)) \times H^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(d)) \times H^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(e)),$$

we have the product  $X \times \mathbb{A}$ , the closed subscheme of this product consisting of points  $(x \in X, Z, F, G)$  where  $L(x)G(x) = 0$ , and its open complement  $\mathcal{V}[1/g]_{univ}$ , consisting of points  $(x \in X, Z, F, G)$  where  $L(x)G(x)$  is invertible. We view  $\mathcal{V}[1/g]_{univ}$  as fibred over  $\mathbb{A}$ , say

$$\pi_{univ} : \mathcal{V}[1/g]_{univ} \rightarrow \mathbb{A}.$$

The triples  $(Z, F, G) \in \mathbb{A}$  which satisfy our four transversality conditions with respect to  $X$  form a dense open set  $\mathcal{U} \subset \mathbb{A}$ . Over this open set  $\mathcal{U} \subset \mathbb{A}$ , the pullback  $\mathcal{V}[1/g]$  of  $\mathcal{V}[1/g]_{univ}$  is an affine smooth  $\mathcal{U}$ -scheme, say

$$\pi : \mathcal{V}[1/g] \rightarrow \mathcal{U}.$$

with geometrically connected fibres of dimension  $n$ , whose fibre over a point  $(Z, F, G)$  is  $V[1/g] = X[1/LG]$ .

On  $\mathcal{V}[1/g]$ , we have the lisse sheaf  $\mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)}$ . The sheaf

$$\mathcal{N} := R^n \pi_1(\mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)})$$

is then a sheaf of perverse origin on  $\mathcal{U}$ , cf. [Ka-SCMD, Introduction and Cor. 5]. For a sheaf of perverse origin, one knows [Ka-SCMD, Prop.'s 11, 12] that the stalk at any point has rank at most the generic rank, and that the open set  $\mathcal{U}_{lisse}$  where the sheaf is lisse consists precisely of the points  $\mathcal{U}_{max}$  where the stalk has this maximum rank.

The stalk of  $\mathcal{N}$  at a  $k$ -valued point  $(Z, F, G) \in \mathcal{U}(k)$  is the cohomology group

$$H_c^n(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)}).$$

The supplementary condition on  $(Z, F, G)$  that  $X \cap G$  be smooth of codimension 1 in  $X$  defines a dense open set  $\mathcal{U}_1 \subset \mathcal{U}$ . By the second part of Theorem 3.1, the stalk of  $\mathcal{N}$  at any point of  $\mathcal{U}_1$  has rank  $C(X, d, e)$ , and this stalk is pure of weight  $n$ . [Let us note in passing that this proves part (3) of Theorem 2.1.]

Therefore the generic rank of  $\mathcal{N}$  must be  $C(X, d, e)$ . So for any  $k$ -valued point  $(Z, F, G) \in \mathcal{U}(k)$ , we have

$$\dim H_c^n(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)}) \leq C(X, d, e).$$

As this group is mixed of weight  $\leq n$ , and all other  $H_c^i$  vanish, we have

$$\sum_{x \in V(k)} \psi(f(x)) \chi(g(x)) = (-1)^n \text{Trace}(\text{Frob}_k | H_c^n),$$

so we get the estimate

$$\left| \sum_{x \in V(k)} \psi(f(x)) \chi(g(x)) \right| \leq C(X, d, e) (\#k)^{n/2}.$$

This proves part (1) of Theorem 2.1.

On the dense open set  $\mathcal{U}_1$ ,  $\mathcal{N}$  is punctually pure of weight  $n$ , and has constant rank  $C(X, d, e)$ . Thus we have the inclusion  $\mathcal{U}_1 \subset \mathcal{U}_{lisse} = \mathcal{U}_{max}$ . Now the sheaf  $\mathcal{N}$  is mixed, by [De-Weil II, 3.3.3], so its restriction to  $\mathcal{U}_{lisse}$  is a lisse sheaf which is mixed. Such a sheaf on a lisse  $k$ -scheme is a successive extension of pure lisse sheaves, by [De-Weil II, 3.4.1], so the weights which occur, and their multiplicities, can be read by looking

at any single point in  $\mathcal{U}_{\text{isse}}(\bar{k})$ . Taking a point in  $\mathcal{U}_1(\bar{k})$ , we conclude that  $\mathcal{N}|_{\mathcal{U}_{\text{max}}}$  is pure of weight  $n$ . This proves the second assertion of Theorem 2.1.

#### 4. PROOF OF PART (1) OF THEOREM 3.1

Let us recall the situation. We have  $X/k$  a projective, smooth, and geometrically connected  $k$ -scheme of dimension  $n \geq 1$ , given with a projective embedding  $X \hookrightarrow \mathbb{P}_k^N := \mathbb{P}$ . And we have homogeneous forms  $(Z, F, G)$  of prime-to- $p$  degrees  $1, d, e$  respectively in the ambient  $\mathbb{P}$ , subject to various transversality conditions. We must show that

$$H_c^i := H_c^i(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)})$$

vanishes for  $i \neq n$ . As already noted earlier,  $H_c^i$  vanishes for  $i < n$  by the dual of the Lefschetz affine theorem. So it remains to show that  $H_c^i$  vanishes for  $i > n$ .

We first treat the case where  $\chi^e$  is trivial. In this case, we argue as follows. Consider the finite flat covering  $V_e := V[g^{1/e}]$  of  $V$  gotten by taking the  $e$ 'th root of  $g$ , say

$$\rho : V_e \rightarrow V.$$

Concretely,  $V_e$  is the closed subscheme of  $V \times \mathbb{A}^1$ , with coordinate  $t$  on  $\mathbb{A}^1$ , of equation  $t^e = g$ . The direct image sheaf  $\rho_* \bar{\mathbb{Q}}_\ell$  on  $V$  has a direct sum decomposition, as the direct sum of the constant sheaf on  $V$  with various Kummer sheaves on  $V[1/g]$ , extended by zero. More precisely, denote by

$$j : V[1/g] \subset V$$

the inclusion. We have a direct sum decomposition on  $V$

$$\rho_* \bar{\mathbb{Q}}_\ell = \bar{\mathbb{Q}}_\ell \bigoplus \bigoplus_{\Lambda^{e \text{triv}, \Lambda} \text{ nontriv}} j_* \mathcal{L}_{\Lambda(g)}.$$

By the projection formula, we see that for each  $i$ ,

$$H_c^i(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)})$$

is a direct summand of

$$\begin{aligned} H_c^i(V_e \otimes_k \bar{k}, \rho^* \mathcal{L}_{\psi(f)}) &= H_c^i(V \otimes_k \bar{k}, \rho_* \rho^* \mathcal{L}_{\psi(f)}) = H_c^i(V \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \rho_* \bar{\mathbb{Q}}_\ell) \\ &= H_c^i(V \otimes_k \bar{k}, \mathcal{L}_{\psi(f)}) \bigoplus \bigoplus_{\Lambda^{e \text{triv}, \Lambda} \text{ nontriv}} H_c^i(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\Lambda(g)}). \end{aligned}$$

[We note for later use that this same projection formula argument shows that for each  $i$ , the ordinary cohomology group

$$H^i(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)})$$

is a direct summand of the ordinary cohomology group

$$H^i(V_e \otimes_k \bar{k}, \rho^* \mathcal{L}_{\psi(f)}).]$$

So it suffices to show that the cohomology groups

$$H_c^i(V_e \otimes_k \bar{k}, \rho^* \mathcal{L}_{\psi(f)})$$

vanish for  $i > n$ . We will see that this results from the “nonsingular” case ( $\epsilon = \delta = -1$ ) of [Ka-ESES, Thm. 4]. For this, we argue as follows. We began with  $X \subset \mathbb{P}$ . To fix ideas, think of this ambient  $\mathbb{P} = \mathbb{P}^N$  as having homogeneous coordinates  $(X_0, \dots, X_N)$ . In the projective space  $\mathbb{P}^{N+1}$ , with homogeneous coordinates  $(T, X_0, \dots, X_N)$ , we consider the closed subscheme  $X_e$  defined by the equations which defined  $X$ , together with the new equation

$$T^e - G = 0.$$

Then  $(X_0, \dots, X_N)$  define a map from  $X_e$  to  $X$ , which makes  $X_e$  a finite flat covering of  $X$  of degree  $e$ . [A more intrinsic way to view  $X_e$  is as follows. On  $X$ , we have the invertible  $\mathcal{O}_X$ -module  $\mathcal{M} := \mathcal{O}_X(1)$ , and the global section  $m := G$  of  $\mathcal{M}^{\otimes e}$ . Then  $X_e$  represents the functor on  $X$ -schemes which attaches to an  $X$ -scheme  $\pi : Y \rightarrow X$  the set

$$\{z \in H^0(Y, \pi^* \mathcal{M}) \mid z^e = \pi^* m \text{ in } H^0(Y, \pi^* \mathcal{M}^{\otimes e}).\}$$

Inside  $X_e$ ,  $V_e$  is the open set  $X_e - X_e \cap Z = X_e[1/Z]$ , and  $\rho^* \mathcal{L}_{\psi(f)}$  on  $V_e$  is just  $\mathcal{L}_{\psi(f)}$ , for  $f$  the “same” function  $F/Z^d$ , but now viewed on  $V_e = X_e[1/Z]$ . A rereading of [Ka-ESES, Lemma 10 made cohomological, Cor. 14(1), and the first paragraph of the proof of Thm. 16], shows that the asserted vanishing of the cohomology groups

$$H_c^i(V_e \otimes_k \bar{k}, \rho^* \mathcal{L}_{\psi(f)})$$

for  $i > n$  is proven (though not explicitly stated!) in [Ka-ESES], provided that the following three conditions hold.

- (1)  $X_e$  is Cohen-Macaulay and equidimensional of dimension  $n$ .
- (2)  $X_e \cap Z$  is smooth of dimension  $n - 1$ .
- (3)  $X_e \cap Z \cap F$  is smooth of dimension  $n - 2$  ( $:=$  empty, if  $n=1$ ).

To see that these three conditions hold, we argue as follows. To show  $X_e$  is Cohen-Macaulay and equidimensional of dimension  $n$ , we argue as follows. The scheme  $X_e$  is the finite flat covering of  $X$  defined by taking the  $e$ 'th root of  $G$ . The open set  $X_e[1/G] \subset X$  is finite etale over  $X[1/G]$ , so is itself smooth. And over an open neighborhood  $U$  of a point  $x \in X$  where  $G(x) = 0$ , the covering  $X_e$  is a hypersurface in the smooth scheme  $U \times \mathbb{A}^1$ , so is Cohen-Macaulay, cf. [A-K, Chpt. III, Cor. 4.5]. To see that  $X_e \cap Z$  is smooth of dimension  $n - 1$ , view it as the covering of  $X \cap Z$  defined by taking the  $e$ 'th root of  $G$ . By hypothesis  $G = 0$  defines a smooth hypersurface in the smooth scheme  $X \cap Z$  of dimension  $n - 1$ , and  $e$  is prime to  $p$ , so the total space  $X_e \cap Z$



of this covering is itself smooth. Similarly,  $X_e \cap Z \cap F$  is the covering of  $X \cap Z \cap F$  defined by taking the  $e$ 'th root of  $G$ , and we argue as above, now using the assumed smoothness of both  $X \cap Z \cap F$  and of  $X \cap Z \cap F \cap G$ . This concludes the proof of the first part of Theorem 3.1, in the case when  $\chi^e$  is trivial.

We now explain how to reduce the general case to this one. The asserted vanishing of the cohomology groups is a geometric statement, so we may extend scalars at will from the original finite field  $k$  to any finite extension. Our first task is to show that after such an extension, we can find a particularly nice coordinate system  $(Y_0, \dots, Y_N)$  in the ambient  $\mathbb{P}$ , which is suitably transverse to the situation  $(X, Z, F, G)$ . We will inductively find these homogeneous coordinates, or rather the hyperplanes they define. We start by defining

$$Y_0 := Z.$$

We wish to find a coordinate system  $(Z = Y_0, Y_1, \dots, Y_N)$  in  $\mathbb{P}$  such that the following conditions hold.

- (1)  $X$  is transverse to the coordinate system  $(Y_0, \dots, Y_N)$ , in the sense that for any subset  $I \subset \{0, 1, \dots, N\}$ , the intersection  $X \cap \bigcap_{i \in I} Y_i$  is smooth of dimension  $\dim X - \#I$  ( $:=$  empty if  $\#I > \dim X$ ).
- (2) If  $X \cap G$  is smooth, then it is transverse to the coordinate system  $(Y_0, \dots, Y_N)$ .
- (3) Each of  $X \cap Z$ ,  $X \cap Z \cap F$ ,  $X \cap Z \cap G$ , and  $X \cap Z \cap F \cap G$ , viewed as a closed smooth subscheme of  $\mathbb{P} \cap Z$ , is transverse to the coordinate system  $(Y_1, \dots, Y_N)$ .

It is standard that, over  $\bar{k}$ , given any finite list of smooth, equidimensional subschemes  $W_i \subset \mathbb{P}$ , we can find a hyperplane  $Y_1 = 0$  in  $\mathbb{P}$  which is transverse to each  $W_i$ , in the sense that  $W_i \cap Y_1$  is smooth of codimension 1 in  $W_i$  ( $:=$  empty, if  $\dim(W_i) = 0$ ). We apply this with the list taken to be  $Z, X, X \cap Z, X \cap Z \cap F, X \cap Z \cap G, X \cap Z \cap F \cap G$ , and, in the second part of Theorem 3.1,  $X \cap G$  itself. This produces the desired  $Y_1$ . To define  $Y_2$ , we consider this list of  $W_i$ 's, augmented by adding their intersections, when nonempty, with  $Y_1$ . We then continue, at each step keeping the terms on our previous list of smooth subschemes of  $\mathbb{P}$  and adding on their intersections, when nonempty, with the previously obtained hyperplane. In this way, we get the desired coordinate system  $(Z = Y_0, Y_1, \dots, Y_N)$  in the ambient  $\mathbb{P}$ , defined over some finite extension of  $k$ , which is suitably transverse to our original situation. Thus it suffices to treat the case where our original

coordinate system  $(X_0, \dots, X_N)$  has  $Z = X_0$  and is suitably transverse to  $(X, Z, F, G)$  as above.

Pick a prime-to- $p$  integer  $r$  such that  $\chi^r$  is trivial (e.g., one might take  $r$  to be  $\#k - 1$ ). Consider the “ $r$ ’th power map”

$$[r] : \mathbb{P} \rightarrow \mathbb{P}, (X_0, \dots, X_N) \mapsto (X_0^r, \dots, X_N^r).$$

It is finite and flat of degree  $r^N$ , and finite étale over the dense open set where all  $Y_i$  are invertible.

**Lemma 4.1.** *We have the following results.*

- (1) *Suppose we are given a closed subscheme  $W \subset \mathbb{P}$  which is smooth and equidimensional, and which is transverse to the coordinate system  $(X_0, \dots, X_N)$ , in the sense that for any subset  $I \subset \{0, 1, \dots, N\}$ , the intersection  $W \cap \bigcap_{i \in I} X_i$  is smooth of dimension  $\dim W - \#I$  ( $:=$  empty if  $\#I > \dim W$ ). Then its inverse image  $W_r$  in the covering  $[r] : \mathbb{P} \rightarrow \mathbb{P}$ , is smooth.*
- (2) *For any closed subscheme  $W \subset \mathbb{P}$ , the intersection  $W_r \cap Z$  is the inverse image of  $W \cap Z \subset \mathbb{P} \cap Z$  under the “ $r$ ’th power map”*

$$[r : Z] : \mathbb{P} \cap Z \rightarrow \mathbb{P} \cap Z, (X_1, \dots, X_N) \mapsto (X_1^r, \dots, X_N^r).$$

- (2) *For any closed subscheme  $W \subset \mathbb{P}$  such that  $W \cap Z$  is smooth,  $W_r \cap Z$  is smooth.*

*Proof.* (1) Since  $k$  is perfect, it suffices to show that  $W_r$  is a regular scheme. Over a  $\bar{k}$ -valued point  $w$  of  $W$  where all the  $X_i$  are invertible, our covering is finite étale. Over a  $\bar{k}$ -valued point  $w$  of  $W$  where precisely the  $X_i, i \in I$  vanish, with  $\#I \geq 1$ , pick some index  $j$  with  $X_j$  invertible at  $w$ , and consider the functions  $x_i := X_i/X_j$ . By the transversality hypothesis, these  $x_i$  are part of a system of parameters at  $w$ . Our covering over an open neighborhood of  $w$  is an étale covering of degree  $r^{N-\#I}$  of the finite flat covering obtained by extracting the  $r$ ’th roots of the  $x_i$ . In this finite flat covering, there is a unique point over  $w$ , whose local ring is visibly regular. Thus  $W_r$  is a regular scheme.

(2) This is a tautology.

(3) By (2), this results from (1) applied to  $W \cap Z \subset \mathbb{P} \cap Z$  and the map

$$[r : Z] : \mathbb{P} \cap Z \rightarrow \mathbb{P} \cap Z, (X_1, \dots, X_N) \mapsto (X_1^r, \dots, X_N^r).$$

□

We now consider the pullback of our situation  $(X, Z, F, G)$  by the map  $[r] : \mathbb{P} \rightarrow \mathbb{P}$ . We obtain  $(X_r, Z_r = Z^r, F_r, G_r)$ . Here  $F_r(X_i) := F(X_i^r)$ ,  $G_r(X_i) := G(X_i^r)$ . We have  $Z_r = Z^r$  because by construction

we have  $Z = X_0$ . We put  $V_r := X_r[1/Z_r] = X_r[1/Z]$ ,  $f_r := F_r/Z^{rd}$ , and  $g_r := G_r/Z^{re}$ . We have a finite flat map

$$[r]_{V[1/g]} : V_r[1/g_r] \rightarrow V[1/g]$$

of degree  $r^N$ . By the projection formula, for each  $i$  the cohomology group

$$H_c^i(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)})$$

is a direct summand of the cohomology group

$$H_c^i(V_r[1/g_r] \otimes_k \bar{k}, \mathcal{L}_{\psi(f_r)} \otimes \mathcal{L}_{\chi(g_r)}).$$

[We remark for later use that this same projection formula argument shows that for each  $i$  the ordinary cohomology group

$$H^i(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)})$$

is a direct summand of the ordinary cohomology group

$$H^i(V_r[1/g_r] \otimes_k \bar{k}, \mathcal{L}_{\psi(f_r)} \otimes \mathcal{L}_{\chi(g_r)}).]$$

We claim that the data  $(X_r, Z, F_r, G_r)$  satisfies all the transversality conditions of section 2, but now with degrees  $(d, e)$  replaced by degrees  $(dr, er)$ . First of all,  $X_r$  is geometrically connected, because at any of the finitely many points where exactly  $n$  of the  $X_i$  intersect  $X$ , the covering  $[r]_X : X_r \rightarrow X$  is fully ramified. But if  $X_r$  were not geometrically connected, each of its connected components would map onto  $X$ .

The transversality hypotheses of section 2 are that  $X \cap Z$ ,  $X \cap Z \cap F$ ,  $X \cap Z \cap G$ , and  $X \cap Z \cap F \cap G$ , are all smooth of the correct dimension (:=empty, if that dimension is negative). Their inverse images under  $[r] : \mathbb{P} \rightarrow \mathbb{P}$  are the schemes  $X_r \cap Z$ ,  $X_r \cap Z \cap F_r$ ,  $X_r \cap Z \cap G_r$ , and  $X_r \cap Z \cap F_r \cap G_r$ . That these inverse images (and also  $X_r \cap G_r$ , if  $X \cap G$  is assumed smooth) are all smooth of the correct dimension (:=empty, if that dimension is negative) results from Lemma 4.1.

But in this situation,  $\chi^{er}$  is trivial, so the cohomology groups

$$H_c^i(V_r[1/g_r] \otimes_k \bar{k}, \mathcal{L}_{\psi(f_r)} \otimes \mathcal{L}_{\chi(g_r)})$$

vanish for  $i \neq n$ . And hence their direct summands

$$H_c^i(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)})$$

vanish for  $i \neq n$ . This concludes the proof of part (1) of Theorem 3.1.

## 5. PROOF OF PART (2) OF THEOREM 3.1

Let us recall the situation. We start with  $(X, Z, F, G)$ , but now we assume that  $X \cap Z$ ,  $X \cap Z \cap F$ ,  $X \cap Z \cap G$ ,  $X \cap Z \cap F \cap G$ , and *in addition*  $X \cap G$ , are all smooth of the correct dimension (:=empty, if that dimension is negative). We first show that

$$H_c^n(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)})$$

is pure of weight  $n$ .

We first explain how to reduce to the case when  $\chi^e$  is trivial. Exactly as in the previous section, we pick a prime-to- $p$  integer  $r$  so that  $\chi^r$  is trivial, extend scalars so that  $(Z = X_0, X_1, \dots, X_N)$  is a suitably transverse coordinate system, and pass to the situation  $(X_r, Z, F_r, G_r)$ , for which all of these smoothness assumptions still hold. Our cohomology group

$$H_c^n(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)})$$

is a direct factor of

$$H_c^n(V_r[1/g_r] \otimes_k \bar{k}, \mathcal{L}_{\psi(f_r)} \otimes \mathcal{L}_{\chi(g_r)}).$$

So we are reduced to proving that  $H_c^n(V_r[1/g_r] \otimes_k \bar{k}, \mathcal{L}_{\psi(f_r)} \otimes \mathcal{L}_{\chi(g_r)})$  is pure of weight  $n$ .

So it suffices to considering the situation  $(X, Z, F, G)$  of the paragraph above, but under the additional hypothesis that  $\chi^e$  is trivial. In this case, we return to the considerations of the first part of section 4, where we introduced the covering  $V_e$  defined by taking the  $e$ 'th root of  $g$ , and saw that our cohomology group

$$H_c^n(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)})$$

was a direct factor of

$$H_c^n(V_e \otimes_k \bar{k}, \mathcal{L}_{\psi(f)}).$$

In this situation, the key observation is that  $X_e$  is in fact smooth. Indeed, it is the covering of  $X$  defined by extracting the  $e$ 'th root of  $G$ . But  $e$  is prime to  $p$ ,  $X$  is smooth, and  $X \cap G$  is smooth, so it follows that  $X_e$  is regular, and hence smooth. It is geometrically connected, because it is fully ramified over  $X$  at any point of  $X \cap G$ . We have already seen in the first part of section 4 that  $X_e \cap Z$  and  $X_e \cap Z \cap F$  are both smooth of the correct dimension. So the purity of

$$H_c^n(V_e \otimes_k \bar{k}, \mathcal{L}_{\psi(f)})$$

now results from [Ka-SE, 5.1.1(2)].

To conclude the proof of part (2a) of Theorem 3.1, it remains to compute the dimension of

$$H_c^n(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)}).$$

Since this is the only nonvanishing cohomology group, its dimension is equal to  $(-1)^n \times$  the Euler characteristic

$$\chi_c(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)}).$$

By standard arguments of reducing the  $L$  function mod various primes  $\lambda$  of  $\mathbb{Z}[\zeta_p, \zeta_{\#k-1}]$  of residue characteristic  $\neq p$  which divide the order of  $\chi$  and considering the degree of the resulting mod  $\lambda$   $L$ -function, we see that this Euler characteristic is independent of the particular choice of  $\chi$ , and is the same with  $\chi$  replaced by the trivial character:

$$\chi_c(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)}) = \chi_c(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)}).$$

On the other hand, we have

$$\begin{aligned} & \chi_c(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)}) \\ &= \chi_c(V \otimes_k \bar{k}, \mathcal{L}_{\psi(f)}) - \chi_c(V \cap (g=0) \otimes_k \bar{k}, \mathcal{L}_{\psi(f)}). \end{aligned}$$

Now  $\chi_c(V \otimes_k \bar{k}, \mathcal{L}_{\psi(f)})$  is the additive character euler characteristic attached to the situation  $(X, L, F)$ , with  $f = F/Z_d$  on  $V = X[1/L]$ . Similarly,  $\chi_c(V \cap (g=0) \otimes_k \bar{k}, \mathcal{L}_{\psi(f)})$  is the additive character euler characteristic attached to the situation  $(X \cap G, L, F)$ , with  $f = F/Z_d$  on  $V \cap (g=0) = (X \cap G)[1/L]$ . So from [Ka-SE, 5.1.1 and Remarque on page 166], we have the formulas

$$\begin{aligned} \chi_c(V \otimes_k \bar{k}, \mathcal{L}_{\psi(f)}) &= \int_X \frac{c(X)}{(1+L)(1+dL)}, \\ \chi_c(V \cap (g=0) \otimes_k \bar{k}, \mathcal{L}_{\psi(f)}) &= \int_{X \cap G} \frac{c(X \cap G)}{(1+L)(1+dL)} \\ &= \int_X \frac{eLc(X)}{(1+L)(1+dL)(1+eL)}. \end{aligned}$$

Subtracting, we find

$$\begin{aligned} & \chi_c(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)}) \\ &= \int_X \frac{c(X)}{(1+L)(1+dL)(1+eL)} := (-1)^n C(X, d, e), \end{aligned}$$

as required.

It remains to prove part (2b) of Theorem 3.1, that the “forget supports” map is an isomorphism

$$H_c^n(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)}) \cong H^n(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)}).$$

The right hand group  $H^n(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)})$  is, up to a Tate twist, the Poincaré dual of  $H_c^n(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\bar{\psi}(f)} \otimes \mathcal{L}_{\bar{\chi}(g)})$ , so (by part (2a) of Theorem 3.1, applied with  $\bar{\psi}$  and  $\bar{\chi}$ ) it has the same dimension,  $(-1)^n C(X, d, e)$ , as the left hand group. Therefore it suffices to show that the “forget supports” map is injective:

$$H_c^n(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)}) \hookrightarrow H^n(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)}).$$

For this we first reduce to the case when  $\chi^e$  is trivial, by passing to the covering  $V_r$  and looking at the commutative diagram

$$\begin{array}{ccc} H_c^n(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)}) & \xrightarrow{\text{forget}} & H^n(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)}) \\ \cap & & \cap \\ H_c^n(V_r[1/g_r] \otimes_k \bar{k}, \mathcal{L}_{\psi(f_r)} \otimes \mathcal{L}_{\chi(g_r)}) & \xrightarrow{\text{forget}} & H^n(V_r[1/g_r] \otimes_k \bar{k}, \mathcal{L}_{\psi(f_r)} \otimes \mathcal{L}_{\chi(g_r)}). \end{array}$$

So it suffices to treat the case when  $\chi^e$  is trivial. In this case, we pass to the covering  $V_e$ , and look at the commutative diagram

$$\begin{array}{ccc} H_c^n(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)}) & \xrightarrow{\text{forget}} & H^n(V[1/g] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \mathcal{L}_{\chi(g)}) \\ \cap & & \cap \\ H_c^n(V_e \otimes_k \bar{k}, \mathcal{L}_{\psi(f)}) & \xrightarrow{\text{forget}} & H^n(V_e \otimes_k \bar{k}, \mathcal{L}_{\psi(f)}). \end{array}$$

This bottommost “forget supports” map is in fact bijective, by [Ka-SE, 5.1.1, part (0)].

## 6. APPENDIX: THE CASE OF $r \geq 1$ $g$ 'S

We begin by stating the generalization of Theorem 1.2 analogous to Theorem 2.1. As in that theorem,  $X/k$  is a projective, smooth, and geometrically connected  $k$ -scheme of dimension  $n \geq 1$ , given with a projective embedding  $X \hookrightarrow \mathbb{P}_k^N := \mathbb{P}$ . We fix integers  $d \geq 1$  and  $e_1, \dots, e_r \geq 1$ , all prime to  $p$ . We are given a linear form

$$Z \in H^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(1)),$$

a degree  $d$  form

$$F \in H^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(d)),$$

and, for  $i = 1, \dots, r$ , a degree  $e_i$  form

$$G_i \in H^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(e_i)),$$

all on the ambient projective space  $\mathbb{P}$ . We assume that the following transversality hypotheses hold.

- (1)  $X \cap Z$  is lisse of codimension 1 in  $X$ .
- (2)  $X \cap Z \cap F$  is lisse of codimension 1 in  $X \cap Z$  ( $:=$  empty, if  $n = 1$ ).

- (3) For any nonempty subset  $I \subset \{1, \dots, r\}$ ,  $X \cap Z \cap \bigcap_{i \in I} G_i$  is lisse of codimension  $\#I$  in  $X \cap Z$  ( $:=$  empty, if  $\#I \geq n$ ).
- (4) For any nonempty subset  $I \subset \{1, \dots, r\}$ ,  $X \cap Z \cap F \cap \bigcap_{i \in I} G_i$  is lisse of codimension  $1 + \#I$  in  $X \cap Z$  ( $:=$  empty, if  $1 + \#I \geq n$ ).

To this data, we attach the smooth affine  $k$ -scheme

$$V := X - X \cap Z = X[1/Z],$$

and the functions

$$f := F/Z^d : V \rightarrow \mathbb{A}_k^1$$

and

$$g_i := G_i/Z^{e_i} : V \rightarrow \mathbb{A}_k^1.$$

We denote by  $c(X)$  the total Chern class of  $X$ , and by  $L$  the class of  $\mathcal{O}_X(1)$ . We define the constant  $C(X, d, e_1, \dots, e_r)$  by

$$C(X, d, e_1, \dots, e_r) := (-1)^n \int_X \frac{c(X)}{(1+L)(1+dL) \prod_i (1+e_i L)}.$$

Thus when  $X$  is  $\mathbb{P}^n$  with the identity embedding of itself into  $\mathbb{P} = \mathbb{P}^n$ ,  $C(X, d, e_1, \dots, e_r)$  is the constant  $C(n, d, e_1, \dots, e_r)$  of Theorem 1.2.

We have the following generalization of Theorem 2.1.

**Theorem 6.1.** *Suppose that  $(X, Z, F, G_1, \dots, G_r)$  are as above. Then we have the following results.*

- (1) *We have the estimate*

$$\left| \sum_{x \in V(k)} \psi(f(x)) \prod_i \chi_i(g_i(x)) \right| \leq C(X, d, e_1, \dots, e_r) (\#k)^{n/2}.$$

*The associated  $L$  function is a polynomial  $P(T)$  (for  $n$  odd) or a reciprocal polynomial  $1/P(T)$  (for  $n$  even) of degree  $\leq C(X, d, e_1, \dots, e_r)$ , which is mixed of weight  $\leq n$ .*

- (2) *If  $P(T)$  has degree  $= C(X, d, e_1, \dots, e_r)$ , then  $P(T)$  is pure of weight  $n$ .*
- (3) *If, for any nonempty subset  $I \subset \{1, \dots, r\}$ ,  $X \cap \bigcap_{i \in I} G_i$  is lisse of codimension  $\#I$  in  $X$  ( $:=$  empty, if  $\#I > n$ ), then  $P(T)$  has degree  $= C(X, d, e_1, \dots, e_r)$ , and is pure of weight  $n$ .*

Exactly as in section 3, Theorem 6.1 follows from the following generalization of Theorem 3.1.

**Theorem 6.2.** *Suppose that  $(X, Z, F, G_1, \dots, G_r)$  are as in Theorem 6.1. Then we have the following results.*

- (1)  $H_c^i := H_c^i(V[1/\prod_i g_i] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes (\otimes_i \mathcal{L}_{\chi_i(g_i)}))$  vanishes for  $i \neq n$ .

- (2) If, for any nonempty subset  $I \subset \{1, \dots, r\}$ ,  $X \cap \bigcap_{i \in I} G_i$  is lisse of codimension  $\#I$  in  $X$  ( $:=$  empty, if  $\#I > n$ ), then we have the following results.
- (2a)  $H_c^n$  has dimension  $C(X, d, e_1, \dots, e_r)$ , and is pure of weight  $n$ .
- (2b) The “forget supports” map is an isomorphism

$$\begin{aligned} & H_c^n(V[1/\prod_i g_i] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes (\otimes_i \mathcal{L}_{\chi_i(g_i)})) \\ & \cong H^n(V[1/\prod_i g_i] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes (\otimes_i \mathcal{L}_{\chi_i(g_i)})). \end{aligned}$$

To prove the first part of Theorem 6.2, it suffices, exactly as in section 4, to prove the vanishing of  $H_c^i$  for  $i > n$ . We first reduce to the case when all  $\chi_i^{e_i}$  are trivial. Extending scalars, we can find a coordinate system  $(Z = Y_0, Y_1, \dots, Y_N)$  in the ambient  $\mathbb{P}$  which is transversal to  $X$ , to each  $X \cap \bigcap_{i \in I} G_i$  which is smooth, to  $X \cap Z$ ,  $X \cap Z \cap F$ , to every nonempty  $X \cap Z \cap \bigcap_{i \in I} G_i$ , and to every nonempty  $X \cap Z \cap F \cap \bigcap_{i \in I} G_i$ . So it suffices to treat the case when the original coordinate system  $(Z = X_0, X_1, \dots, X_N)$  has all these transversality properties. Then with  $q := \#k$ , we consider the “ $q - 1$ ’th power mapping”

$$[q - 1] : \mathbb{P} \rightarrow \mathbb{P}, (X_0, \dots, X_N) \mapsto (X_0^{q-1}, \dots, X_N^{q-1}).$$

It is finite and flat of degree  $(q - 1)^N$ , and finite etale over the dense open set where all  $X_i$  are invertible. Exactly as in section 4, it suffices to treat the pullback situation  $(X_{q-1}, Z, F_{q-1}, G_{1,q-1}, \dots, G_{r,q-1})$  by this map. This completes the reduction to the case when all  $\chi_i^{e_i}$  are trivial.

When all the  $\chi_i^{e_i}$  are trivial, we pass to the covering  $X_{e_1, \dots, e_r}$  of  $X$  defined by extracting, for each  $i = 1, \dots, r$ , the  $e_i$ ’th root of  $G_i$ . On this covering, we have the pullbacks  $Z$  and  $F$  of their namesakes on  $X$ . Exactly as in section 4, the “nonsingular” case ( $\epsilon = \delta = -1$ ) of [Ka-ESES, Thm. 4], applied now to the data  $(X_{e_1, \dots, e_r}, Z, F)$ , gives the vanishing of  $H_c^i$  for  $i > n$ .

To prove the second part of Theorem 6.2, we observe that under the additional transversality hypotheses, the covering  $X_{e_1, \dots, e_r}$  of the previous paragraph is itself smooth, so the purity of  $H_c^n$  again results from [Ka-SE, 5.1.1(2)]. Exactly as in section 5, the dimension of  $H_c^n$  is  $(-1)^n \times$  the Euler characteristic

$$\begin{aligned} & \chi_c(V[1/\prod_i g_i] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)} \otimes \otimes_i (\mathcal{L}_{\chi_i(g_i)})) \\ & = \chi_c(V[1/\prod_i g_i] \otimes_k \bar{k}, \mathcal{L}_{\psi(f)}). \end{aligned}$$



The asserted formula for this Euler formula now follows by inclusion-exclusion from the formulas of [Ka-SE, 5.1.1 and Remarque on page 166]. This proves part (2a). The proof of part (2b) is entirely analogous to the proof of part (2b) of Theorem 3.1 given in section 5.

## REFERENCES

- [A-K] Altman, A., and Kleiman, S., Introduction to Grothendieck duality theory. Lecture Notes in Mathematics, Vol. 146 Springer-Verlag, Berlin-New York 1970 ii+185 pp.
- [De-ST] Deligne, P., Applications de la formule des traces aux sommes trigonométriques, pp. 168-232 in SGA 4 1/2, cited below.
- [De-Weil I] Deligne, P., La conjecture de Weil. Publ. Math. IHES 43 (1974), 273-307.
- [De-Weil II] Deligne, P., La conjecture de Weil II. Publ. Math. IHES 52 (1981), 313-428.
- [Gr-Rat] Grothendieck, A., Formule de Lefschetz et rationalité des fonctions  $L$ . Séminaire Bourbaki, Vol. 9, Exp. No. 279, 41-55, Soc. Math. France, 1995.
- [Ka-ENSMCS] Katz, N., Estimates for nonsingular multiplicative character sums. Int. Math. Res. Not. 2002, no. 7, 333-349.
- [Ka-ESES] Katz, N., Estimates for “singular” exponential sums, IMRN 16, 1999, 875-899.
- [Ka-SCMD] Katz, N., A semicontinuity result for monodromy under degeneration. Forum Math. 15 (2003), no. 2, 191-200.
- [Ka-SE] Katz, N., *Sommes Exponentielles*, rédigé par G. Laumon, Astérisque 79, 1980.
- [Ray] Raynaud, M., Caractéristique d’Euler-Poincaré d’un faisceau et cohomologie des variétés abéliennes. Séminaire Bourbaki, Vol. 9, Exp. No. 286, 129-147, Soc. Math. France, Paris, 1995.
- [SGA 4 1/2] Cohomologie Etale. Séminaire de Géométrie Algébrique du Bois Marie SGA 4 1/2. par P. Deligne, avec la collaboration de J. F. Boutot, A. Grothendieck, L. Illusie, et J. L. Verdier. Lecture Notes in Mathematics, Vol. 569, Springer-Verlag, 1977.
- [SGA 4 Tome 3] Théorie des Topos et Cohomologie Etale des Schémas, Tome 3. Séminaire de Géométrie Algébrique du Bois Marie 1963-1964 (SGA 4). Dirigé par M. Artin, A. Grothendieck, J. L. Verdier. Lecture Notes in Mathematics, Vol.305, Springer-Verlag, 1973.

PRINCETON UNIVERSITY, MATHEMATICS, FINE HALL, NJ 08544-1000, USA  
*E-mail address:* nmk@math.princeton.edu