

# A NOTE ON GALOIS REPRESENTATIONS WITH BIG IMAGE

NICHOLAS M. KATZ

## CONTENTS

Introduction	1
1. The case $N \geq 3$	2
2. An alternate approach to the case $N \geq 3, N \neq 4$	6
3. Yet another variant, for $N \geq 3$	8
4. Working over $\mathbb{Q}$ , when $\ell \equiv 1 \pmod{3}$	12
5. Working over $\mathbb{Q}$ , when $\ell \equiv 1 \pmod{4}$	14
6. Independence of $\ell$	16
7. Examples in higher dimension	17
8. Explicit examples in higher dimension	19
9. Some open questions	22
References	22

## INTRODUCTION

Given an integer  $N \geq 3$ , we will first construct reasonably “motivic” representations

$$\rho : \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}(\zeta_N)) \rightarrow \text{GL}(n, \mathbb{Q}_\ell)$$

with open image, for any  $\ell$  which is 1 mod  $N$  and for certain  $n$ . We will do this in three different ways. The third of them has a descent to  $\mathbb{Q}$  when  $N$  is 3 or 4. This provides us with motivic galois representations of  $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  with open image in  $\text{GL}(n, \mathbb{Q}_\ell)$  for any even  $n \geq 6$  and any  $\ell$  with  $\ell \equiv 1 \pmod{3}$  or  $\pmod{4}$ .

The underlying idea in all the cases considered is this. First, we find families of (projective, smooth, geometrically connected) curves over open sets of  $\mathbb{P}^1$  over suitable rings of  $S$ -integers together with an action of a finite cyclic group  $G$  such that suitable  $G$ -isotypical components of the  $H^1$  along the fibres have complex monodromy groups which are Zariski dense in  $SL(n, \mathbb{C})$  (for  $n$  the rank of the isotypical

component in question). The Zariski density is proven by specializing into finite characteristic, where on the one hand old results of the author (especially [Ka-ESDE] and [Ka-ACT]) give such Zariski density results for the geometric monodromy groups in characteristic  $p$ , and where on the other hand a fundamental semicontinuity theorem of Pink, [Ka-ESDE, 8.18.2] or [Ka-ST, 2.1], tells us that knowing this Zariski density in all but finitely many characteristics forces its truth in characteristic zero. The second step is to apply a form of Hilbert irreducibility, along the lines of Serre [Se-MW, Theorem, page 149] or Terasoma [Ter, Thm. 2]. This shows the existences of infinitely many rational points in the parameter space such the isotypical component of the  $H^1$  at that rational point is a galois representation with the desired large image. We should point out that our method does **not** give explicit rational points with this property.

The descents from  $\mathbb{Q}(\zeta_N)$  to  $\mathbb{Q}$  in the cases  $N = 3$  and  $N = 4$  are in some sense tricks. They make explicit use of well-chosen elements of  $PGL(2, \mathbb{Q})$  of orders 3 and 4 respectively, namely  $W \mapsto \frac{1}{1-W}$  and  $W \mapsto \frac{W-1}{W+1}$ . We do not understand the general setting to which these tricks should belong.

The problem of constructing galois representations with large image has also been considered by Upton [Up] in the case  $GL_3$ , by Greenberg [Gre] for  $GL_n$ , and by Cornut-Ray [C-R] for more general groups. The construction of Upton is motivic, those of Greenberg and Cornut-Ray are spectacularly non-motivic. Motivic galois representations with images which are Zariski dense in exceptional groups have been constructed by Dettweiler-Reiter [D-R] (for  $G_2$ , by local system methods), by Yun [Yun] (for  $G_2, E_7, E_8$ , by automorphic methods), and by Boxer-Celegari-Emerton- Levin-Madapusi Pera-Patrikis [BCELMP] (for  $E_6$ , also by automorphic methods).

It is a pleasure to thank Peter Sarnak and Laurent Clozel, who awakened, respectively reawakened, my interest in this question.

## 1. THE CASE $N \geq 3$

In this section, we fix an integer  $N \geq 3$ , an integer  $n \geq 5$  such that  $n + 1$  is nonzero mod  $N$ , a prime number  $\ell$  which is 1 mod  $N$ , and an embedding of  $\mathbb{Q}(\zeta_N)$  into  $\mathbb{Q}_\ell$ . We also fix a monic polynomial

$$f = f_{n+1}(X) \in \mathbb{Z}[X]$$

which, over  $\mathbb{C}$ , is a **Morse polynomial**, meaning that the derivative  $f'(X)$  has  $n$  distinct zeroes, say  $\alpha_1, \dots, \alpha_n$ , and  $f$  separates these zeroes:

$$f(\alpha_i) \neq f(\alpha_j) \text{ if } i \neq j.$$

The values  $f(\alpha_i)$  are called the critical values of  $f$ . The critical values are integral over  $\mathbb{Z}[1/(n+1)]$ . We define a polynomial  $F_{crit.val.}(T) \in \mathbb{Z}[1/(n+1)][T]$  by

$$F_{crit.val.}(T) := \prod_{\text{roots } \alpha_i \text{ of } f'} (T - f(\alpha_i))$$

and denote by

$$\Delta_{Morse}(f) := \text{Discriminant}(F_{crit.val.}).$$

For any  $t \in \mathbb{C}$  which is not a critical value of  $f$ , the polynomial  $t - f(X)$  has all distinct roots. For any prime  $p$  which is prime to  $n+1$  and to the numerator of  $\Delta_{Morse}(f)$ , the coefficient-wise reduction mod  $p$  of  $f$  is a Morse polynomial in characteristic  $p$ .

For example, the polynomial  $X^{n+1} - (n+1)X$  is a Morse polynomial. The zeroes of its derivative are the  $n$ 'th roots of unity. Its critical values are  $\{-n\zeta\}_{\zeta \in \mu_n}$ , and  $F_{crit.val.}(T) = T^n - (-n)^n$ . Its reduction mod any prime  $p$  not dividing  $n(n+1)$  is a Morse polynomial in characteristic  $p$ .

Over the parameter space

$$S := \text{Spec}(\mathbb{Z}[\zeta_N, 1/N, 1/(n+1), 1/\Delta_{Morse}(f)][T][1/F_{crit.val.}(T)])$$

we have the one parameter family of curves  $\pi : \mathcal{C} \rightarrow S$  given by

$$\mathcal{C} : Y^N = T - f(X).$$

The group  $\mu_N := \mu_N(\mathbb{Z}[\zeta_N, 1/N])$  acts on this family, by  $(X, Y) \mapsto (X, \zeta Y)$ . The sheaf  $\mathcal{F}_\ell := R^1\pi_!\mathbb{Q}_\ell$  is lisse on  $S[1/\ell]$ , and carries the action of  $\mu_N$ . For a character  $\chi$  of  $\mu_N$  of full order  $N$ , the  $\chi$ -isotypical component  $\mathcal{F}_\ell^\chi$  of  $\mathcal{F}_\ell$  is lisse on  $S[1/\ell]$  of rank  $n$  and pure of weight one. [It is to insure the purity that we need  $n+1$  to be nonzero mod  $N$ , otherwise the rank of  $\mathcal{F}_\ell^\chi$  remains  $n$  but its quotient of (highest) weight one has rank  $n-1$ , cf. [Ka-ACT, 5.16 and 5.18]. ] We denote by

$$\rho_{\ell, f, \chi} : \pi_1(S[1/\ell]) \rightarrow GL(n, \mathbb{Q}_\ell)$$

the  $\ell$ -adic representation which "is"  $\mathcal{F}_\ell^\chi$ .

**Theorem 1.1.** *The image of  $\pi_1(S[1/\ell])$  in  $GL(n, \mathbb{Q}_\ell)$  is open. Moreover, if we embed  $\mathbb{Z}[\zeta_N]$  into  $\mathbb{C}$ , the image of*

$$\pi_1^{geom}(S[1/\ell]) := \pi_1(S \otimes_{\mathbb{Z}[\zeta_N]} \mathbb{C})$$

*contains an open subgroup of  $SL(n, \mathbb{Q}_\ell)$ .*

*Proof.* It suffices to prove the "moreover", as the determinant, being pure of weight  $n \neq 0$ , certainly has open image in  $\mathbb{Z}_\ell^\times$ . Let us denote

by  $\Gamma$  the image of  $\pi_1^{geom}(S[1/\ell])$  in  $GL(n, \mathbb{Q}_\ell)$ . Fix a prime  $p$  not dividing either  $n + 1$  or the numerator of  $\Delta_{Morse}(f)$ , and a  $p$ -adic place of  $\mathbb{Z}[\zeta_N, 1/N]$  with residue field some  $\mathbb{F}_q$  with  $q \equiv 1 \pmod{N}$ . We denote by  $\Gamma(p)$  the image of  $\pi_1^{geom}(S[1/\ell] \otimes \mathbb{F}_q)$  in  $GL(n, \mathbb{Q}_\ell)$ . By Pink's specialization theorem, cf. [Ka-ESDE, 8.18.2] or [Ka-ST, 2.1],  $\Gamma$  contains (a conjugate of)  $\Gamma(p)$ . So it suffices to show that  $\Gamma(p)$  contains an open subgroup of  $SL(n, \mathbb{Q}_\ell)$ . For this, we argue as follows. It is proven in [Ka-ACT, 5.13] that the Zariski closure  $G_{geom,p}$  of  $\Gamma(p)$  sits in

$$SL(n) \subset G_{geom,p} \subset GL(n)$$

and has determinant of finite order. Let us denote by

$$\Gamma(p)_1 \subset \Gamma(p)$$

the subgroup of elements of determinant one. This is a subgroup of finite index, so its Zariski closure contains the identity component  $G_{geom,p}^0 = SL(n)$ . Hence  $\Gamma(p)_1$  is Zariski dense in  $SL(n)$ . We then conclude by the following well-known lemma, cf [Se-pDiv, Cor, page 120], which we record for ease of reference.

**Lemma 1.2.** *Over  $\mathbb{Q}_\ell$ , let  $G \subset SL(n)$  be an irreducible connected (and hence semisimple) algebraic group. Suppose  $\Gamma \subset G(\mathbb{Q}_\ell)$  is a closed subgroup which is Zariski dense in  $G$ . Then  $\Gamma$  contains an open subgroup of  $G(\mathbb{Q}_\ell)$ .*

*Proof.* The group  $\Gamma$  is an  $\ell$ -adic Lie group. Every open subgroup  $\Gamma_1$  of  $\Gamma$  is also Zariski dense in  $G$  (because  $G$  is connected, and replacing  $\Gamma$  by  $\Gamma_1$  does not change the identity component of the Zariski closure). Its Lie algebra  $\text{Lie}(\Gamma) = \text{Lie}(\Gamma_1)$  is reductive, because  $\Gamma_1$  has, via  $G$ , an irreducible representation. Because this irreducible representation has trivial determinant,  $\text{Lie}(\Gamma)$  has no nonzero abelian factor. Thus  $\text{Lie}(\Gamma)$  is semisimple. One knows that a semisimple Lie algebra is algebraic. Visibly we have the inclusion  $\text{Lie}(\Gamma) \subset \text{Lie}(G)$ . Thus  $\text{Lie}(\Gamma)$  is  $\text{Lie}(H)$  for some connected semisimple group  $H \subset G$ , and for a small enough  $\Gamma_1$ ,  $\Gamma_1$  is open in  $H(\mathbb{Q}_\ell)$  (because they have the same Lie algebra) and hence is Zariski dense in  $H$ . Therefore  $H = G$ .  $\square$

This concludes the proof of the theorem.  $\square$

**Remark 1.3.** In fact, for any  $\ell$  which is 1 mod  $N$ , the image of  $\rho_{\ell,f,X}$  contains  $SL(n, \mathbb{Z}_\ell)$  (via some  $\mathbb{Z}_\ell$ -lattice in  $\mathcal{F}_{\ell,f,X}$ ). [See [H-L] for another approach to this sort of question.] To see this, we use the (never written!) theory of middle convolution with mod  $\ell$  coefficients, applied in any good characteristic  $p$ . The sheaf  $\mathcal{F}_{\ell,f,X}$  is the middle additive convolution

$$\mathcal{L}_X \star_{mid,+} (f_* \mathbb{Q}_\ell / \mathbb{Q}_\ell).$$

Its reduction mod  $\ell$  is the middle additive convolution

$$\mathcal{L}_\chi \star_{mid,+} (f_\star \mathbb{F}_\ell / \mathbb{F}_\ell).$$

Because  $f$  is Morse, the second factor is irreducible, tame at  $\infty$ , and all of its local monodromies at finite distance are reflections. Therefore the middle convolution is irreducible, tame at  $\infty$ , and all of its local monodromies at finite distance are pseudoreflections with determinant  $\chi \chi_{quad}$ , which is always of order  $\geq 3$  because  $N \geq 3$ . Thus the mod  $\ell$  image of  $\Gamma(p)$  is an irreducible subgroup of  $GL(n, \mathbb{F}_\ell)$  which is generated by pseudoreflections of order  $\geq 3$ . By a theorem of Wagner [Wag, Thm. 1.2 and following paragraph], an irreducible subgroup of  $GL(n, \mathbb{F}_\ell)$  generated by pseudoreflections of order  $\geq 3$  necessarily contains  $SL(n, \mathbb{F}_\ell)$  provided that  $n \geq 5$ . The inverse image, call it  $\Gamma(p)_1$ , in  $\Gamma(p)$  of  $SL(n, \mathbb{F}_\ell)$  lies in  $SL(n, \mathbb{Z}_\ell)$  (because the determinants of elements of  $\Gamma(p)$  are roots of unity of order dividing  $\ell - 1$ ). Then  $\Gamma(p)_1$  is a closed subgroup of  $SL(n, \mathbb{Z}_\ell)$  which maps onto  $SL(n, \mathbb{F}_\ell)$ . One knows [Se-AbLRep, Exc. 1, page IV-27] that if  $n \geq 2$  and  $\ell \geq 5$ , the only closed subgroup of  $SL(n, \mathbb{Z}_\ell)$  which maps onto  $SL(n, \mathbb{F}_\ell)$  is  $SL(n, \mathbb{Z}_\ell)$  itself. Thus  $\Gamma(p)_1$  is  $SL(n, \mathbb{Z}_\ell)$ . Hence  $\Gamma(p)$  contains  $SL(n, \mathbb{Z}_\ell)$ . As  $\Gamma$  contains  $\Gamma(p)$ , we are done.

Here is another approach to this question, which gives the result for all but a finite set of  $\ell$  and does not use the theory of mod  $\ell$  middle convolution. Over the Riemann surface  $S^{an} := (S \otimes_{\mathbb{Z}[\zeta_N]} \mathbb{C})^{an}$ , with

$$K := \mathbb{Q}(\zeta_n)$$

we have the  $K$ -local system

$$\mathcal{F}_{K,f} := R(\pi^{an})!K$$

and its  $\chi$  component  $\mathcal{F}_{K,f,\chi}$ . We know that each of its local monodromies at finite distance is a pseudoreflection of known order  $\geq 3$ , and that (correctly chosen conjugates of each of) these elements generate the entire image of  $\pi_1(S^{an})$  in  $GL(n, K)$ . So if we invert some highly divisible integer  $M$ , we can find an  $\mathcal{O}_K[1/M]$ -lattice  $\mathcal{F}_{\mathcal{O}_K[1/M],f,\chi}$  in  $\mathcal{F}_{K,f,\chi}$  whose monodromy representation lands in  $GL(n, \mathcal{O}_K[1/M])$ . Because our  $K$ -representation was absolutely irreducible, the entire matrix ring  $Mat_n(K)$  is spanned over  $K$  by the the image of the  $K$ -grouping of  $\pi_1(S^{an})$ . At the expense of increasing  $M$ , we may assume that  $Mat_n(\mathcal{O}_K[1/M])$  is spanned by the image of the  $\mathcal{O}_K[1/M]$ -grouping. Once we are in this situation, then the reduction mod any prime  $\mathcal{P}$  of  $\mathcal{O}_K[1/M]$  of this  $\mathcal{O}_K[1/M]$ -form of our monodromy representation will be absolutely irreducible. Take a  $\mathcal{P}$  whose residue field is a prime field  $\mathbb{F}_\ell$  and whose completion is  $\mathbb{Q}_\ell$ . We apply Wagner's theorem to

conclude that the monodromy representation of  $\mathcal{F}_{\mathcal{O}_K[1/M],f,\chi} \otimes \mathbb{Z}_\ell$  on  $S^{an}$  has image containing  $SL(n, \mathbb{Z}_\ell)$ . On the scheme

$$S_{\mathbb{C}} := S \otimes_{\mathbb{Z}[\zeta_N]} \mathbb{C},$$

we have our  $\mathbb{Q}_\ell$  local system  $\mathcal{F}_\ell^\chi$ , which is the “partner” (in the sense of [SGA4, XI, 4.4] via [SGA4, XVI, 4.1]) of the transcendental local system  $\mathcal{F}_{\mathcal{O}_K[1/M],f,\chi} \otimes \mathbb{Q}_\ell$  on  $S^{an}$ . So the image of the monodromy representation of  $\mathcal{F}_\ell^\chi$  on the scheme  $S_{\mathbb{C}}$  contains  $SL(n, \mathbb{Z}_\ell)$ .

**Corollary 1.4.** *There exist infinitely many  $t \in \mathbb{Q}(\zeta_N)$  with  $F_{crit.val.}(t) \neq 0$  for which the representation of  $Gal(\overline{\mathbb{Q}}/\mathbb{Q}(\zeta_N))$  given by specializing  $T$  to  $t$  (i.e. by composing the maps*

$$Gal(\overline{\mathbb{Q}}/\mathbb{Q}(\zeta_N)) = \pi_1(Spec(\mathbb{Q}(\zeta_N))) \rightarrow \pi_1(S \otimes_{\mathbb{Z}} \mathbb{Q}) \rightarrow \pi_1(S[1/\ell]) \rightarrow GL(n, \mathbb{Q}_\ell),$$

*the last map being the representation  $\rho_{\ell,f,\chi}$ ) has open image, indeed the same open image as the representation  $\rho_{\ell,f,\chi}$ .*

*Proof.* This is Hilbert Irreducibility, cf.[Ter, Thm.2] or [Se-MW, Theorem, page 149].  $\square$

**Remark 1.5.** When we specialize  $T$  to  $t \in \mathbb{Q}(\zeta_N)$ , the  $\ell$ -adic representation we get is the  $\chi$ -component of

$$H_c^1(\mathcal{C}_t \otimes \overline{\mathbb{Q}}, \mathbb{Q}_\ell)$$

for  $\mathcal{C}_t$  the curve

$$\mathcal{C}_t : Y^N = t - f(X)$$

over  $\mathbb{Q}(\zeta_N)$ .

## 2. AN ALTERNATE APPROACH TO THE CASE $N \geq 3, N \neq 4$

Fix an integer  $n \geq 1$  such that  $n + 1$  is nonzero mod  $N$ . Fix  $f = f_n(X) \in \mathbb{Z}[X]$  a monic polynomial of degree  $n$  whose discriminant  $\Delta(f_n)$  is nonzero. [For example, we might take  $\prod_{i=1}^n (X - i)$ , whose discriminant is nonzero mod every prime  $p \geq n$ .] Over the parameter space

$$S := Spec(\mathbb{Z}[\zeta_N, 1/N][T][1/f_n(T)][1/\Delta(f_n)])$$

we have the family of affine curves

$$\pi : \mathcal{C} \rightarrow S$$

of equation

$$\mathcal{C} : Y^N = f_n(X)(T - X).$$

The group  $\mu_N(\mathbb{Q}(\zeta_N))$  has an obvious action on this family, namely  $(x, y) \mapsto (x, \zeta y)$ . For any  $\ell$ , the relevant  $H^1$  along the fibres,

$$\mathcal{F}_\ell := R^1 \pi_!(\mathbb{Q}_\ell),$$

is lisse on  $S[1/\ell]$ , and carries an action of  $\mu_N(\mathbb{Q}(\zeta_N))$ .

Suppose now that  $\ell$  is 1 mod  $N$ . Then we can choose an embedding of  $\mathbb{Q}(\zeta_N)$  into  $\mathbb{Q}_\ell$ , and diagonalize the action. For a character  $\chi$  of  $\mu_N(\mathbb{Q}(\zeta_N))$  of full order  $N$ , the  $\chi$  isotypical component  $\mathcal{F}_\ell^\chi$  of  $\mathcal{F}_\ell$  is lisse of rank  $n$  and pure of weight one.

**Theorem 2.1.** *Fix an integer  $n \geq 5$  such that  $n + 1$  is nonzero mod  $N$ . Suppose that  $\ell$  is 1 mod  $N$ . Let  $\chi$  be a character of order  $N$ . Then the local system  $\mathcal{F}_\ell^\chi$  on  $S[1/\ell]$ , viewed as a representation*

$$\rho_\chi : \pi_1(S[1/\ell]) \rightarrow GL(n, \mathbb{Q}_\ell)$$

*has open image. Moreover, after extension of scalars from  $\mathbb{Q}(\zeta_N)$  to  $\mathbb{C}$  (e.g. by mapping  $\zeta_N$  to  $\exp(2\pi i/N)$ ), the image of  $\pi_1^{geom}(S[1/\ell]) := \pi_1(S \otimes_{\mathbb{Z}[\zeta_N, 1/N]} \mathbb{C})$  contains an open subgroup of  $SL(N, \mathbb{Q}_\ell)$ , and for all but at most finitely many  $\ell$  the image contains  $SL(n, \mathbb{Z}_\ell)$ .*

*Proof.* The overall structure of the proof is the same as that of Theorem 1.1. It suffices to prove the “moreover” statement, and for this it is enough to specialize to a good characteristic  $p \neq \ell$  (one not dividing  $\Delta(f_n)$ ) and a residue field  $\mathbb{F}_q$  at a  $p$ -adic place of  $\mathbb{Q}(\zeta_N)$ . Then our local system, pulled back to  $\text{Spec}(\mathbb{F}_q[T][1/f_n(T)])$ , is tame at  $\infty$  (as it comes from characteristic zero, cf. [Ka-SE, 4.7.1]). Its trace function is as follows: for  $k/\mathbb{F}_q$  a finite extension, and  $\chi$  extended to  $k^\times$  by composition with the norm,

$$\text{Trace}(\text{Frob}_{t,k} | \mathcal{F}_\ell^\chi) = - \sum_{x \in k} \chi(f_n(x)(t - x)).$$

This is the trace function of the middle additive convolution [Ka-RLS]

$$\mathcal{L}_\chi \star_{mid,+} \mathcal{L}_{\chi(f_n(x))}.$$

Thus our local system is geometrically irreducible [Ka-RLS, 2.9.6 and 2.9.8], and its local monodromies at finite distance (i.e., at the zeroes of  $f_n(t)$ ) are all pseudoreflections of determinant  $\chi^2$ , cf. [Ka-RLS, 3.3.6]. Because  $N \geq 3$  but  $N \neq 4$ ,  $\chi^2$  has order  $\geq 3$ . It then follows from [Ka-ACT, 5.11] that in our chosen characteristic  $p$ , the image  $\Gamma(p)$  of  $\pi_1^{geom}(S \otimes \mathbb{F}_q)$  has Zariski closure  $G_{geom,p}$  with  $G_{geom,p}^0 = SL(n)$ . Exactly as in the proof of Theorem 1.1, we infer that  $\Gamma(p)$  contains an open subgroup of  $SL(n, \mathbb{Q}_\ell)$ . To show that, for all but at most finitely many  $\ell$ , it contains all of  $SL(n, \mathbb{Z}_\ell)$  for a suitable  $\mathbb{Z}_\ell$ -lattice in  $\mathcal{F}_\ell^\chi$ , we repeat that Wagner argument of Remark 1.3.  $\square$

**Remark 2.2.** On  $S_{\mathbb{C}}^{an}$ , the  $\chi$  component of the local system  $\mathcal{F}_{\mathbb{C}} := R^1\pi_1^{an}\mathbb{C}$  is the sheaf of germs of local holomorphic solutions of a Pochhammer hypergeometric equation, cf. [De-Mos, bottom of page 6] and [Poch, pp. 322-325].

**Remark 2.3.** Exactly as in the previous section, Hilbert Irreducibility ensures that there are infinitely many  $t \in \mathbb{Q}(\zeta_N)$  with  $f_n(t) \neq 0$  for which the  $\ell$ -adic representation given by the  $\chi$  component of the  $H_c^1$  of the curve

$$Y^N = f_n(X)(t - X)$$

over  $\mathbb{Q}(\zeta_N)$  has open image in  $GL(n, \mathbb{Q}_{\ell})$ .

### 3. YET ANOTHER VARIANT, FOR $N \geq 3$

In this section, we fix an integer  $N \geq 3$ , a nonzero element  $A \in \mathbb{Z}[\zeta_N]$ , an integer  $n \geq 2$ , a prime number  $\ell$  which is 1 mod  $N$ , an embedding of  $\mathbb{Q}(\zeta_N)$  into  $\mathbb{Q}_{\ell}$ , and a monic polynomial  $f = f_n(X) \in \mathbb{Z}[\zeta_N][X]$  which, over  $\mathbb{C}$  (by any choice of embedding) is a Morse polynomial. For any  $t \in \mathbb{C}$  which is not a critical value of either  $f$  or of  $f + A$ , the rational function

$$\frac{t - f(X) - A}{t - f(X)}$$

has  $n$  simple zeroes and  $n$  simple poles at finite distance, no other zeroes or poles, and takes the value 1 at  $\infty$ . We denote by

$$F := F_{crit.val.}(T)$$

the monic polynomial whose roots are the  $n - 1$  distinct critical values of  $f$ . We make the additional hypothesis

(CritDiff<sub>A</sub>) The polynomial  $F(T)F(T - A)$  has  $2n - 2$  i.e., all distinct roots.

Equivalently, the condition on  $f$  is that  $A$  is not the difference of two distinct critical values of  $f$ . We denote by  $\Delta_{crit,A} \in \mathbb{Z}[\zeta_N, 1/(2n)]$  the discriminant of  $F(T)F(T - A)$ .

Notice that if  $f$  is a Morse polynomial satisfying (CritDiff<sub>A</sub>), then for any constant  $a \in \mathbb{Z}[\zeta_N]$ , the polynomial  $f(X) + a$  is also a Morse polynomial satisfying (CritDiff<sub>A</sub>).

Here is an example, to show that for any  $A \neq 0$ , we can find a Morse  $f = f_n$  which satisfies (CritDiff<sub>A</sub>). For an integer  $M \neq 0$ , consider the polynomial

$$X^n - nM^{n-1}X.$$

Its critical values are  $\{M^n(1 - n)\zeta\}_{\zeta \in \mu_{n-1}}$ . Using the archimedean inequality

$$|e^{i\alpha} - 1| \geq 2\alpha/\pi \quad \text{for } 0 \leq \alpha \leq \pi,$$



we see that for every complex embedding of  $\mathbb{Q}(\zeta_{n-1})$ , and any two distinct  $n - 1$  roots of unity  $\zeta$  and  $\zeta'$ , we have

$$|(n - 1)(\zeta - \zeta')| \geq 4.$$

So taking  $M$  large enough that  $4M^n$  exceeds every archimedean absolute value of  $A$  provides a suitable  $f_n$ .

More generally, if  $f = f_n$  is a Morse polynomial, then for any integer  $M$ , the polynomial  $M^n f(X/M)$  has critical values  $M^n$  times those of  $f$  (and critical points  $M$  times those of  $f$ ). So if we take  $M$  sufficiently large,  $M^n f(X/M)$  will satisfy  $(\text{CritDiff}_A)$ .

Fix now a Morse function  $f$  and an  $A \neq 0$  in  $\mathbb{Z}[\zeta_N]$ , such that  $(\text{CritDiff}_A)$  holds for  $f$ . Over the parameter space

$$S := \text{Spec}(\mathbb{Z}[\zeta_N, 1/N, 1/(2n), 1/A, 1/\Delta_{\text{crit},A}(T)][T][1/(F(T)F(T - A))])$$

we have the complete nonsingular model (add  $N$  sections along  $\infty$ )

$$\pi : \bar{\mathcal{C}} \rightarrow S$$

of the family of smooth affine curves

$$\mathcal{C} : y^N = \frac{T - f(X) - A}{T - f(X)}.$$

The sheaf  $\mathcal{F}_\ell := R^1\pi_1\mathbb{Q}_\ell$  is lisse on  $S[1/\ell]$  of rank  $(N - 1)(2n - 2)$  and pure of weight one. For any character  $\chi$  of  $\mu_N$  of full order  $N$ , the  $\chi$  component  $\mathcal{F}_\ell^\chi$  is lisse of rank  $2n - 2$  and pure of weight one. [In fact, this is true for every nontrivial  $\chi$  of order dividing  $N$ .]

We denote by

$$\rho_{\ell,f,A,\chi} : \pi_1(S[1/\ell]) \rightarrow GL(2n - 2, \mathbb{Q}_\ell)$$

the  $\ell$ -adic representation which “is”  $\mathcal{F}_\ell^\chi$ .

**Theorem 3.1.** *Suppose we are in one of the two following situations.*

- (1)  $n \geq 4$ ,  $f$  satisfies  $(\text{CritDiff}_A)$  and is of the form  $X^n - M^{n-1}nX + a$  for some  $a \in \mathbb{Z}[\zeta_N]$ .
- (2)  $n \geq 6$ ,  $f$  satisfies  $(\text{CritDiff}_A)$  and is of the form  $M^n h(X/M) + a$  for some  $a \in \mathbb{Z}[\zeta_N]$  and some monic  $h = h_n \in \mathbb{Z}[X]$  whose derivative  $h'$  has Galois group over  $\mathbb{Q}$  the full symmetric group  $S_{n-1}$ .

*Then the image of  $\pi_1(S[1/\ell])$  in  $GL(2n - 2, \mathbb{Q}_\ell)$  is open. Moreover, if we embed  $\mathbb{Z}[\zeta_N]$  into  $\mathbb{C}$ , the image of*

$$\pi_1^{\text{geom}}(S[1/\ell]) := \pi_1(S \otimes_{\mathbb{Z}[\zeta_N]} \mathbb{C})$$

*contains an open subgroup of  $SL(2n - 2, \mathbb{Q}_\ell)$ .*

*Proof.* Exactly as in the proof of Theorem 1.1, it suffices to prove the “moreover”. And for this it suffices to pass to an  $\mathbb{F}_q$ -valued point of  $\mathbb{Z}[\zeta_N, 1/N, 1/(2n), 1/\Delta_{crit,A}, 1/\ell]$ , and show that over  $S \otimes \mathbb{F}_q$  the image of  $\pi_1^{geom}(S \otimes \mathbb{F}_q)$  contains an open subgroup of  $SL(2n-2, \mathbb{Q}_\ell)$ . We will choose such a point whose characteristic  $p$  is sufficiently large that, in case (1), [Ka-ESDE, 7.10.5] applies, and in case (2) that [Ka-ESDE, 7.10.6] applies. What this choice of  $p$  ensures is that in our characteristic  $p$  situation, when we consider the Fourier Transform

$$FT(f_*\mathbb{Q}_\ell/\mathbb{Q}_\ell)$$

its geometric monodromy group  $G_{geom}$  sits in

$$Sp(n-1) \subset G_{geom} \subset \mu_p Sp(n-1), \text{ in case [1], } n \text{ odd,}$$

$$SL(n-1) \subset G_{geom} \subset \pm 1 \mu_p SL(n-1), \text{ in case [1], } n \text{ even,}$$

$$SL(n-1) \subset G_{geom} \subset \pm 1 \mu_p SL(n-1), \text{ in case [2], any } n.$$

[The references cited apply to an  $f$  to which a suitable constant has been added. The effect of such an addition is to perform an additive translation on  $f_*\mathbb{Q}_\ell/\mathbb{Q}_\ell$ , which in turn has the effect of tensoring its Fourier Transform by a character of order  $p$ , hence the  $\mu_p$  factor in the above statement. ]

Using Lemma 1.2, it suffices to show that  $G_{geom}^0 = SL(2n-2)$  in this characteristic  $p$  situation.

Henceforth we work in characteristic  $p$ . The trace function of our sheaf  $\mathcal{F}_\ell^x$  at points  $t$  of a finite field extension  $k/\mathbb{F}_q$  with  $F(t)F(t-A) \neq 0$  is the character sum

$$-1 - \sum_{x \in k} (\chi \circ \text{Norm}_{k/\mathbb{F}_q}) \left( \frac{t - f(x) - A}{t - f(x)} \right),$$

with the usual convention that  $\chi(0) = \chi(\infty) = 0$ .

Write

$$\frac{t - f(x) - A}{t - f(x)} = 1 - \frac{A}{t - f(x)}.$$

Then we see that this character sum is the additive convolution

$$- \sum_{u \in k} (\chi \circ \text{Norm}_{k/\mathbb{F}_q}) (1 - A/(t-u)) (\#\{x \in k \mid f(x) = u\} - 1).$$

of the trace functions of  $\mathcal{L}_\chi(1 - A/x)$  and  $f_*\mathbb{Q}_\ell/\mathbb{Q}_\ell$ . In other words,  $\mathcal{F}_\ell^x$  has the same trace function on  $S \otimes \mathbb{F}_q$  as the ! additive convolution

$$\mathcal{L}_\chi(1 - A/x) \star_{!,+} f_*\mathbb{Q}_\ell/\mathbb{Q}_\ell.$$

Let us admit for a moment the following key lemma.

**Lemma 3.2.** *The canonical surjection of perverse sheaves is an isomorphism*

$$\mathcal{L}_\chi(1 - A/x) \star_{!,+} f_\star \mathbb{Q}_\ell/\mathbb{Q}_\ell \cong \mathcal{L}_\chi(1 - A/x) \star_{mid,+} f_\star \mathbb{Q}_\ell/\mathbb{Q}_\ell.$$

*This middle convolution is geometrically irreducible, and all its monodromies at finite distance are pseudoreflections of determinant  $\chi\chi_{quad}$ .*

Then  $\mathcal{F}_\ell^\chi$  has the same trace function on  $S \otimes \mathbb{F}_q$  as the geometrically, and hence arithmetically, irreducible perverse middle convolution above. By Chebotarev, it follows that  $\mathcal{F}_\ell^\chi$  is isomorphic to this middle convolution. Thus  $\mathcal{F}_\ell^\chi$  is lisse on  $S \otimes \mathbb{F}_q$  of rank  $2n - 2 \geq 6$ , geometrically irreducible, with all finite monodromies pseudoreflections of order  $\geq 3$ . That  $\mathcal{F}_\ell^\chi$  is tame at  $\infty$  results from its “coming from characteristic zero”, cf. [Ka-SE, 4.7.1 (i)]. The result then follows from [Ka-ACT, 5.11].  $\square$

It remains to give the proof of the key lemma.

*Proof.* To show that the ! convolution is equal to the middle convolution, it is equivalent to show that its Fourier Transform FT is a middle extension, and that its Fourier Transform is geometrically irreducible. Now FT turns ! convolution into  $\otimes$ , so the FT of the ! convolution is

$$FT(\mathcal{L}_\chi(1 - A/x) \otimes FT(f_\star \mathbb{Q}_\ell/\mathbb{Q}_\ell)).$$

Both factors are lisse on  $\mathbb{G}_m$ . At 0, the first factor has unipotent local monodromy  $Unip(2)$  and the second factor has local monodromy  $\oplus_{\chi:\chi^n=1, \chi \neq 1} \mathcal{L}_\chi$ , so their tensor product correctly vanishes at 0.

It remains to show that this tensor product is geometrically irreducible. The first factor, being pure, has  $G_{geom}$  a (not necessarily connected) semisimple subgroup of  $GL(2)$ . So its identity component is either trivial or  $SL(2)$ . It is not trivial, because it contains  $Unip(2)$ .

The second factor has  $G_{geom}^0$  either  $SL(n - 1)$  or, if  $n$  is odd, possibly  $Sp(n - 1)$ . As  $n \geq 4$  in all cases, its Lie algebra is a simple Lie algebra which is not that of  $SL(2)$ . So by the Goursat-Kolchin-Ribet theorem [Ka-ESDE, 1.8.2], the direct sum object

$$FT(\mathcal{L}_\chi(1 - A/x) \oplus FT(f_\star \mathbb{Q}_\ell/\mathbb{Q}_\ell))$$

has

$$G_{geom}^0 = SL(2) \times (\text{either } SL(n - 1) \text{ or, if } n \text{ is odd, possibly } Sp(n - 1)).$$

Consequently, the tensor product is irreducible.

That its local monodromies at finite distance are pseudoreflections of determinant  $\chi\chi_{quad}$  is proven in [Ka-MMP, 6.1.18].

This concludes the proof of the key lemma, and with it the proof of the theorem.  $\square$

**Remark 3.3.** Exactly as in the previous sections, Hilbert Irreducibility ensures that there are infinitely many  $t \in \mathbb{Q}(\zeta_N)$  with  $\Delta_{crit,A}(t) \neq 0$  for which the  $\ell$ -adic representation given by the  $\chi$  component of the  $H_c^1$  of the complete nonsingular model (add  $\mu_N$  at  $\infty$ ) of the curve

$$y^N = \frac{t - f(X) - A}{t - f(X)}$$

over  $\mathbb{Q}(\zeta_N)$  has open image in  $GL(2n - 2, \mathbb{Q}_\ell)$ .

#### 4. WORKING OVER $\mathbb{Q}$ , WHEN $\ell \equiv 1 \pmod{3}$

Consider the automorphism of the rational function field  $\mathbb{Q}(W)$  given by the fractional linear transformation

$$\sigma : W \mapsto \frac{1}{1 - W}.$$

One checks easily that  $\sigma^3$  is the identity. We define the trace rational function

$$S(W) := W + \sigma(W) + \sigma^2(W) = W + \frac{1}{1 - W} + 1 - \frac{1}{W} = \frac{W^3 - 3W + 1}{W(W - 1)},$$

which is, of course,  $\sigma$ -invariant.

On the other hand, after we extend scalars to  $\mathbb{Q}(\zeta)(W)$ , for  $\zeta = \zeta_3$  a primitive cube root of unity, the rational function

$$R(W) := \frac{W + \zeta^2}{W + \zeta}$$

is easily checked to satisfy

$$\sigma(R(W)) = \zeta R(W),$$

and hence  $R(W)^3$  is  $\sigma$ -invariant.

We have the following miraculous identity, whose verification is left to the reader.

$$R(W)^3 := \left(\frac{W + \zeta^2}{W + \zeta}\right)^3 = \frac{S(W) + 3\zeta^2}{S(W) + 3\zeta}.$$

With these preliminaries out of the way, we fix an integer  $n \geq 4$ , an integer  $M \geq 2$ , and take

$$f(X) := X^n - nM^{n-1}X.$$

Consider the one parameter family of curves in  $(X, W)$  space over  $\mathbb{Q}(T)$

$$T - f(X) = S(W).$$

This family has an automorphism of order three, which we will denote  $\sigma$  if no confusion can occur, given by

$$(X, W) \mapsto (X, \sigma(W)) = (X, 1/(1 - W)).$$

If we extend scalars to  $\mathbb{Q}(\zeta)(T)$ , we can write this curve as

$$\frac{T - f(X) + 3\zeta^2}{T - f(X) + 3\zeta} = \left(\frac{W + \zeta^2}{W + \zeta}\right)^3.$$

Now define

$$Y := \frac{W + \zeta^2}{W + \zeta}.$$

Then in  $(X, Y)$  space, we have the curve

$$Y^3 = \frac{T - f(X) + 3\zeta^2}{T - f(X) + 3\zeta} = 1 - \frac{3\zeta - 3\zeta^2}{T - f(X) + 3\zeta},$$

on which the automorphism of order three has become the obvious automorphism  $(X, Y) \mapsto (X, \zeta Y)$ .

For  $F(T) := F_{crit.val.}(T)$  the critical value polynomial for  $f(X) - 3\zeta_3$ , and for  $A := 3\zeta - 3\zeta^2$ , we saw in Section 3 that this last family has a projective smooth model over

$$S := \text{Spec}(\mathbb{Z}[\zeta_3, 1/3, 1/(2n), 1/A, 1/\Delta_{crit,A}(T)][T][1/(F(T)F(T-A))]).$$

So if we replace each of the quantities

$$1/A, 1/\Delta_{crit,A}(T), (F(T)F(T-A))$$

by its Norm from  $\mathbb{Q}(\zeta)$  down to  $\mathbb{Q}$ , we find that our family

$$T - f(X) = S(W)$$

has a projective smooth model

$$\pi : \mathcal{C} \rightarrow S_0$$

over

$$S_0 := \text{Spec}(\mathbb{Z}[1/3, 1/(2n), 1/\text{Norm}(A), 1/\text{Norm}(\Delta_{crit,A}(T))][T][1/\text{Norm}(F(T)F(T-A))]).$$

For any  $\ell$ , we have the lisse  $\mathbb{Q}_\ell$  sheaf  $\mathcal{F}_\ell := R^1\pi_1\mathbb{Q}_\ell$  on  $S_0[1/\ell]$ . When  $\ell \equiv 1 \pmod{3}$ , and  $\chi$  is a character of full order three, we can extract the  $\chi$ -component  $\mathcal{F}_\ell^\chi$ , which is lisse of rank  $2n - 2$  and pure of weight one. View it as an  $\ell$ -adic representation

$$\rho_{f,\sigma} : \pi_1(S_0[1/\ell]) \rightarrow GL(2n - 2, \mathbb{Q}_\ell).$$

**Theorem 4.1.** *the image of  $\pi_1(S_0[1/\ell])$  in  $GL(2n - 2, \mathbb{Q}_\ell)$  is open. Moreover the image of*

$$\pi_1^{geom}(S_0[1/\ell]) := \pi_1(S_0 \otimes_{\mathbb{Z}} \mathbb{C})$$

*contains an open subgroup of  $SL(2n - 2, \mathbb{Q}_\ell)$ .*

*Proof.* As always, the key statement is the "moreover". But  $S_0 \otimes_{\mathbb{Z}} \mathbb{C}$  with its  $\mathcal{F}_\ell^\chi$  is just the same as the  $S \otimes_{\mathbb{Z}[\zeta_3]} \mathbb{C}$  with its  $\mathcal{F}_\ell^\chi$  in Theorem 3.1, applied with  $f(X) - 3\zeta$ .  $\square$

**Remark 4.2.** Exactly as in the previous sections, Hilbert Irreducibility ensures that there are infinitely many  $t \in \mathbb{Q}$  with  $\text{Norm}(F(t)F(t) - A) \neq 0$  for which the  $\ell$ -adic representation given by the  $\chi$  component of the  $H^1$  of the complete nonsingular model of the curve

$$t - f(X) = S(W)$$

over  $\mathbb{Q}$  has open image in  $GL(2n - 2, \mathbb{Q}_\ell)$ .

## 5. WORKING OVER $\mathbb{Q}$ , WHEN $\ell \equiv 1 \pmod{4}$

This section is almost identical in idea and structure to the previous section, so we simply explain the relevant miraculous identity and leave the rest to the reader.

We have the automorphism of the rational function field  $\mathbb{Q}(W)$  given by the fractional linear transformation

$$\sigma : W \mapsto \frac{W - 1}{W + 1}.$$

The trace rational function

$$S(W) := W + \sigma(W) + \sigma^2(W) + \sigma^3(W) = \frac{W^4 - 6W^2 + 1}{X(X^2 - 1)}$$

is  $\sigma$ -invariant. After extending scalars from  $\mathbb{Q}(W)$  to  $\mathbb{Q}(i)(W)$ ,  $i$  being a primitive fourth root of unity, the quantity

$$R(W) := \frac{W + i}{W - i}$$

transforms under  $\sigma$  by

$$\sigma(R(W)) = iR(W).$$

The miraculous identity is

$$R(W)^4 := \left(\frac{W + i}{W - i}\right)^4 = \frac{S(W) + 4i}{S(W) - 4i}.$$

With these preliminaries out of the way, we fix an integer  $n \geq 4$ , an integer  $M \geq 2$ , and take

$$f(X) := X^n - nM^{n-1}X.$$

Consider the one parameter family of curves in  $(X, W)$  space over  $\mathbb{Q}(T)$

$$T - f(X) = S(W).$$

This family has an automorphism of order four, which we will denote  $\sigma$  if no confusion can occur, given by

$$(X, W) \mapsto (X, \sigma(W)) = (X, (W - 1)/(W + 1)).$$

If we extend scalars to  $\mathbb{Q}(i)(T)$ , we can write our curve as

$$\frac{T - f(X) + 4i}{T - f(X) - 4i} = \left(\frac{W + i}{W - i}\right)^4.$$

Now define

$$Y := \frac{W + i}{W - i}.$$

Then in  $(X, Y)$  space, we have the curve

$$Y^4 = \frac{T - f(X) + 4i}{T - f(X) - 4i} = 1 - \frac{-8i}{T - f(X) - 4i},$$

on which the automorphism of order four has become the obvious automorphism  $(X, Y) \mapsto (X, iY)$ .

So if we replace each of the quantities

$$1/A, 1/\Delta_{crit,A}(T), (F(T)F(T - A))$$

by its Norm from  $\mathbb{Q}(i)$  down to  $\mathbb{Q}$ , we find that our family

$$T - f(X) = S(W)$$

has a projective smooth model

$$\pi : \mathcal{C} \rightarrow S_0$$

over

$$S_0 := \text{Spec}(\mathbb{Z}[1/(2n), 1/\text{Norm}(A), 1/\text{Norm}(\Delta_{crit,A}(T))][T][1/\text{Norm}(F(T)F(T - A))]).$$

For any  $\ell$ , we have the lisse  $\mathbb{Q}_\ell$  sheaf  $\mathcal{F}_\ell := R^1\pi_!\mathbb{Q}_\ell$  on  $S_0[1/\ell]$ . When  $\ell \equiv 1 \pmod{4}$ , and  $\chi$  is a character of full order four, we can extract the  $\chi$ -component  $\mathcal{F}_\ell^\chi$ , which is lisse of rank  $2n - 2$  and pure of weight one. View it as an  $\ell$ -adic representation

$$\rho_{f,\sigma} : \pi_1(S_0[1/\ell]) \rightarrow GL(2n - 2, \mathbb{Q}_\ell).$$

**Theorem 5.1.** *the image of  $\pi_1(S_0[1/\ell])$  in  $GL(2n - 2, \mathbb{Q}_\ell)$  is open. Moreover the image of*

$$\pi_1^{geom}(S_0[1/\ell]) := \pi_1(S_0 \otimes_{\mathbb{Z}} \mathbb{C})$$

*contains an open subgroup of  $SL(2n - 2, \mathbb{Q}_\ell)$ .*

*Proof.* As always, the key statement is the "moreover". But  $S_0 \otimes_{\mathbb{Z}} \mathbb{C}$  with its  $\mathcal{F}_\ell^\chi$  is just the same as the  $S \otimes_{\mathbb{Z}[\zeta_3]} \mathbb{C}$  with its  $\mathcal{F}_\ell^\chi$  in Theorem 3.1, applied with  $f(X) + 4i$ .  $\square$

**Remark 5.2.** Exactly as in the previous sections, Hilbert Irreducibility ensures that there are infinitely many  $t \in \mathbb{Q}$  with  $\text{Norm}(F(t)F(t) - A) \neq 0$  for which the  $\ell$ -adic representation given by the  $\chi$  component of the  $H^1$  of the complete nonsingular model of the curve

$$t - f(X) = S(W)$$

over  $\mathbb{Q}$  has open image in  $GL(2n - 2, \mathbb{Q}_\ell)$ .

## 6. INDEPENDENCE OF $\ell$

In each of the previous sections, we began with a cyclotomic field  $K := \mathbb{Q}(\zeta_N)$ ,  $N \geq 3$ , a number field  $E$ , a proper smooth one-parameter family

$$\pi : \mathcal{C} \rightarrow S$$

of curves over a dense open set  $S$  of  $\text{Spec}(\mathcal{O}_E[T])$ , a character  $\chi$  of either  $\mathbb{Z}/N\mathbb{Z}$  or of  $\mu_N(\mathbb{Q}(\zeta_N))$  with values in  $\mu_N(\mathbb{Q}(\zeta_N))$  of full order  $N$ , and an action of either  $\mathbb{Z}/N\mathbb{Z}$  or of  $\mu_N(\mathbb{Q}(\zeta_N))$  on the family.

Over  $\mathbb{C}$ ,  $\mathcal{F}^{an} := R^1(\pi^{an})_! K$  is a  $K$ -local system on  $S^{an}$  whose  $\chi$  component  $(\mathcal{F}^{an})^\chi$ , under the action of either  $\mathbb{Z}/N\mathbb{Z}$  or of  $\mu_N(\mathbb{Q}(\zeta_N))$ , has rank

$$d := \text{rank}(\mathcal{F}^{an})^\chi.$$

For each finite place  $\lambda$  of  $K$ , with completion  $K_\lambda$ , we have the  $K_\lambda$ -local system  $R^1\pi_! K_\lambda$  on  $S[1/\ell]$ , whose  $\chi$  component has the same rank  $d$ . For each  $t \in S(E)$ , the  $\chi$ -component  $H^1(\mathcal{C}_t \otimes_E \bar{E}, K_\lambda)^\chi$  is a representation of  $\text{Gal}(\bar{E}/E)$ , and for variable  $\lambda$  these form a compatible system of  $\lambda$ -adic representations

$$\rho_{\lambda,t} : \text{Gal}(\bar{E}/E) \rightarrow GL(d, K_\lambda).$$

For each  $\lambda$ , we denote by  $G_{\lambda,t}$  the algebraic group over  $K_\lambda$  which is the Zariski closure in  $GL(d)$  of the image of  $\rho_{\lambda,t}$ .

When we fix a place  $\lambda$  of  $K$  of residue characteristic  $\ell \equiv 1 \pmod{N}$ , then  $K_\lambda$  is  $\mathbb{Q}_\ell$ . For a fixed such  $\lambda$ , we showed the existence of infinitely many  $t \in E$  for which the image of  $\rho_{\lambda,t}$  is open in  $GL(d, \mathbb{Q}_\ell)$ .



**Theorem 6.1.** *Suppose that  $\lambda_1$  is a place of  $K$  of residue characteristic  $\ell_1 \equiv 1 \pmod{N}$ , and  $t \in S(E)$  is such that the image of  $\rho_{\lambda_1, t}$  is open in  $GL(d, \mathbb{Q}_{\ell_1})$ . Then for every place  $\lambda_2$  of residue characteristic  $\ell_2 \equiv 1 \pmod{N}$ , the image of  $\rho_{\lambda_2, t}$  is open in  $GL(d, \mathbb{Q}_{\ell_2})$ .*

*Proof.* For any such  $\lambda_2$  of  $K$ ,  $H^1(\mathcal{C}_t \otimes_E \overline{E}, \mathbb{Q}_{\ell_2})^\times$  is a direct factor of  $H^1(\mathcal{C}_t \otimes_E \overline{E}, \mathbb{Q}_{\ell_2})$ . By Faltings [Fal, §5, Satz 4],  $\rho_{\lambda_2, t}$  is completely reducible, and hence  $G_{\lambda_2, t}$  is reductive. Denote by  $\Gamma_{\lambda_2, t}$  the image of  $\rho_{\lambda_2, t}$ . By Bogomolov [Bog, Thm. 1],  $\Gamma_{\lambda_2, t}$  is open in  $G_{\lambda_2, t}(\mathbb{Q}_{\ell_2})$ . So it suffices to show that  $G_{\lambda_2, t}$  is  $GL(d)$ . According to Hui [Hui, 3.22 and 3.19], the rank of the derived group  $(G_{\lambda, t}^0)^{der}$  is independent of the auxiliary choice of  $\lambda$ . For  $\lambda_1$ , we know that  $G_{\lambda_1, t} = GL(d)$ , so its derived group is  $SL(d)$ , of rank  $d - 1$ . Therefore  $(G_{\lambda_2, t}^0)^{der}$  is a connected semisimple subgroup of  $SL(d)$  of rank  $d - 1$ . The only such subgroup is  $SL(d)$  itself. The determinant of  $\rho_{\lambda_2, t}$  is of infinite order, being pure of weight  $d \neq 0$ . Therefore  $G_{\lambda_2, t}$  must be  $GL(d)$ .  $\square$

**Remark 6.2.** See [H-L, Thm.1] and [C-H-T, Thm.1.2] for other results, in slightly different contexts, of the same type.

## 7. EXAMPLES IN HIGHER DIMENSION

In this section, we give examples built with higher dimensional varieties. For given integers  $n \geq 2$  and  $d \geq 3$ , and  $k$  a field in which  $d$  is invertible, a polynomial  $f(X_1, \dots, X_n) \in k[X_1, \dots, X_n]$  is a “strong Deligne polynomial” if it satisfies the following two conditions.

- (1) The affine hypersurface of equation  $f = 0$  in  $\mathbb{A}^n$  is smooth of dimension  $n - 1$ .
- (2) Write  $f$  as the sum of homogeneous forms  $f = f_d + f_{d-1} + \dots + f_0$ . The projective hypersurface of equation  $f_d = 0$  in  $\mathbb{P}^{n-1}$  is smooth of dimension  $n - 2$ .

Here is another way to think of these conditions. Pass to the homogeneous form

$$F(X_1, \dots, X_n, Z) \in k[X_1, \dots, X_n, Z] := f_d + Zf_{d-1} + Z^2f_{d-2} + \dots + Z^df_0.$$

The conditions on  $F$  are that  $F = 0$  defines a smooth hypersurface  $H_F$  of dimension  $n - 1$  in  $\mathbb{P}^n$  and that the intersection  $H_F \cap (Z = 0)$  defines a smooth hypersurface of dimension  $n - 2$  in  $\mathbb{P}^{n-2}$ .

This point of view makes clear that in the affine space  $\text{Poly}(n, d)$  over  $\mathbb{Z}[1/d]$  of all degree  $d$  polynomials in  $n$  variables (the coefficients being the coordinate functions), the special Deligne polynomials form

a dense (over any field  $k$  in which  $d$  is invertible, the polynomial  $f := 1 + \sum_{i=1}^n X_i^d$  is a strong Deligne polynomial) open set  $SD(n, d)$ .

Now fix an integer  $N \geq 3$ , and write  $K := \mathbb{Q}(\zeta_N)$ . Over the parameter space

$$S := SD(n, d) \otimes_{\mathbb{Z}[1/d]} \mathcal{O}_K[1/N, 1/d]$$

of strong Deligne polynomials  $f$ , we have the family  $\pi : \mathcal{H} \rightarrow S$  of smooth affine hypersurfaces of dimension  $n$  of equation

$$Y^N = f(X_1, \dots, X_N)$$

and the obvious action on it of the group  $\mu_N(K)$ . We fix a character  $\chi$  of  $\mu_N(K)$  of full order  $N$ .

**Theorem 7.1.** *Suppose  $n \geq 2$ ,  $d \geq 3$ ,  $N \geq 3$ , and  $d$  is nonzero mod  $N$  (i.e.,  $\chi^d \neq 1$ ). For each finite place  $\lambda$  of  $K$ , denote by  $\ell$  its residue characteristic. Then we have the following results.*

- (1) *The sheaves  $(R^i \pi_! K_\lambda)^\chi$  on  $S[1/\ell]$  vanish for  $i \neq n$ .*
- (2) *The sheaf  $\mathcal{F}_{\lambda, \chi} := (R^n \pi_! K_\lambda)^\chi$  on  $S[1/\ell]$  is pure of weight  $n$  and lisse of rank  $(d-1)^n$ . Its trace function is given by*

$$\text{Trace}(\text{Frob}_{k,f} | \mathcal{F}_{\lambda, \chi}) = (-1)^n \sum_{x \in k^n} \chi_k(f(x)).$$

- (3) *Over any field  $k$  in which  $d\ell$  is invertible, the geometric monodromy group of  $\mathcal{F}_{\lambda, \chi} | S \otimes k$  contains  $SL((d-1)^n)$ .*

*Proof.* This is proven in [Ka-MMP, 5.1.9, 5.1.14, and 5.2.2 1)].  $\square$

Exactly as in the first section, we get the following corollary.

**Corollary 7.2.** *Under the hypotheses of the theorem, for any  $\lambda$  of residue characteristic  $\ell \equiv 1 \pmod{N}$ , the image of the representation*

$$\rho_{\lambda, \chi} : \pi_1(S[1/\ell]) \rightarrow GL((d-1)^n, \mathbb{Q}_\ell)$$

*which “is” the sheaf  $\mathcal{F}_{\lambda, \chi}$  is open in  $GL((d-1)^n, \mathbb{Q}_\ell)$ .*

Because the parameter space  $S$  is rational, Hilbert irreducibility and the independence of  $\ell$  results give us

**Theorem 7.3.** *Fix a place  $\lambda_1$  of residue characteristic  $\ell_1 \equiv 1 \pmod{N}$ . There exist infinitely many strong Deligne polynomials  $f \in K[X_1, \dots, X_n]$  of degree  $d$  for which the action of  $\text{Gal}(\overline{K}/K)$  on*

$$H_c^n((Y^N = f(x)) \otimes_K \overline{K}, K_{\lambda_1} = \mathbb{Q}_{\ell_1})^\chi$$

*has open image in  $GL((d-1)^n, K_{\lambda_1}) = GL((d-1)^n, \mathbb{Q}_{\ell_1})$ . Moreover, given such an  $f$ , then for any other place  $\lambda_2$  of residue characteristic  $\ell_2 \equiv 1 \pmod{N}$ , the action of  $\text{Gal}(\overline{K}/K)$  on*

$$H_c^n((Y^N = f(x)) \otimes_K \overline{K}, K_{\lambda_2} = \mathbb{Q}_{\ell_2})^\chi$$

has open image in  $GL((d-1)^n, K_{\lambda_2}) = GL((d-1)^n, \mathbb{Q}_{\ell_2})$ .

### 8. EXPLICIT EXAMPLES IN HIGHER DIMENSION

Thus far, we have worked over the **entire** parameter space of **all** strong Deligne polynomials of given degree  $d$  in a given number  $n \geq 2$  of variables. Over a field in which  $d$  is invertible, suppose we have a polynomial  $f(X) := f(X_1, \dots, X_n)$  of degree  $d$  whose leading form  $f_d$  defines a smooth hypersurface in  $\mathbb{P}^{n-1}$  (a ‘‘Deligne polynomial’’) and which has only finitely many critical points (points where all the  $\partial f / \partial f X_i$  vanish). For any  $t$  which is not a critical value (the value of  $f$  at a critical point),  $t - f(X)$  is a strong Deligne polynomial.

As before, we put  $K := \mathbb{Q}(\zeta_N)$ ,  $N \geq 3$ . Fix a degree  $d \geq 3$  which is nonzero mod  $N$ , and a number  $n \geq 2$  of variables. Given a Deligne polynomial  $f \in \mathcal{O}_K[X_1, \dots, X_n]$  of degree  $d$ , with only finitely many critical points (in  $\mathbb{C}^n$ ), denote by

$$F_{crit}(T) \in \mathcal{O}_K[T]$$

a choice of (not necessarily monic) polynomial whose roots are the distinct critical values of  $f$ , and by

$$\Delta_{crit} \in \mathcal{O}_K$$

its discriminant. We will exhibit such an  $f$  such that for any  $\chi$  of full order  $N$ , the one parameter (‘‘ $t$ ’’ the parameter) family of character sums

$$(-1)^n \sum_{x \in k^n} \chi(t - f(x))$$

over

$$S := \text{Spec}(\mathcal{O}_K[1/d, 1/\Delta_{crit}, 1/N, T][1/F_{crit}(T)]),$$

i.e., for  $\pi : \mathcal{H} \rightarrow S$  the family of varieties of equation

$$Y^N = t - f(X_1, \dots, X_n),$$

the  $(R^i \pi_! K_\lambda)^x$  vanish for  $i \neq n$ , and

$$\mathcal{F}_{\chi, \lambda} := (R^n \pi_! K_\lambda)^x$$

is lisse of rank  $(d-1)^n$ , pure of weight  $n$ , and its  $G_{geom}$  contains  $SL((d-1)^n)$ .

Once we have such an  $f$ , then there are infinitely many  $t \in K$  such that for every place  $\lambda_i$  of  $K$  with residue characteristic  $\ell_i \equiv 1 \pmod{N}$ , the action of  $Gal(\overline{K}/K)$  on

$$H_c^n((Y^N = t - f(X)) \otimes_K \overline{K}, K_{\lambda_i} = \mathbb{Q}_{\ell_i})^x$$

has open image in  $GL((d-1)^n, K_{\lambda_i}) = GL((d-1)^n, \mathbb{Q}_{\ell_i})$ .

Here is our  $f(X_1, \dots, X_n)$ , for given  $n$  and  $d$ . It is a variation on the idea behind [Ka-MMP, 6.7]. Define

$$g(X) := X^d - dX,$$

Choose a prime  $p$  which is 1 mod  $d - 1$  and take

$$f(X_1, \dots, X_n) := \sum_{i=1}^n p^i g(X_i).$$

The critical points of this  $f$  are the points  $(a_1, \dots, a_n)$  where each  $a_i$  is a  $d - 1$  root of unity. The critical values are the sums

$$\sum_{i=1}^n p^i g(a_i) = \sum_{i=1}^n p^i (1 - d) a_i.$$

We first show that these  $(d - 1)^n$  sums are all distinct. Because  $p \equiv 1 \pmod{d - 1}$ , the  $a_i$  are Teichmüller points in  $\mathbb{Z}_p$ . After dividing by  $1 - d$ , we are looking at the expression of  $(d - 1)^n$  distinct elements of  $\mathbb{Z}_p$  expanded in the base  $p$ , using 0 and Teichmüller points as “digits”.

The vanishing of the  $(R^i \pi_1 K_\lambda)^\times$  on  $S[1/\ell]$  for  $i \neq n$  results by proper base change from the universal assertion (1) of Theorem 7.1, as does the fact that our  $\mathcal{F}_{\chi, \lambda}$  on  $S[1/\ell]$  is lisse of rank  $(d - 1)^n$  and pure of weight  $n$ . What must be proven is that its  $G_{geom}$  contains  $SL((d - 1)^n)$ .

For  $i = 1, \dots, n$ , and fixed  $\lambda$ , define

$$L_i := ((p^i g)_* \mathbb{Q}_\ell / \mathbb{Q}_\ell) \otimes K_\lambda.$$

Exactly as in [Ka-MMP, 6.7.9-11], the purity of our  $\mathcal{F}_{\chi, \lambda}$  shows that is the iterated middle convolution of the Kummer sheaf  $\mathcal{L}_\chi$  with the  $L_i$ :

$$\mathcal{F}_{\chi, \lambda} \cong \mathcal{L}_\chi \star_{mid,+} L_1 \star_{mid,+} L_2 \dots \star_{mid,+} L_n.$$

This sheaf is tame at  $\infty$ , lisse outside the  $(d - 1)^n$  critical values, and at each critical value its local monodromy is a pseudoreflection of determinant  $\chi \chi_{quad}^n$ .

There is a further property, nonpunctuality, we need of the multiple middle convolution  $L_1 \star_{mid,+} L_2 \dots \star_{mid,+} L_n$ , namely that it is the direct sum of irreducible middle extension sheaves. To see this, we work successively from the right. Applying [Ka-MMP, 6.1.12] successively, it suffices to verify that for each  $m = 1, \dots, n - 1$ ,

$$(**) \quad \text{all ratios } \frac{\sum_{i=1}^m p^i (a_i - b_i)}{a_0 - b_0} \neq 1,$$

for all choices of  $d - 1$  roots of unity  $a_0 \neq b_0, a_1, \dots, a_m, b_1, \dots, b_m$ . To see that this holds, simply cross multiply and rewrite this as

$$b_0 + \sum_{i=1}^m p^i a_i \neq a_0 + \sum_{i=1}^m p^i b_i.$$

Viewed in  $\mathbb{Z}_p$ , this inequality is obvious, already the first digits are different.

If in addition we knew that  $\mathcal{F}_{\chi, \lambda}$  was geometrically irreducible, then by [Ka-ACT, 5.11], we would know that its  $G_{geom}$  contains  $SL((d-1)^n)$  provided that  $n \geq 2$  and either

$$(d - 1)^n > 4$$

or

$$d = 3, n = 2, \text{ and } \chi \text{ does not have order } 3.$$

[When  $d = 3$  and  $n = 2$  the requirement that  $N$  be nonzero mod  $d = 3$  forces our  $\chi$  to have order prime to 3.] To show the geometric irreducibility, it is enough to work in some large characteristic  $P$  which is 1 mod  $d - 1$  (not to be confused with  $p$  used in defining  $f$  as the sum of  $p^i g(X_i)$ ), with  $P > p^{2n(d-1)} - 1$  (to be sure that none of  $p, p^2, \dots, p^n$  is a  $2d - 2$  root of unity mod  $P$ ), with  $P > 2d - 2$ , and such that (\*\*) above holds in  $\mathbb{F}_P$ , and to show there that

$$L_1 \star_{mid,+} L_2 \dots \star_{mid,+} L_n$$

is geometrically irreducible (because middle convolution with  $\mathcal{L}_\chi$  preserves geometric irreducibility). For this, it suffices to show that on  $\mathbb{G}_m$ , the tensor product of the (lisse on  $\mathbb{G}_m$ ) Fourier Transforms

$$FT(L_1) \otimes FT(L_2) \dots \otimes FT(L_n)$$

is geometrically irreducible (its extension by direct image across 0 is the Fourier Transform of  $L_1 \star_{mid,+} L_2 \dots \star_{mid,+} L_n$ , by the nonpunctuality of the latter).

From [Ka-ESDE, 7.10.4], we know that, in our sufficiently large characteristic  $P$ , each individual  $FT(L_i)$  has  $G_{geom}$  either  $Sp(d - 1)$ , if  $d$  is odd, or  $\pm SL(d - 1)$ , if  $d$  is even. So it suffices to show that the  $G_{geom}$  of the direct sum is the  $n$ -fold self product  $(Sp(d - 1))^n$  in the  $d$  odd case, and contains the  $n$ -fold self product  $(SL(d - 1))^n$  in the  $d$  even case. For then the tensor product is the product of irreducible representations of the factors, so is an irreducible representation of that product.

By Goursat-Kolchin-Ribet, cf. [Ka-ESDE, 1.8.2], it suffices to show that for  $i \neq j$ , there is no lisse rank one  $\mathcal{L}$  on  $\mathbb{G}_m$  for which  $FT(L_i) \cong \mathcal{L} \otimes FT(L_j)$  or for which  $FT(L_i)^\vee \cong \mathcal{L} \otimes FT(L_j)$ . In the  $d$  odd case,

$\mathcal{L}$  must have order dividing 2, as the only scalars in  $Sp(d-1)$  are  $\pm 1$ . The  $I(\infty)$  representation of  $FT(L_i)$  is the direct sum

$$\bigoplus_{\zeta \in \mu_{d-1}} \mathcal{L}_{\chi_{quad}} \otimes \mathcal{L}_{\psi(p^i(1-d)\zeta x)}.$$

The only  $\mathcal{L}$  of order dividing 2 is either  $\mathbb{Q}_\ell$  or  $\mathcal{L}_{\chi_{quad}}$ . The second is not allowed, because it “removes” the factor  $\mathcal{L}_{\chi_{quad}}$  from each piece of the  $I(\infty)$ -representation. The first is not allowed, because the  $I(\infty)$ -representation of  $FT(L_j)$  is

$$\bigoplus_{\zeta \in \mu_{d-1}} \mathcal{L}_{\chi_{quad}} \otimes \mathcal{L}_{\psi(p^j(1-d)\zeta x)},$$

and for  $i \neq j$ , no  $p^i\zeta$  is any  $p^j\zeta'$ , for  $\zeta, \zeta'$  any  $d-1$  roots of unity.

In the  $d$  even case, where each factor's  $G_{geom}$  is  $\pm SL(d-1)$ ,  $\mathcal{L}$  must have order dividing  $2(d-1)$ . As  $P > 2d-2$ , such an  $\mathcal{L}$  is necessarily tame, so of the form  $\mathcal{L}_\chi$  for a character  $\chi$  of order dividing  $2d-2$ . If  $\chi$  is nontrivial, we multiply the  $\mathcal{L}_{\chi_{quad}}$  factors in the  $I(\infty)$  summands by  $\chi$ , not allowed. So it remains only to rule out both  $FT(L_i) \cong FT(L_j)$  and  $FT(L_i)^\vee \cong \otimes FT(L_j)$ . Again we compare  $I(\infty)$  summands. The first case cannot happen, for the same reason as above: no  $p^i\zeta$  is any  $p^j\zeta'$ . The second case cannot happen because no  $-p^i\zeta$  is any  $p^j\zeta'$ . This concludes the proof that our  $\mathcal{F}_{\chi,\lambda}$  on  $S[1/\ell]$  is geometrically irreducible.

## 9. SOME OPEN QUESTIONS

Zarhin has shown that for hyperelliptic curves over  $\mathbb{Q}$  (or indeed over any field  $K$  which is finitely generated over  $\mathbb{Q}$ ) of the form  $Y^2 = f(X)$ , with  $f$  a polynomial of degree  $n$ , various explicit conditions on  $n$  and on the galois group of  $f$  over  $K$  guarantee that the  $\ell$ -adic representation on  $H^1$  has image which is open in the group of symplectic similitudes. For example, if  $f$  has degree  $n \geq 5$  and has Galois group over  $\mathbb{Q}$  either  $S_n$  or  $A_n$ , then this holds, cf. [Zar1]. See the papers [Zar2], [Zar3], [Zar4] for more such spectacular results. What if any are the analogues of Zarhin's results for  $\chi$  components of  $H_c^1$  of superelliptic<sup>1</sup> curves over cyclotomic fields, and their descents to  $\mathbb{Q}$ , when they exist?

## REFERENCES

- [A-P] Achter, J.; Pries, R., The integral monodromy of hyperelliptic and trielliptic curves. *Math. Ann.* 338 (2007), no. 1, 187-206.
- [Bbk-Lie I] Bourbaki, N. *Éléments de mathématique*. Fasc. XXVI. Groupes et algèbres de Lie. Chapitre I: Algèbres de Lie. (French) Seconde édition. *Actualités Scientifiques et Industrielles*, No. 1285 Hermann, Paris 1971 146 pp.

<sup>1</sup>See [A-P, Thm.3.8] for the monodromy of the universal family of tri-elliptic curves.

- [BCELMP] Boxer, G.; Calegari, F.; Emerton, M.; Levin, B.; Madapusi Pera, K.; Patrikis, S., Compatible systems of Galois representations associated to the exceptional group  $E_6$ , arXiv:1805.09777.
- [Bog] Bogomolov, F. A., Sur l'algébricité des représentations  $l$ -adiques. C. R. Acad. Sci. Paris Sr. A-B 290 (1980), no. 15, A701-A703.
- [C-H-T] Cadoret, A., Hui, C.Y., Tamagawa, A., Geometric monodromy-semisimplicity and maximality. Ann. of Math. (2) 186 (2017), no. 1, 205-236.
- [C-R] Cornut, C., Ray, J., Generators of the pro- $p$  Iwahori and Galois representations. Int. J. Number Theory 14 (2018), no. 1, 37-53.
- [De-Mos] Deligne, P., Mostow, G.D., Monodromy of hypergeometric functions and non-lattice integral monodromy, Pub. Math. I.H.E.S. 63 (1986), 5-89.
- [De-Weil II] Deligne, P., La conjecture de Weil II, Pub. Math. I.H.E.S. 52 (1981), 313-428.
- [D-R] Dettweiler, M.; Reiter, S., Rigid local systems and motives of type  $G_2$ . With an appendix by Michael Dettweiler and Nicholas M. Katz. Compos. Math. 146 (2010), no. 4, 929-963.
- [Fal] Faltings, G., Endlichkeitssätze für abelsche Varietäten über Zahlkörpern. Invent. Math. 73 (1983), no. 3, 349-366.
- [Gre] Greenberg, R., Galois representations with open image. Ann. Math. Qué. 40 (2016), no. 1, 83-119.
- [Gro-FL] Grothendieck, A., Formule de Lefschetz et rationalité des fonctions  $L$ , Seminaire Bourbaki 1964-65, Exposé 279, reprinted in Dix Exposés sur la cohomologie des schémas, North-Holland, 1968.
- [Hui] Hui, Chun Yin, Monodromy of Galois representations and equal-rank subalgebra equivalence. Math. Res. Lett. 20 (2013), no. 4, 705-728.
- [H-L] Hui, C. Y., Larsen, M., Type A images of Galois representations and maximality. Math. Z. 284 (2016), no. 3-4, 989-1003.
- [Ka-ACT] Katz, N., Affine cohomological transforms, perversity, and monodromy. J. Amer. Math. Soc. 6 (1993), no. 1, 149-222.
- [Ka-ESDE] Katz, N., Exponential sums and differential equations. Annals of Mathematics Studies, 124. Princeton Univ. Press, Princeton, NJ, 1990. xii+430 pp.
- [Ka-GKM] Katz, N., Gauss Sums, Kloosterman Sums, and Monodromy Groups, Annals of Mathematics Studies, 116. Princeton Univ. Press, Princeton, NJ, 1988. ix+246 pp.
- [Ka-MMP] Katz, N., Moments, Monodromy, and Perversity. Annals of Mathematics Studies, 159. Princeton University Press, Princeton, NJ, 2005. viii+475 pp.
- [Ka-RLS] Katz, N., Rigid Local Systems. Annals of Mathematics Studies, 139. Princeton University Press, Princeton, NJ, 1996. viii+223 pp.

- [Ka-SE] Katz, N., Sommes Exponentielles, Astérisque 79, Course taught at the University of Paris, Orsay, Fall 1979. With a preface by Luc Illusie. Notes written by Gérard Laumon. Astérisque 79. Société Mathématique de France, Paris, 1980.
- [Ka-ST] Katz, N., Sato-Tate in the higher dimensional case: elaboration of 9.5.4 in Serre's  $NX(p)$  book. *Enseign. Math.* 59 (2013), no. 3-4, 359-377
- [Poch] Pochhammer, L., Über hypergeometrische Functionen höherer Ordnung, *J. r. und ang. Math (Crelle)* 71 (1870), 316-362.
- [Se-AbLRep] Serre, J. P., Abelian  $l$ -adic representations and elliptic curves. With the collaboration of Willem Kuyk and John Labute. Second edition. *Advanced Book Classics*. Addison-Wesley Publishing Company, Advanced Book Program, Redwood City, CA, 1989. xxiv+184 pp.
- [Se-pDiv] Serre, J. P., Sur les groupes de Galois attachés aux groupes  $p$ -divisibles. 1967 *Proc. Conf. Local Fields (Driebergen, 1966)* pp. 118-131 Springer, Berlin.
- [Se-MW] Serre, J. P., *Lectures on the Mordell-Weil Theorem, Third Edition, Aspects of Mathematics E15*, Springer Fachmedien Wiesbaden GmbH, 1997.
- [SGA4] Artin, M., Grothendieck, A., Verdier, J. L., Séminaire de Géométrie Algébrique du Bois Marie, SGA 4, Tome III, Springer Lecture Notes in Mathematics 305, Springer Verlag, 1973.
- [Ter] Terasoma, T., Complete intersections with middle Picard number 1 defined over  $\mathbb{Q}$ . *Math. Z.* 189 (1985), no. 2, 289-296.
- [Up] Upton, M. G., Galois representations attached to Picard curves. *J. Algebra* 322 (2009), no. 4, 1038-1059.
- [Wag] Wagner, A., Collineation groups generated by homologies of order greater than 2, *Geom. Dedicata* 7 (1978), no. 4, 387-398.
- [Yun] Yun, Z., Motives with exceptional Galois groups and the inverse Galois problem, *Invent. Math.* 196 (2014), no. 2, 267-337.
- [Zar1] Zarhin, Y., Very simple 2-adic representations and hyperelliptic Jacobians. Dedicated to Yuri I. Manin on the occasion of his 65th birthday. *Mosc. Math. J.* 2 (2002), no. 2, 403-431.
- [Zar2] Zarhin, Y., Families of absolutely simple hyperelliptic Jacobians. *Proc. Lond. Math. Soc.* (3) 100 (2010), no. 1, 24-54.
- [Zar3] Zarhin, Y., Galois groups of Mori trinomials and hyperelliptic curves with big monodromy. *Eur. J. Math.* 2 (2016), no. 1, 360-381
- [Zar4] Zarhin, Y., Two-dimensional families of hyperelliptic Jacobians with big monodromy. *Trans. Amer. Math. Soc.* 368 (2016), no. 5, 3651-3672.
- E-mail address:* `nmk@math.princeton.edu`