

Asymptotic structure. IV. A counterexample to the weak coarse Menger conjecture

Tung Nguyen¹
Princeton University,
Princeton, NJ 08544, USA

Alex Scott²
University of Oxford,
Oxford, UK

Paul Seymour³
Princeton University,
Princeton, NJ 08544, USA

July 20, 2025; revised July 31, 2025

¹Supported by AFOSR grants A9550-19-1-0187 and FA9550-22-1-0234, and by NSF grants DMS-1800053 and DMS-2154169.

²Supported by EPSRC grant EP/X013642/1

³Supported by AFOSR grants A9550-19-1-0187 and FA9550-22-1-0234, and by NSF grants DMS-1800053 and DMS-2154169.

Abstract

Coarse graph theory concerns finding “coarse” analogues of graph theory theorems, replacing disjointness with being far apart. One of the most interesting open questions is to find a coarse analogue of Menger’s theorem (the latter characterizes when there are k vertex-disjoint paths between two given sets S, T of vertices of a graph). We showed in an earlier paper that the most natural such analogue is false, but a weaker statement remained as a popular open question. Here we show that it is also false.

More exactly, suppose that S, T are sets of vertices of a graph G , and there do not exist k paths between S, T , pairwise at distance at least c . To make an analogue of Menger’s theorem, one would like to prove that there must be a small set $X \subseteq V(G)$ such that every $S - T$ path of G passes close to a member of X : but how small and how close? In view of Menger’s theorem, one would hope for $|X| < k$ and “close” some function of k, c (and indeed, this was conjectured by two groups, Georgakopoulos and Papasoglu, and independently, by Albrechtsen, Huynh, Jacobs, Knappe and Wollan); but we showed that this is false, even if $c = 3$ and $k = 3$.

Here we upgrade the counterexample: we show that, even if $c = k = 3$, no pair of constants (for “small” and “close”) work. For all ℓ, m , there is a graph G and $S, T \subseteq V(G)$, such that there do not exist three $S - T$ paths pairwise with distance at least three, and yet there is no X with $|X| \leq m$ such that every $S - T$ path passes within distance at most ℓ of X .

1 Introduction

The “disjoint paths problem” asks when there is a set of k vertex-disjoint paths between sets S, T of vertices of a graph G ; and it is answered by a theorem of K. Menger from 1927 [8], that such paths exist if and only if there is no subset $X \subseteq V(G)$ of size $< k$ such that every $S - T$ path has a vertex in X . But what if we want the paths to be at least a certain distance from one another?¹ This question is motivated both by the developing area of “coarse graph theory”, which is concerned with the large-scale geometric structure of graphs (see Georgakopoulos and Papasoglu [5]), and by the algorithmic question of deciding whether such paths exist (see Bienstock [3], Kawarabayashi and Kobayashi [7], and Baligács and J. MacManus [2]).

A coarse analogue of Menger’s theorem was conjectured by Albrechtsen, Huynh, Jacobs, Knappe and Wollan [1], and independently by Georgakopoulos and Papasoglu [5]:

1.1 False conjecture: *For all integers $k, c \geq 1$ there exists $\ell > 0$ with the following property. Let G be a graph and let $S, T \subseteq V(G)$. Then either*

- *there are k paths between S, T , pairwise at distance at least c ; or*
- *there is a set $X \subseteq V(G)$ with $|X| \leq k - 1$ such that every path between S, T contains a vertex with distance at most ℓ from some member of X .*

Both sets of authors proved the result for $k = 2$, but we showed in [9] that 1.1 is false for all $k \geq 3$, even if $c = 3$ and G has maximum degree three. The case $c = 3$ is of special interest, because it is easy to see that if the result is true when $c = 3$ (for some value of k) then it is true for all $c \geq 3$ and the same value of k (apply the result when $c = 3$ to the c th power of G). And indeed, the conjecture remains open when $c = 2$, and we have nothing to say about that case in this paper.

Since the most natural extension of Menger’s theorem is false, we fall back onto what seems the next most natural, the following weaker statement:

1.2 False conjecture: *For all integers $k, c \geq 1$ there exist $m, \ell > 0$ with the following property. Let G be a graph and let $S, T \subseteq V(G)$. Then either*

- *there are k paths between S, T , pairwise at distance at least c ; or*
- *there is a set $X \subseteq V(G)$ with $|X| \leq m$ such that every path between S, T contains a vertex with distance at most ℓ from some member of X .*

We proposed this in [9], but now we will show that this too is false, even if $c = k = 3$. More exactly, we will show:

1.3 *For all integers $\ell, m \geq 1$, there is a graph G and subsets $S, T \subseteq V(G)$ such that:*

- *there do not exist three paths between S, T that pairwise have distance at least three; and*
- *for every set $X \subseteq V(G)$ with $|X| \leq m$, there is a path P between S, T such that $\text{dist}_G(P, X) > \ell$.*

Our counterexample has some vertices of large degree, but it can easily be modified into a counterexample with only one vertex of degree more than three, while keeping $c = k = 3$. We will explain this in section 3.

¹If X and Y are vertices, or sets of vertices, or subgraphs, of a graph G , then $\text{dist}_G(X, Y)$ denotes the distance between X, Y , that is, the number of edges in the shortest path of G with one end in X and the other in Y .

2 The counterexample

For all integers $\ell, m \geq 1$, we will give a construction for a quadruple (G, r, S, T) called (in this paper) an (ℓ, m) -*block*, where G is a graph, $r \in V(G)$, and S, T are disjoint subsets of $V(G) \setminus \{r\}$, both of size m . Later we will prove (by induction on m) that it has the properties that:

- $\text{dist}_G(u, v) > 2\ell$ for every two vertices $u, v \in \{r\} \cup S \cup T$;
- for every choice of $X \subseteq V(G)$ with $|X| < m$, there is a path P between S, T such that $\text{dist}_G(P, X \cup \{r\}) > \ell$; and
- for every two vertex-disjoint $S-T$ paths P, Q , either one of P, Q contains r , or $\text{dist}_G(P, Q) \leq 2$.

This will prove 1.3. (To see this, replace S, T by $S \cup \{r\}, T \cup \{r\}$.) We call r the *root*.

The construction for an $(\ell, 1)$ -block is easy: let G' be a path of length $2\ell + 1$ with ends s, t , and let $S = \{s\}$ and $T = \{t\}$; let r be a new vertex, and let G be obtained from G' by adding r as a vertex of degree zero. Then (G, r, S, T) is an $(\ell, 1)$ -block.

The construction for $(\ell, 2)$ -blocks was given in [9], but here it is again (slightly modified for convenience), illustrated in figure 1. Each dotted curve in the figure represents a path of length




Figure 1: The dotted curves represent paths of length $2\ell + 1$.

$2\ell + 1$, with interiors that are pairwise disjoint; let us call them “dotted paths”. If we delete the interiors of the “horizontal” dotted paths at the bottom of the figure, we obtain a subdivision of a uniform binary tree, rooted at r , of depth 6 in the figure²; and to make an $(\ell, 2)$ -block, we need this tree to have depth at least $2\ell + 2$. The path formed by the horizontal edges in the figure is called its *base path*; it starts in S and ends in T , and has no other vertices in $S \cup T$. Let $s_0 \in S$ and $t_0 \in T$ be the vertices not in the base path. In the figure there are 32 vertices shown in the base path; they are the leaves of the binary tree. In general there are $2^{2\ell+1}$ such vertices, if the binary tree has depth $2\ell + 2$. We call these the *anchors* of the base path. (The base path has many vertices that are not anchors, since it is a union of dotted paths). Let us number the anchors of the base path $v_1 \cdots v_n$ in order, where $v_1 \in S$ and $v_n \in T$.

An $(\ell, m + 1)$ -block is defined to be a certain combination of (ℓ, m) -blocks, explained below. Roughly, we replace each anchor of an $(\ell, 2)$ -block by a set of m vertices, and replace each dotted path between anchors by an (ℓ, m) -block, and then we identify all the roots.

Here then, more exactly, is the inductive construction. We assume that we have an (ℓ, m) -block (H, r, S, T) , and we will assemble copies of it to make an $(\ell, m + 1)$ -block. Take an $(\ell, 2)$ -block

²The “depth” of a uniform binary tree is the number of vertices in paths from root to leaf.

(H_0, r_0, S_0, T_0) as in Figure 1, and number the anchors of its base path v_1, \dots, v_n , as described above. Let $S_0 = \{v_1, s_0\}$ and $T_0 = \{v_n, t_0\}$. For $1 \leq i \leq n$, let V_i be a set of m new vertices; and for $1 \leq i < n$, let (H_i, r_0, V_i, V_{i+1}) be a copy of (H, r, S, T) , where all vertices of H_i are new except those in $V_i \cup V_{i+1} \cup \{r_0\}$. Let J be obtained from H_0 by deleting all vertices of its base path and all internal vertices of all its dotted paths. Let G be the graph obtained from the union of J, H_1, \dots, H_{n-1} by adding a path of length $2\ell + 1$ between u and each vertex of V_i , for each $u \in V(J)$ and each v_i with $1 \leq i \leq n$ such that u, v_i are joined by a dotted path of H_0 . We call these last *spines* of G . Let $S = V_1 \cup \{s_0\}$ and $T = V_n \cup \{y_0\}$. Then (G, r_0, S, T) is an $(\ell, m + 1)$ -block. (We illustrate this when $m = 2$ in figure 2.)




Figure 2: An $(\ell, 3)$ -block (except that the binary tree should have depth $2\ell + 2$, and not just five as in the figure). The dotted curves represent paths of length $2\ell + 1$. Each gray area (together with r_0) is an $(\ell, 2)$ -block, and each contains neighbours of r_0 (not shown in the figure.)

It is easy to check that if (G, r, S, T) is an (ℓ, m) -block, then every two vertices in $\{r\} \cup S \cup T$ have distance at least $2\ell + 1$, and now we will check the other properties mentioned in the bullets at the start of this section.

2.1 *If (G, r_0, S, T) is an $(\ell, m + 1)$ -block, and P, Q are paths between S, T , then either one of P, Q contains r_0 , or $\text{dist}_G(P, Q) \leq 2$.*

Proof. We proceed by induction on m , and the result is true when $m = 1$ by the result of [9], so we assume it is true for (ℓ, m) -blocks. We use the notation given in the construction for an $(\ell, m + 1)$ -block. Roughly, G was obtained from H_0 by blowing up the vertices and edges of its base path; now we want to shrink them back to H_0 , and carry P, Q to some paths P', Q' of H_0 . More exactly, let $Z = V(J) \cup V_1 \cup \dots \cup V_n$. The ends of P both belong to Z ; so P is the concatenation of a sequence of paths of G , each with distinct ends in Z and no internal vertex in Z , say P_1, \dots, P_g . Each P_i is either an edge of J , or a spine, or a path of some H_j . We may assume that no P_i contains r_0 . For $1 \leq i < n$, let B_i be the dotted path of H_0 with ends v_i, v_{i+1} . For $1 \leq i \leq g$, define a subgraph P'_i of H_0 as follows.

- If P_i is an edge of J let $P'_i = P_i$.

- If P_i is a spine with ends $u \in V(J)$ and some vertex in some V_j , let P'_i be the dotted path of H_0 between u, v_i .
- If P_i is a path of some H_j with both ends in V_j , let P'_i be the one-vertex path with vertex v_j .
- If P_i is a path of some H_j with both ends in V_{j+1} , let P'_i be the one-vertex path with vertex v_{j+1} .
- If P_i is a path of some H_j with one end in V_j and the other in V_{j+1} , let $P'_i = B_j$.

It is not necessarily true that $P'_1 \cup \dots \cup P'_g$ is a path of H_0 from S_0 to T_0 , because it might pass through the same vertex more than once, and indeed P'_1, \dots, P'_g need not all be distinct. But concatenating P'_1, \dots, P'_g yields a walk from S_0 to T_0 , and therefore $P'_1 \cup \dots \cup P'_g$ includes a path P' from S_0 to T_0 . Define Q' similarly, with P replaced by Q . We may assume that no internal vertex of P' or of Q' belongs to $S_0 \cup T_0$.

(1) *We may assume that there exists $i \in \{1, \dots, n\}$ such that $v_i \in V(P') \cap V(Q')$.*

Since (H_0, r_0, S_0, T_0) is an $(\ell, 2)$ -block, and neither of P', Q' contains r_0 , it follows that $\text{dist}_{H_0}(P', Q') \leq 2$. Now there are two cases, depending whether P', Q' are vertex-disjoint or not. If P', Q' are vertex-disjoint, then since $\text{dist}_{H_0}(P', Q') \leq 2$, it follows that the corresponding path of H_0 joining them uses no dotted paths, and so is a path of J ; and hence $\text{dist}_G(P, Q) \leq 2$, and the result holds. So we may assume that some vertex $x \in V(P') \cap V(Q')$. If $x \in V(J)$, then $x \in V(P) \cap V(Q)$ and the result holds, so we may assume that $x \notin V(J)$. It remains to show that we may choose x to be an anchor of H_0 . If not, then x belongs to the interior of some dotted path of H_0 ; but then the whole of that path belongs to $P' \cap Q'$, and one end of the path is an anchor. This proves (1).

(2) *We may assume that there exists $i \in \{1, \dots, n-1\}$ such that $B_i \subseteq P' \cap Q'$.*

By (1) we may assume that $v_i \in V(P' \cap Q')$ for some $i \in \{1, \dots, n\}$. Suppose first that $i = 1$. Thus v_1 is the first vertex of both P', Q' . Neither of P', Q' contain s_0 , since no internal vertex of P' or of Q' belongs to $S_0 \cup T_0$; so both P', Q' contain B_1 and the claim holds. Similarly the claim holds if $i = n$, so we assume that $2 \leq i \leq n-1$. Now both P', Q' include two of the three dotted paths of H_0 incident with v_i , and so include one in common. If that one is a spine, then its end in $V(J)$ belongs to $P' \cap Q'$ and we are done as before; and otherwise one of $B_i, B_{i-1} \in P' \cap Q'$. This proves (2).

Consequently both P, Q have subpaths with one end in V_i , one end in V_{i+1} , and all internal vertices in H_i . Since (H_i, r_0, V_i, V_{i+1}) is an (ℓ, m) -block, and neither of P, Q contains r_0 , it follows that $\text{dist}_{H_i}(P, Q) \leq 2$, and hence $\text{dist}_G(P, Q) \leq 2$. This proves 2.1. \blacksquare

Next we need to show:

2.2 *If (G, r_0, S, T) is an $(\ell, m+1)$ -block, and $X \subseteq V(G)$ with $|X| \leq m$, then there is a path P of G between S, T with $\text{dist}_G(P, X \cup \{r_0\}) > \ell$.*

Proof. Again, we may assume the result holds for (ℓ, m) -blocks, and use the notation of the construction. For each $x \in X$, let C_x be the set of vertices v of G with $\text{dist}_G(v, x) \leq \ell$, let $C = \bigcup_{x \in X} C_x$, and let C_0 be the set of all v with $\text{dist}_G(v, r_0) \leq \ell$. We need to show that there is an $S - T$ path disjoint from $C \cup C_0$. In fact we will prove that there is such a path within the union of the H_i 's and the spines. In Figure 3 we show a section of the union of the H_i 's and spines. Let us label the vertices of J that are ends of spines as in the figure. (Thus, $s_0 = u_1$, and $t_0 = u_{n-1}$.)




Figure 3: Part of the construction for an $(\ell, m + 1)$ -block. The dotted curves represent paths of length $2\ell + 1$. Each gray area (together with r_0) is an (ℓ, m) -block, and really has m (not four as in the figure) vertices at either end. Each contains neighbours of r_0 (not shown in the figure.)

We may assume that $m \geq 2$, and so every vertex in $S \cup T$ has degree at least three in G . We begin with:

- (1) *We may assume that for every path R of G of length at least $2\ell + 1$ in which all internal vertices have degree two in G , no internal vertex of R belongs to X .*

If $x \in X$ is an internal vertex of such a path R , then by extending R as much as possible, we may assume that both ends of R have degree at least three. Let R have ends a, b . Since R has length at least $2\ell + 1$, not all of $V(R)$ belongs to C_x , and we may assume that some vertex of R between x, b is not in C_x . Let $X' = (X \setminus \{x\}) \cup \{a\}$. Every path of G within distance ℓ of x , and with no end in the interior of R , is within distance ℓ of a . So if the result is true for X' then it is true for X , so by repeating this at most $|X|$ times, we may assume there is no such x . This proves (1).

- (2) *We may assume that for some $i \in \{1, \dots, n - 1\}$, every path of $H_i \setminus r_0$ between V_i, V_{i+1} intersects C .*

Suppose not, and for $1 \leq i \leq n - 1$ let P_i be a path of $H_i \setminus r_0$ between V_i, V_{i+1} that is vertex-disjoint from C . For $2 \leq i < n$, the end of P_{i-1} in V_i and the end of P_i in V_i are joined by a path (Q_i say) of H_i of length $2\ell + 1$ (or zero, if the two ends are equal) and all its internal vertices have degree two, so by (1), no vertex of Q_i belongs to X . Since its ends are not in C , no vertex of Q_i is in C ; and

$$P_1 \cup Q_2 \cup P_2 \cup Q_2 \cup \dots \cup Q_{n-1} \cup P_{n-1}$$

is a path satisfying the theorem. This proves (2).

Let i be as in (2). For each $x \in X$, C_x contains at most one vertex in $V_i \cup V_{i+1}$, since they pairwise have distance at least $2\ell + 1$. Consequently $C \cap V(H_i)$ is the union of at most $|X|$ balls of H_i of radius ℓ . But from the inductive hypothesis, for any set of at most $m - 1$ such balls, there is a path of $H_i \setminus r_0$ between V_i, V_{i+1} avoiding their union: so $C_x \cap V(H_i) \neq \emptyset$ for each $x \in X$. It follows that $X \subseteq V(H_{i-1}) \cup V(H_i) \cup V(H_{i+1})$ (defining H_0, H_n to be null), since no vertex of X is in a spine by (1). Moreover, $V_{i-1} \cap C = \emptyset$ (again, defining $V_0, V_{n+1} = \emptyset$), since $C_x \cap V(H_i) \neq \emptyset$ for each $x \in X$; and similarly $V_{i+2} \cap C = \emptyset$. If i is odd, the vertex called u_i (see Figure 3) exists, and provides a way to avoid C , via two spines incident with u_i ; so we assume that i is even. Thus H_{i-1}, H_{i+1} exist. We may assume that C does not contain V_i (because if it does, then $i - 1$ also satisfies (2) and is odd). Since $C_x \cap V(H_i) \neq \emptyset$ for each $x \in X$, and each of the sets $C_x \cap V_i$ ($x \in X$) as size at most one, it follows that $C \cap V(H_{i-1})$ is the union of at most $|C \cap V_i| \leq |V_i| - 1 = m - 1$ balls of radius ℓ , and so from the inductive hypothesis, there is a path of $H_{i-1} \setminus \{r_0\}$ between V_{i-2}, V_{i-1} that is disjoint from C . But then we can extend this path via two spines incident with u_{i+1} to provide a route that avoids C . This proves 2.2. ■

From 2.1 and 2.2, it follows that $(\ell, m - 1)$ -blocks provide a counterexample to 1.2 for each value of $\ell, m \geq 1$, and so this proves 1.3.

3 What remains?

In view of this counterexample, is there anything even weaker that might be true and provide some sort of coarse extension of Menger's theorem? There are several possibilities to consider:

Bounded degree

Imposing an upper bound on the maximum degree looks promising at first, because Gartland, Korhonen and Lokshantov [4] and Hendrey, Norin, Steiner, and Turcotte [6] proved it with $c = 2$: they proved

3.1 *For every integer $\Delta \geq 1$ there exists $C > 0$ with the following property. Let G be a graph, let $k \geq 1$ be an integer, and let $S, T \subseteq V(G)$; then either*

- *there are k paths between S, T , pairwise at distance at least two; or*
- *there is a set $X \subseteq V(G)$ with $|X| < kC$ such that every path between S, T contains a vertex of X .*

But when $c \geq 3$, we can nearly make a counterexample. In the definition of an $(\ell, m + 1)$ -block, there are currently vertices of large degree: the vertices of J that are the ends of spines, and the root, and the vertices in the sets V_i . It is easy to modify the construction to keep the vertices in V_i of degree at most three: attach a leaf to each vertex in $S \cup T$, and call these new vertices S and T instead. (When we chain them together to make an $(\ell, m + 1)$ -block, they become of degree three instead of degree one, but that is fine.) For a vertex $u \in V(J)$ that is an end of spines, it is (in general) currently incident with $2m$ spines, and one edge of J . But we can partition these spines into

two groups in the natural way, and replace each group with a binary tree with root u and leaves the corresponding set V_i ; so that problem goes away as well. Thus, the only problem is the root vertex. In summary, we can make a counterexample to 1.2 in which only one vertex has degree more than three, but we don't see how to fix this last vertex.

Bounded pathwidth

Our counterexamples contain arbitrarily large binary trees, and so have unbounded pathwidth. We will prove in a later paper [10] that not only 1.2 but also 1.1 is true in graphs of bounded pathwidth:

3.2 *For all $c, k, w \geq 1$, there exists ℓ such that if G has pathwidth at most w , and $S, T \subseteq V(G)$, then either:*

- *there are k paths between S, T , pairwise at distance at least c ; or*
- *there is a set $X \subseteq V(G)$ with $|X| \leq k - 1$ such that every path between S, T contains a vertex with distance at most ℓ from some member of X .*

Our counterexample to 1.2 contains subdivisions of arbitrarily large complete graphs, and indeed contains them with arbitrarily large “fatness” (see [5] for definition). Is 1.2 true for graphs of bounded tree-width say? Or for every proper minor-closed class?

Planar graphs

That suggests that we study whether 1.2 is true for planar graphs, but in that case, it seems possible that 1.1 itself is true. Indeed, 1.1 might be true for graphs of bounded genus: our counterexample to 1.1 in Figure 1 has unbounded genus, because it contains arbitrarily many vertex-disjoint nonplanar subgraphs.

We have begun to work on trying to prove 1.1 for planar graphs, and have been able to prove it in the case when the graph is planar and all the vertices in $S \cup T$ are incident with the infinite region [11]. (Despite its appearance, that was difficult!) We think we can also prove it in the cylinder case, when the graph is planar and some two regions include $S \cup T$: and perhaps there is an approach to the general coarse Menger conjecture for planar graphs that is like that in [12, 13]. But this needs much further work.

References

- [1] S. Albrechtsen, T. Huynh, R. W. Jacobs, P. Knappe, and P. Wollan, “A Menger-type theorem for two induced paths”, *SIAM Journal on Discrete Mathematics* **38** (2024), 1438–1450, [arXiv:2305.04721v5](#).
- [2] J. Baligács and J. MacManus, “The metric Menger problem”, [arXiv:2403.05630](#).
- [3] D. Bienstock, “On the complexity of testing for odd holes and induced odd paths”, *Discrete Math.* **90** (1991), 85–92 (see also D. Bienstock, “Corrigendum: On the complexity of testing for odd holes and induced odd paths”, *Discrete Math.* **102** (1992), 109).

- [4] P. Gartland, T. Korhonen and D. Lokshtanov, “On induced versions of Menger’s theorem on sparse graphs”, [arXiv:2309.08169](#).
- [5] A. Georgakopoulos and P. Papasoglu, “Graph minors and metric spaces”, [arXiv:2305.07456](#).
- [6] K. Hendrey, S. Norin, R. Steiner, and J. Turcotte, “On an induced version of Menger’s theorem”, *Electronic J. Combinatorics* **31**, #P4.28, [arXiv:2309.07905](#).
- [7] K. Kawarabayashi and Y. Kobayashi, “The induced disjoint paths problem”, *Integer Programming and Combinatorial Optimization: 13th International Conference, IPCO 2008*, Bertinoro, Italy, May 26-28, 2008, Lecture Notes in Computer Science **5035**, Springer Berlin Heidelberg, 2008.
- [8] K. Menger, “Zur allgemeinen kurventheorie”, *Fundamenta Mathematicae* **10** (1927), 96–115.
- [9] T. Nguyen, A. Scott and P. Seymour, “A counterexample to the coarse Menger conjecture”, *J. Combinatorial Theory, Ser. B*, **173** (2025), 58–82. [arXiv:2401.06685](#).
- [10] T. Nguyen, A. Scott and P. Seymour, “Asymptotic structure. V. The coarse Menger conjecture is true if pathwidth is bounded”, in preparation.
- [11] T. Nguyen, A. Scott and P. Seymour, “Asymptotic structure. VI. Distant paths across a disc”, in preparation.
- [12] N. Robertson and P. Seymour, “Graph minors. VI. Disjoint paths across a disc”, *J. Combinatorial Theory, Ser. B*, **41** (1986), 115–138.
- [13] N. Robertson and P. Seymour, “Graph minors. VII. Disjoint paths on a surface”, *J. Combinatorial Theory, Ser. B*, **45** (1988), 212–254.