

Pure pairs. V. Excluding some long subdivision

Alex Scott

Mathematical Institute, University of Oxford, Oxford OX2 6GG, UK

Paul Seymour¹

Princeton University, Princeton, NJ 08544

Sophie Spirkl²

Princeton University, Princeton, NJ 08544

December 13, 2019; revised December 8, 2020

¹Supported by AFOSR grant A9550-19-1-0187 and NSF grant DMS-1800053.

²This material is based upon work supported by the National Science Foundation under Award No. DMS-1802201.

Abstract

A “pure pair” in a graph G is a pair A, B of disjoint subsets of $V(G)$ such that A is complete or anticomplete to B . Jacob Fox showed in [1] that for all $\varepsilon > 0$, there is a comparability graph G with n vertices, where n is large, in which there is no pure pair A, B with $|A|, |B| \geq \varepsilon n$. He also proved that for all $c > 0$ there exists $\varepsilon > 0$ such that for every comparability graph G with $n > 1$ vertices, there is a pure pair A, B with $|A|, |B| \geq \varepsilon n^{1-c}$; and conjectured that the same holds for every perfect graph G . We prove this conjecture and strengthen it in several ways.

In particular, we show that for all $c > 0$, and all $\ell_1, \ell_2 \geq 4c^{-1} + 9$, there exists $\varepsilon > 0$ such that, if G is an $(n > 1)$ -vertex graph with no hole of length exactly ℓ_1 and no antihole of length exactly ℓ_2 , then there is a pure pair A, B in G with $|A| \geq \varepsilon n$ and $|B| \geq \varepsilon n^{1-c}$. This is further strengthened, replacing excluding a hole by excluding some “long” subdivision of a general graph.

1 Introduction

Graphs in this paper are finite, and without loops or parallel edges. Let $A, B \subseteq V(G)$ be disjoint. We say that A is *complete* to B , or A, B are *complete*, if every vertex in A is adjacent to every vertex in B , and similarly A, B are *anticomplete* if no vertex in A has a neighbour in B . We say A *covers* B if every vertex in B has a neighbour in A . A *pure pair* in G is a pair A, B of disjoint subsets of $V(G)$ such that A, B are complete or anticomplete.

Jacob Fox [1] proved:

1.1 *For every sufficiently large positive integer n :*

- *for every n -vertex comparability graph G , there is a pure pair A, B in G with $|A|, |B| > \frac{n}{4 \log_2 n}$;*
- *there is an n -vertex comparability graph G such that there is no pure pair A, B in G with $|A|, |B| \geq \frac{15n}{\log_2 n}$.*

There is also a slightly stronger asymmetric result, by Fox, Pach and Toth [2]:

1.2 *There exists $\varepsilon > 0$ such that for every n -vertex comparability graph G with $n > 1$, either there is a complete pair A, B with $|A|, |B| \geq cn$, or there is an anticomplete pair A, B with $|A|, |B| \geq cn / \log n$.*

Comparability graphs are perfect, and Fox conjectured that something like 1.1 holds for all perfect graphs; more exactly:

1.3 Conjecture: *For every sufficiently large positive integer n and every n -vertex perfect graph G , there is a pure pair A, B in G with $|A|, |B| \geq n^{1-o(1)}$.*

We will prove this conjecture, and several strengthenings. To prove 1.3 itself, we will show that

1.4 *For all $c > 0$, and all sufficiently large n , if G is an n -vertex perfect graph, then there is a pure pair A, B in G with $|A|, |B| \geq n^{1-c}$.*

We denote the number of vertices of a graph G by $|G|$. We can replace the “sufficiently large” condition in 1.4 with a multiplicative constant; 1.4 is equivalent to:

1.5 *For all $c > 0$ there exists $\varepsilon > 0$ such that if G is a perfect graph with $|G| > 1$, then there is a pure pair A, B in G with $|A|, |B| \geq \varepsilon |G|^{1-c}$.*

This can be strengthened; we will show:

1.6 *For all $c > 0$ there exists $\varepsilon > 0$ such that if G is a perfect graph with $|G| > 1$, then there is a pure pair A, B in G with $|A| \geq \varepsilon |G|$ and $|B| \geq \varepsilon |G|^{1-c}$.*

The complement graph of G is denoted by \overline{G} . A *hole* in G is an induced cycle of length at least four; and an *antihole* in G is an induced subgraph whose complement graph is a hole in \overline{G} . Perfect graphs have no holes or antiholes of odd length, but we will show that it is not necessary to exclude all odd holes and odd antiholes to have the result 1.6; it is enough to exclude one of each, of sufficient length. The next result is a strengthening of 1.6:

1.7 Let $c > 0$ with $1/c$ an integer, and let $\ell_1, \ell_2 \geq 4c^{-1} + 5$ be integers. Then there exists $\varepsilon > 0$ such that if G is a graph with $|G| > 1$, with no hole of length ℓ_1 and no antihole of length ℓ_2 , then there is a pure pair A, B in G with $|A| \geq \varepsilon|G|$ and $|B| \geq \varepsilon|G|^{1-c}$.

This can be further strengthened, as follows. Let us say G contains H if some induced subgraph of H is isomorphic to H , and G is H -free otherwise. If $X \subseteq V(G)$, $G[X]$ denotes the subgraph induced on X . We say that a graph H has *branch-length* at least ℓ if every cycle of H has length at least ℓ , and every two vertices of H with degree at least three have distance at least ℓ in H . Since a cycle of length ℓ_1 has branch-length ℓ_1 , the next result strengthens 1.7 and is the main result of the paper:

1.8 Let $c > 0$ with $1/c$ an integer, and let H_1, H_2 be graphs with branch-length at least $4c^{-1} + 5$. Then there exists $\varepsilon > 0$ such that if G is a graph with $|G| > 1$ that is H_1 -free and $\overline{H_2}$ -free, then there is a pure pair A, B in G with $|A| \geq \varepsilon|G|$ and $|B| \geq \varepsilon|G|^{1-c}$.

We have found a kind of strengthening of 1.8, that we state without proof (we will not use it). For $\ell \geq 2$, Let us say a graph H is ℓ -handled if there are induced subgraphs P_0, \dots, P_k of H , for some $k \geq 1$, such that:

- P_0 is a forest;
- every path of P_0 has length at most ℓ ;
- P_1, \dots, P_k are pairwise vertex-disjoint paths, each of length at least ℓ ;
- for $1 \leq i \leq k$, $V(P_i \cap P_0)$ consists exactly of the two ends of P_i ; and
- $H = P_0 \cup P_1 \cup \dots \cup P_k$.

Then:

1.9 There exists $\gamma > 0$ with the following property. Let $c > 0$ with $1/c$ an integer, and let H_1, H_2 be γc^{-1} -handled graphs. Then there exists $\varepsilon > 0$ such that if G is a graph with $|G| > 1$ that is H_1 -free and $\overline{H_2}$ -free, then there is a pure pair A, B in G with $|A| \geq \varepsilon|G|$ and $|B| \geq \varepsilon|G|^{1-c}$.

Then the essentials of 1.8 follow from 1.9 by taking P_0 to be the subgraph of H induced on the set of all vertices of degree at least three and their neighbours. But we feel that 1.9 is not very satisfactory, because if the forest P_0 has long paths, the hypothesis requires the paths P_1, \dots, P_k to be long too. We would prefer a version of 1.9 where we omit the second bullet from the definition of exterior-width, but so far we cannot prove it.

A weaker form of 1.8 will be proved for a wider class of graphs in [5]. Let H be a graph. If $E(H) \neq \emptyset$, we define the *congestion* of H to be the maximum of $1 - (|J| - 1)/|E(J)|$, taken over all subgraphs J of H with at least one edge; and if $E(H) = \emptyset$, we define the congestion of H to be zero. Thus the congestion of H is always non-negative, and equals zero if and only if H is a forest; and, for instance, long cycles have smaller congestion than short cycles.

In [5] we will prove:

1.10 Let $c > 0$, and let H_1, H_2 be graphs with congestion at most $c/(9 + 15c)$. Then there exists $\varepsilon > 0$ such that if G is a graph with $|G| > 1$ that is H_1 -free and $\overline{H_2}$ -free, then there is a pure pair A, B in G with $|A|, |B| \geq \varepsilon|G|^{1-c}$.

This is pleasing because of the following weak converse (easily proved with a random graph argument that we omit):

1.11 *Let $c > 0$, and let H_1, H_2 be graphs both with congestion more than c . There is no $\varepsilon > 0$ such that for every graph G with $|G| > 1$ that is H_1 -free and $\overline{H_2}$ -free, there is a pure pair A, B in G with $|A|, |B| \geq \varepsilon|G|^{1-c}$.*

The result 1.10 does not contain 1.8, because in 1.10 neither of A, B have to have linear cardinality. What if we ask for a strengthened version of 1.10 that would contain 1.8? We pose that as a conjecture:

1.12 Conjecture: *For all $c > 0$, there exists $\sigma > 0$ with the following property. Let H_1, H_2 be graphs with congestion at most σ . There exists $\varepsilon > 0$ such that if G is a graph with $|G| > 1$ that is H_1 -free and $\overline{H_2}$ -free, then there is a pure pair A, B in G with $|A| \geq \varepsilon|G|$ and $|B| \geq \varepsilon|G|^{1-c}$.*

2 Reduction to the sparse case

Let us say a graph G is ε -sparse if every vertex has degree less than $\varepsilon|G|$. We say G is (α, β) -coherent if there do not exist disjoint subsets A, B of $V(G)$, anticomplete to each other, such that $|A| \geq \alpha$ and $|B| \geq \beta$.

A one-vertex graph is ε -sparse for all $\varepsilon > 0$, and (α, β) -coherent for all $\alpha, \beta > 0$, so our standard hypothesis that G is suitably coherent and suitably sparse does not exclude the case $|G| = 1$, and we always need to assume separately that $|G| > 1$. But we observe:

2.1 *If $c, \varepsilon > 0$, and $\varepsilon \leq 1/2$, and G is ε -sparse and $(\varepsilon|G|^{1-c}, \varepsilon|G|)$ -coherent, and $|G| > 1$, then $|G| > 1/\varepsilon$.*

Proof. Suppose that $|G| \leq 1/\varepsilon$. If some distinct $u, v \in V(G)$ are non-adjacent, $\{u\}, \{v\}$ form an anticomplete pair, both of cardinality at least $\varepsilon|G|$, a contradiction. So G is a complete graph; but its maximum degree is less than $\varepsilon|G|$ and $\varepsilon \leq 1/2$, which is impossible since $|G| > 1$. This proves 2.1. ■

If G is a graph and $v \in V(G)$, a G -neighbour of v means a vertex of G adjacent to v in G . A theorem of Rödl [4] implies the following:

2.2 *For every graph H and all $\eta > 0$ there exists $\delta > 0$ with the following property. Let G be an H -free graph. Then there exists $X \subseteq V(G)$ with $|X| \geq \delta|G|$, such that one of $G[X], \overline{G}[X]$ is η -sparse.*

Consequently, in order to prove 1.8, it suffices to prove the following:

2.3 *Let $c > 0$ with $1/c$ an integer, and let H be a graph with branch-length at least $4c^{-1} + 5$. Then there exists $\varepsilon > 0$ such that every ε -sparse $(\varepsilon|G|^{1-c}, \varepsilon|G|)$ -coherent graph G with $|G| > 1$ contains H .*

Proof of 1.8, assuming 2.3. Let $c > 0$ with $1/c$ an integer, and let H_1, H_2 have branch-length at least $4c^{-1} + 5$. For $i = 1, 2$, choose ε_i such that 2.3 holds with $\varepsilon = \varepsilon_i$ and $H = H_i$. Let $\eta = \min(\varepsilon_1, \varepsilon_2, 1/2)$. Choose δ such that 2.2 holds taking $H = H_1$. Let $\varepsilon = \eta\delta$. We claim that ε satisfies 1.8.

Let G be a graph with $|G| > 1$ that is H_1 -free and $\overline{H_2}$ -free. We must show that there is a pure pair A, B in G with $|A| \geq \varepsilon|G|$ and $|B| \geq \varepsilon|G|^{1-c}$. From the choice of δ , there exists $X \subseteq V(G)$ with $|X| \geq \delta|G|$, such that one of $G[X], \overline{G}[X]$ is η -sparse; and by 2.1 we may assume that $|G| > 1/\varepsilon \geq \delta$, and so $|X| > 1$. In the first case, since $\eta \leq \varepsilon_1$, 2.3 applied to $G[X]$ implies that there is an anticomplete pair A, B in $G[X]$ with $|A| \geq \eta|X|$ and $|B| \geq \eta|X|^{1-c}$. Thus

$$|A| \geq \eta|X| \geq \eta\delta|G| = \varepsilon|G|$$

and

$$|B| \geq \eta|X|^{1-c} \geq \eta\delta^{1-c}|G|^{1-c} \geq \eta\delta|G|^{1-c} = \varepsilon|G|^{1-c},$$

as required. In the second case we argue similarly, working in $\overline{G}[X]$. This proves 1.8. ▀

The remainder of the paper is devoted to proving 2.3.

3 Finding a path of specified length

A *levelling* in G is a sequence (L_0, L_1, \dots, L_k) of disjoint subsets of $V(G)$ with $k \geq 1$ such that

- $|L_0| = 1$;
- L_{i-1} covers L_i for $1 \leq i \leq k$; and
- $L_0 \cup \dots \cup L_{i-2}$ is anticomplete to L_i for all $i \in \{2, \dots, k\}$.

We denote $L_0 \cup L_1 \cup \dots \cup L_k$ by $V(\mathcal{L})$. We call L_k the *base* of the levelling $\mathcal{L} = (L_0, L_1, \dots, L_k)$, and $V(\mathcal{L}) \setminus L_k$ is called the *heart* of \mathcal{L} . We call k the *height* of the levelling, and the unique vertex in L_0 is the *apex*. We call L_{k-1} the *penultimate term* of the levelling (for want of a better name). A path P is \mathcal{L} -*vertical* if $V(P) \subseteq V(\mathcal{L})$ and $|V(P) \cap L_i| \leq 1$ for $0 \leq i \leq k$.

We need the following lemma:

3.1 *Let $\rho \geq 1$ be some real number, let $K, k > 0$ be integers with $K > k$, and let n_1, \dots, n_K be non-negative integers, all less than $\rho^{K/k-2-1/k}$. Then there exists $i \in \{1, \dots, K-k\}$ such that $\rho n_i \geq n_j$ for $j = i+1, \dots, i+k$.*

Proof. Suppose not; then for each $i \in \{1, \dots, K-k\}$ there exists $f(i)$ such that $i < f(i) \leq i+k$ and $\rho n_i < n_{f(i)}$. Define $x_1 = 1$ and $x_{i+1} = f(x_i)$ provided $x_i \leq K-k$. Let x_1, \dots, x_t be defined by this process; thus $K-k < x_t \leq K$. Since $x_{i+1} - x_i \leq k$ for each i , it follows that $tk \geq K-1$. Since $n_{x_2} > \rho n_{x_1}$ and n_{x_2} is an integer, it follows that $n_{x_2} \geq 1$. Thus for $2 \leq i \leq t$, $n_{x_i} \geq \rho^{i-2}$, and so $n_{x_t} \geq \rho^{t-2} \geq \rho^{K/k-2-1/k}$, contrary to the hypothesis. This proves 3.1. ▀

Next we need:

3.2 *Let $c > 0$ such that c^{-1} is an integer, and define $r = 1 + 1/c$. Let $\ell \geq 1$ be an integer, and define $K = r^\ell - 1$, and $k = r^{\ell-1} - 1$. Let $\varepsilon > 0$, and let G be an ε -sparse ($\varepsilon|G|^{1-c}, \varepsilon|G|$)-coherent graph. Let $B_0, B_1, \dots, B_K \subseteq V(G)$ be disjoint, where $B_0 \neq \emptyset$ and B_1, \dots, B_K each have cardinality at least $r^{2\ell}\varepsilon|G|$. Then either:*

- there is an induced path of length ℓ , with vertices p_0, p_1, \dots, p_ℓ in order, and

$$1 \leq t_1 < t_2 < \dots < t_\ell \leq K,$$

such that $p_0 \in B_0$, and $p_i \in B_{t_i}$ for $1 \leq i \leq \ell$; or

- $|B_0| \leq K\varepsilon|G|^{1-c}$, and there are sets C_1, \dots, C_{K-k} with union B_0 , such that for each i with $1 \leq i \leq K-k$, and each j with $i \leq j \leq i+k$, at least $r^{2\ell-2}\varepsilon|G|$ vertices in B_j have no neighbour in C_i .

Proof. We proceed by induction on ℓ . Suppose first that $\ell = 1$. If there is an edge between B_0 and $B_1 \cup \dots \cup B_K$, then the first bullet holds; and if B_0 is anticomplete to $B_1 \cup \dots \cup B_K$, then since H is $(\varepsilon|G|^{1-c}, \varepsilon|G|)$ -coherent and $|B_1| \geq r^{2\ell}\varepsilon|G| \geq \varepsilon|G|$, it follows that $|B_0| < \varepsilon|G|^{1-c}$, and the second bullet holds, taking $C_1, \dots, C_{K-k} = B_0$. Thus we may assume that $\ell \geq 2$ and the result holds for $\ell - 1$. Define $\rho = |G|^c$.

Let $B_0 = \{v_1, \dots, v_n\}$. For $1 \leq i \leq K-k$ and $i \leq j \leq i+k$, define $A_{i,j}^0 = \emptyset$, and $A^0 = \emptyset$. Inductively for $h = 1, \dots, n$ we will define

- the sets $X_i^h \subseteq B_i$ for $1 \leq i \leq K$;
- the *type* of v_h (one of the numbers $1, \dots, K-k$);
- the sets $A_{i,j}^h \subseteq B_j$ for $1 \leq i \leq K-k$ and $i \leq j \leq i+k$; and
- the set A^h , which is the union of $A_{i,j}^h$ over all $i \in \{1, \dots, K-k\}$ and all $j \in \{i, \dots, i+k\}$

as follows. Suppose that $1 \leq h \leq n$, and A^{h-1} and $A_{i,j}^{h-1}$ are defined for all i, j with $1 \leq i \leq K-k$ and $i \leq j \leq i+k$. For $1 \leq i \leq K$ let X_i^h be the set of vertices in $B_i \setminus A^{h-1}$ adjacent to v_h . Since $(K+1)/(k+1) = 2 + 1/c$, it follows that $K > (2 + 1/c)k + 1$, and so $1/c < K/k - 2 - 1/k$. Hence for $1 \leq i \leq K$, $|X_i^h| \leq |G| < \rho^{K/k-2-1/k}$. By 3.1 applied to the numbers $|X_1^h|, \dots, |X_K^h|$, there exists i with $1 \leq i \leq K-k$ such that $\rho|X_i^h| \geq |X_j^h|$ for $j = i, \dots, i+k$. Choose some such i . We define i to be the type of v_h . For $i \leq j \leq i+k$ define $A_{i,j}^h = A_{i,j}^{h-1} \cup X_j^h$. This completes the inductive definition.

(1) $\rho|A_{i,j}^h| \geq |A_{i,i}^h|$ for all $h \in \{1, \dots, n\}$ and all $i \in \{1, \dots, K-k\}$ and all $j \in \{i, \dots, i+k\}$.

$A_{i,j}^h$ is the disjoint union of the sets X_j^h for all $h \in \{1, \dots, n\}$ such that v_h has type i ; and $A_{i,i}^h$ is the disjoint union of X_i^h for the same values of h . Since $\rho|X_i^h| \geq |X_j^h|$ for each such h , this proves (1).

(2) If v_h has type i , then every vertex of B_j adjacent to v_h belongs to A^h , for all $h \in \{1, \dots, n\}$, all $1 \leq i \leq K-k$, and all $j \in \{i, \dots, i+k\}$.

Let $x \in B_j$ be adjacent to v_h . If $x \in A^{h-1}$, then the claim holds since $A^{h-1} \subseteq A^h$. If $x \notin A^{h-1}$ then $x \in X_j^h$ from the definition of X_j^h , and since v_h has type i , it follows that

$$x \in X_j^h \subseteq A_{i,j}^h \subseteq A^h.$$

This proves (2).

For $1 \leq i \leq K - k$, let C_i be the set of vertices in B_0 that have type i . Thus C_1, \dots, C_{K-k} are pairwise disjoint and have union B_0 . We note that

$$r^{2\ell} - r^{2\ell-2} = (3 + 4/c + 1/c^2)(k+1)^2 \geq 2(k+1)^2.$$

(3) We may assume that $|A_{i,j}^n| > k\varepsilon|G|$ for some $i \in \{1, \dots, K - k\}$ and some $j \in \{i, \dots, i + k\}$.

Suppose not. Let $1 \leq j \leq K$. Since $A^n \cap B_j$ is the union of the sets $A_{i,j}^n$ for all $i \in \{1, \dots, K\}$ with $j - i \in \{0, \dots, k\}$, it follows that $|A^n \cap B_j| \leq k(k+1)\varepsilon|G|$. Now let $1 \leq i \leq K - k$; by (2), C_i is anticomplete to $B_j \setminus A^n$, for all $j \in \{i, \dots, i + k\}$. Since

$$|B_j \setminus A^n| = |B_j| - |B_j \cap A^n| \geq r^{2\ell}\varepsilon|G| - k(k+1)\varepsilon|G| \geq r^{2\ell-2}\varepsilon|G| \geq \varepsilon|G|$$

and G is $(\varepsilon|G|^{1-c}, \varepsilon|G|)$ -coherent, it follows that $|C_i| < \varepsilon|G|^{1-c}$. Hence $|B_0| \leq K\varepsilon|G|^{1-c}$; and so the second bullet of the theorem holds. This proves (3).

From (3), we may choose $h \in \{1, \dots, n\}$ minimum such that $|A_{i,j}^h| > k\varepsilon|G|$ for some $i \in \{1, \dots, K - k\}$ and some $j \in \{i, \dots, i + k\}$. Define D to be the set of all $v_{h'} \in C_i$ with $1 \leq h' \leq h$. From the minimality of h , and since G is ε -sparse, it follows that $|A_{i,j}^h| \leq (k+1)\varepsilon|G|$ for all $i \in \{1, \dots, K - k\}$ and all $j \in \{i, \dots, i + k\}$. Consequently $|A^h \cap B_i| \leq (k+1)^2\varepsilon|G|$ for $1 \leq i \leq K$. Now choose $i \in \{1, \dots, K - k\}$ such that $|A_{i,j}^h| > k\varepsilon|G|$ for some $j \in \{i, \dots, i + k\}$. By (1), $|A_{i,i}^h| > k\varepsilon|G|/\rho = k\varepsilon|G|^{1-c}$. For $j = i + 1, \dots, i + k$, let D_j be the set of all vertices in B_j that have no neighbour in D . Thus $B_j \setminus A^h \subseteq D_j$, and so

$$|D_j| \geq r^{2\ell}\varepsilon|G| - (k+1)^2\varepsilon|G| \geq r^{2\ell-2}\varepsilon|G|.$$

Since $|A_{i,i}^h| > k\varepsilon|G|^{1-c}$, it follows from the inductive hypothesis (with ℓ replaced by $\ell - 1$, and B_0 replaced by $A_{i,i}^h$, and B_1, \dots, B_K replaced by D_{i+1}, \dots, D_{i+k}) that there is an induced path of length $\ell - 1$, with vertices p_1, \dots, p_ℓ in order, and

$$i + 1 \leq t_2 < \dots < t_\ell \leq i + k,$$

such that $p_1 \in A_{i,i}^h$, and $p_i \in B_{t_i}$ for $2 \leq i \leq \ell$. Choose $p_0 \in D$ adjacent to p_1 , and define $t_1 = i$; then the path with vertices p_0, p_1, \dots, p_ℓ is induced and satisfies the first bullet of the theorem. This proves 3.2. ■

The main result of this section is the following:

3.3 Let $c > 0$, such that c^{-1} is an integer. Let $\ell, s, t > 0$ be integers, and let $d > 0$. Then there exists $\varepsilon > 0$ with the following property. Let G be an ε -sparse $(\varepsilon|G|^{1-c}, \varepsilon|G|)$ -coherent graph, and for $i = 1, 2$, let \mathcal{L}_i be a levelling in G , with heart H_i , apex a_i , and base B_i , satisfying:

- $V(\mathcal{L}_1) \cap V(\mathcal{L}_2) = \{a_1\} \cap \{a_2\}$;
- $V(\mathcal{L}_2) \setminus \{a_2\}$ is anticomplete to H_1 , and if $a_1 \neq a_2$ then a_2 is anticomplete to H_1 ;

- $\mathcal{L}_1, \mathcal{L}_2$ have heights s, t respectively; and
- $|B_i| \geq d|G|$ for $i = 1, 2$.

Then there is an induced path (or cycle, if $a_1 = a_2$) of length $\ell + s + t$ between a_1, a_2 , with vertex set a subset of $H_1 \cup B_1 \cup H_2 \cup B_2$.

Proof. For each integer $i \geq 0$, let $k_i = (2 + 1/c)^i - 1$. Choose ε such that $k_\ell k_{\ell+1} \dots k_{\ell+t-h} \varepsilon < d$ and $k_{\ell+t}(k_{2\ell+2t} + 2)\varepsilon \leq d$. Define $d_i = (2 + 1/c)^{2i} \varepsilon$ for each integer $i \geq 0$.

Let $\mathcal{L}_1 = (L_0, \dots, L_s)$ and $\mathcal{L}_2 = (M_0, \dots, M_t)$ say; thus $L_s = B_1$ and $M_t = B_2$. Let $Z_0 = \emptyset$. For $i = 1, \dots, k_{\ell+t}$, we will inductively define $Z_i \subseteq L_{s-1}$ with $X_{i-1} \subseteq Z_i$, and $D_i \subseteq L_s$ with D_1, \dots, D_i pairwise disjoint, satisfying

- $d_{\ell+t}|G| \leq |D_i| \leq (d_{\ell+t} + \varepsilon)|G|$
- D_i is the set of all vertices in L_s that have a neighbour in Z_i and have no neighbour in Z_{i-1} (and so $D_1 \cup \dots \cup D_i$ is the set of all vertices in L_s that have a neighbour in Z_i).

Thus, suppose that $1 \leq k_{\ell+t}$, and Z_0, \dots, Z_{i-1} and D_1, \dots, D_{i-1} are defined satisfying the conditions above. It follows that

$$|D_1 \cup \dots \cup D_{i-1}| \leq (i-1)(d_{\ell+t} + \varepsilon)|G| \leq (d - d_{\ell+t})|G|$$

and so at least $d_{\ell+t}|G|$ vertices in L_s do not belong to $|D_1 \cup \dots \cup D_{i-1}|$. All these vertices have a neighbour in $L_{s-1} \setminus Z_{i-1}$ and have no neighbour in Z_{i-1} ; and so there exists Z_i with $Z_{i-1} \subseteq Z_i \subseteq L_{s-1}$, minimal such that at least $d_{\ell+t}|G|$ vertices in L_s have a neighbour in Z_i and have none in Z_{i-1} . Let this set of vertices be D_i . Since G is ε -sparse, the minimality of Z_i implies that $|D_i| \leq (d_{\ell+t} + \varepsilon)|G|$. This completes the inductive definition.

Let $\mathcal{Q} = (Q_0, \dots, Q_t)$ be a levelling in G . We say it is a *sub-levelling* of \mathcal{L}_2 if $Q_i \subseteq M_i$ for $0 \leq i \leq t$. For $0 \leq h \leq t$, we say that such a sub-levelling $\mathcal{Q} = (Q_0, \dots, Q_t)$ is *h-good* if

- there exists $g \in \{1, \dots, k_{\ell+t} - k_{\ell+t-h} + 1\}$, and for each $j \in \{g, \dots, g + k_{\ell+t-h} - 1\}$ there exists $F_j \subseteq D_j$, such that F_j is anticomplete to $Q_0 \cup Q_1 \cup \dots \cup Q_{h-1}$, and $|F_j| \geq d_{\ell+t-h}|G|$; and
- $|Q_t| > k_\ell k_{\ell+1} \dots k_{\ell+t-h} \varepsilon |G|^{1-c}$.

Since $d|G| > k_\ell k_{\ell+1} \dots k_{\ell+t-h} \varepsilon |G|^{1-c}$ it follows that \mathcal{L}_2 is 0-good. Choose $h \leq t$ maximum such that some sub-levelling $\mathcal{Q} = (Q_0, \dots, Q_t)$ of \mathcal{L}_2 is *h-good*, and let g and the sets F_j ($j \in \{g, \dots, g + k_{\ell+t-h} - 1\}$) be as in the definition. Let $K = k_{\ell+t-h}$. Since each $|F_j| \geq d_{\ell+t-h}|G|$, we may apply 3.2, replacing B_0 by Q_h , and replacing ℓ by $\ell + t - h$, and replacing the sequence B_1, \dots, B_{k_ℓ} by F_g, \dots, F_{g+K-1} . There are two possible outcomes of 3.2.

The first outcome is: there is an induced path P of length $\ell + t - h$, with vertices $p_0, p_1, \dots, p_{\ell+t-h}$ in order, and

$$g \leq t_1 < t_2 < \dots < t_{\ell+t-h} \leq g + K - 1,$$

such that $p_0 \in Q_h$, and $p_i \in F_{t_i}$ for $1 \leq i \leq \ell + t - h$. In this case, choose a \mathcal{Q} -vertical path Q between a_2 and p_0 (therefore of length h); choose a neighbour v of $p_{\ell+t-h}$ in $Z_{t_{\ell+t-h}}$; and choose an \mathcal{L}_1 -vertical path R between a_1, v (therefore of length $s - 1$). We claim that

$$a_2-Q-p_0-p_1-p_{\ell+t-h}-v-R-a_1$$

is an induced path or cycle. To show this, we must check that

- $V(P) \cap V(Q) = \{p_0\}$, and $V(P) \setminus \{p_0\}$ is anticomplete to $V(Q) \setminus \{p_0\}$; this is true since Q_0, \dots, Q_{h-1} are anticomplete to F_g, \dots, F_{g+K-1} from the definition of h -good.
- $V(P) \cap V(R) = \emptyset$, and the edge with ends $p_{\ell+t-h}$ and v is the only edge between $V(P)$ and $V(R)$; this is true since L_0, \dots, L_{s-2} are anticomplete to L_s , and $v \in Z_{t_{\ell+t-h}}$ is anticomplete to $D_{t_1}, \dots, D_{t_{\ell+t-h-1}}$.
- $V(Q) \cap V(R) = \{a_1\} \cap \{a\}$, and every edge between $V(Q)$ and $V(R)$ has an end in $\{a_1\} \cap \{a\}$; this is true from the hypothesis.

This proves the path or cycle is indeed induced, and since it has length ℓ , the theorem holds.

The second outcome of 3.2 is: $\ell + t - h > 0$, and $|Q_h| \leq K\varepsilon|G|^{1-c}$, and, writing $k = k_{\ell+t-h-1}$, there are sets $C_g, \dots, C_{g+K-k-1}$ with union Q_h , such that for each i with $g \leq i \leq g+K-k-1$, and each j with $i \leq j \leq i+k$, at least $d_{\ell+t-h-1}|G|$ vertices in F_j have no neighbour in C_i . Since

$$|Q_h| \leq K\varepsilon|G|^{1-c} < k_\ell k_{\ell+1} \dots k_{\ell+t-h} \varepsilon |G|^{1-c} < |Q_t|$$

it follows that $h < t$. For $g \leq i \leq g+K-k-1$, let X_i be the set of vertices in Q_t that are joined to a vertex in C_i by a \mathcal{Q} -vertical path. Since \mathcal{Q} is a levelling and $C_g, \dots, C_{g+K-k-1}$ have union Q_h , it follows that $X_g, \dots, X_{g+K-k-1}$ have union Q_t ; and since $|Q_t| > k_\ell k_{\ell+1} \dots k_{\ell+t-h} \varepsilon |G|^{1-c}$, there exists i with $g \leq i \leq g+K-k-1$ such that

$$|X_i| \geq |Q_t|/K > k_\ell k_{\ell+1} \dots k_{\ell+t-h-1} \varepsilon |G|^{1-c}.$$

For $h \leq h' \leq t$ let $Q'_{h'}$ be the set of vertices in $Q_{h'}$ that are joined to a vertex in C_i by a \mathcal{Q} -vertical path. Thus $Q'_h = C_i$, and

$$(Q_0, \dots, Q_{h-1}, Q'_h, Q'_{h+1}, \dots, Q'_t)$$

is an $(h+1)$ -good sub-levelling of \mathcal{L}_2 , a contradiction. This proves 3.3. ▀

The next result is a form of 3.3 with the same hypotheses except that the bases of the two levellings might not be disjoint.

3.4 *Let $c > 0$, such that c^{-1} is an integer. Let $\ell, s, t > 0$ be integers, and let $d > 0$. Then there exist $\varepsilon > 0$ with the following property. Let G be an ε -sparse ($\varepsilon|G|^{1-c}, \varepsilon|G|$)-coherent graph. For $i = 1, 2$, let \mathcal{L}_i be a levelling in G , with heart H_i , apex a_i , and base B_i . Suppose that:*

- for $i = 1, 2$, $|B_i| \geq d|G|$;
- $V(\mathcal{L}_1) \cap V(\mathcal{L}_2) = (\{a_1\} \cap \{a_2\}) \cup (B_1 \cap B_2)$; and
- every edge between H_1 and $V(\mathcal{L}_2)$ has both ends in $V(\mathcal{L}_1)$.

Let $\mathcal{L}_1, \mathcal{L}_2$ have heights s, t respectively. Then there is an induced path (or cycle, if $a_1 = a_2$) of length $\ell + s + t$ between a_1, a_2 , with vertex set a subset of $H_1 \cup B_1 \cup H_2 \cup B_2$.

Proof. Given $d > 0$ let $d' = d/3$, and choose ε to satisfy 3.3 with d replaced by d' . We may assume that $\varepsilon \leq d'$ by reducing ε . We will show that ε satisfies the theorem. Let $G, \mathcal{L}_1, \mathcal{L}_2$ satisfy the hypotheses of the theorem, and let H_i, a_i, B_i ($i = 1, 2$) and s, t be as above. Let $\mathcal{L}_1 = (L_0, \dots, L_s)$; thus $L_s = B_1$. Choose $L'_{s-1} \subseteq L_{s-1}$ minimal such that at least $d'|G|$ vertices in B_1 have a neighbour in L'_{s-1} . Let L'_s be the set of vertices in L_s that have a neighbour in L'_{s-1} . Thus $d'|G| \leq |L'_s| \leq (d' + \varepsilon)|G| \leq 2d'|G|$.

Let \mathcal{L}'_1 be the levelling $(L_0, \dots, L_{s-1}, L'_{s-1}, L'_s)$. Let \mathcal{L}'_2 be the levelling obtained from \mathcal{L}_2 by replacing its base by $B_2 \setminus L'_s$. Then $|L'_s| \geq d'|G|$, and $|B_2 \setminus L'_s| \geq d|G| - 2d'|G| \geq d'|G|$. Hence $\mathcal{L}'_1, \mathcal{L}'_2$ satisfy the hypotheses of 3.3, and the result follows. This proves 3.4. \blacksquare

When we apply 3.4, in the final section, it will be to levellings $\mathcal{L}_1, \mathcal{L}_2$ such that the only edges between $V(\mathcal{L}_1), V(\mathcal{L}_2)$ are either incident with the common apex (if there is one) or between the base of one and one of the last two terms of the other; so 3.4 is stronger than we need.

4 Expansion

If $X \subseteq V(G)$, $N(X)$ denotes the set of vertices in $V(G) \setminus X$ with a neighbour in X , and $N[X] = N(X) \cup X$. A graph G is τ -*expanding* if $|N[X]| \geq \min(\tau|X|, |G|/2)$ for every subset $X \subseteq V(G)$.

4.1 *Let $c > 0$, and let G be a $(|G|^{1-c}/4, |G|/4)$ -coherent graph. Then there exists $Y \subseteq V(G)$ with $|Y| \leq |G|^{1-c}/4$ such that $G \setminus Y$ is $|G|^c$ -expanding.*

Proof. Let $\alpha = |G|^{1-c}/4$ and $\tau = |G|^c$. Choose $Y \subseteq V(G)$ maximal such that $|Y| \leq \alpha$ and $|N[Y]| \leq \tau|Y|$ (possibly $Y = \emptyset$). Let $W = V(G) \setminus Y$. If $G[W]$ is τ -expanding then the theorem holds, so we assume not. Thus there exists $X \subseteq W$ such that $|N[X] \cap W| < \min(\tau|X|, |W|/2)$. Consequently $X \neq \emptyset$. But

$$|N[X \cup Y]| \leq |N[Y]| + |N[X] \cap W| \leq \tau|Y| + \tau|X|,$$

and so from the maximality of Y , it follows that $|X \cup Y| > \alpha$. Now $|N[Y]| \leq \tau|Y| \leq \tau\alpha = |G|/4$, and $|N[X] \cap W| \leq |W|/2 \leq |G|/2$; so

$$|N[X \cup Y]| \leq |N[Y]| + |N[X] \cap W| \leq 3|G|/2.$$

Let $U = V(G) \setminus N[X \cup Y]$; then $|U| \geq |G|/4$. But $X \cup Y$ is anticomplete to U , contradicting that G is $(|G|^{1-c}/4, |G|/4)$ -coherent. This proves 4.1. \blacksquare

If u, v are vertices of a graph G , it is sometimes convenient to call the distance between u, v in G the $|G|$ -*distance* between u, v . We deduce:

4.2 *Let $c > 0$, and let G be a $(|G|^{1-c}/4, |G|/4)$ -coherent graph. Then there exists $u \in W$ and an integer $k < 1 + 1/c$, such that:*

- *at most $|G|/2$ vertices have G -distance less than k from u ; and*

- at least $|G|/4$ vertices have G -distance exactly k from u .

Proof. By 4.1, there exists $Y \subseteq V(G)$ with $|Y| \leq |G|^{1-c}/4$ such that $G[G \setminus Y]$ is τ -expanding, where $\tau = |G|^c$. Choose $u \in V(G) \setminus Y$, and for each integer $i \geq 0$ let M_i be the set of vertices of G that have G -distance at most i from u . Since $G \setminus Y$ is τ -expanding, it follows that for all $i \geq 0$, $|M_{i+1} \setminus Y| \geq \min(\tau|M_i \setminus Y|, |W \setminus Y|/2)$. For each $i \geq 1$, let $L_i = M_i \setminus M_{i-1}$.

(1) *There exists $k \leq 1 + 1/c$ such that $|L_k| \geq |W|/4$.*

Since $G \setminus Y$ is τ -expanding, it is connected, and so there exists i such that $V(G) \setminus Y \subseteq M_i$. Since $|V(G) \setminus Y| \geq 3|G|/4$, we may choose $j \geq 0$ minimum such that $|M_j| \geq |G|^{1-c}/4$. Hence for each $i \in \{1, \dots, j-1\}$, $|M_i| < |G|^{1-c}/4 < |V(G) \setminus Y|/2$, and so $|M_i \setminus Y| \geq \tau|M_{i-1} \setminus Y|$ since $G \setminus Y$ is τ -expanding. Since $|M_0 \setminus Y| = 1$, it follows that $|M_{j-1} \setminus Y| \geq \tau^{j-1}$. Hence

$$|G|^{(j-1)c} = \tau^{j-1} \leq |M_{j-1} \setminus Y| \leq |M_{j-1}| < |G|^{1-c}/4,$$

and so $(j-1)c < 1-c$, that is, $j < 1/c$.

Since G is $(|G|^{1-c}/4, |G|/4)$ -coherent, and M_j is anticomplete to $V(G) \setminus N(M_j)$, it follows that $|V(G) \setminus N(M_j)| < |G|/4$. But also $|M_{j-1}| < |G|^{1-c}/4$ (or $j=0$), and so $|L_j \cup L_{j+1}| \geq |G| - |G|/4 - |G|^{1-c}/4 \geq |G|/2$. Thus some $k \in \{j, j+1\}$ satisfies the claim. This proves (1).

Choose k as in (1), minimum. Thus $|L_{k-1}| < |G|/4$, and $|M_{k-2}| < |G|^{1-c}/4$ since G is $(|G|^{1-c}/4, |G|/4)$ -coherent. Thus $|M_{k-1}| \leq |G|/2$. This proves 4.2. ■

5 Covering sequences

Let us say a *covering* \mathcal{L} in G is a triple (a, H, B) where H, B are disjoint subsets of $V(G)$, $a \in H$, H covers B , and $G[H]$ is connected. We call a the *apex*, H the *heart*, and B the *base* of the covering. If for every vertex $v \in B$ there is a path of length at most r between a, v of length at most r , we say that (a, H, B) has *height* at most r , and the least such r is the height. For instance, if (L_0, \dots, L_k) is a levelling with $k > 0$, and $L_0 = \{a\}$, then $(a, L_0 \cup \dots \cup L_{k-1}, L_k)$ is a covering of height k .

A *covering sequence* in G is a sequence $(\mathcal{L}_1, \dots, \mathcal{L}_n)$ of coverings in G , with hearts H_1, \dots, H_n say, such that H_1, \dots, H_n are pairwise disjoint and pairwise anticomplete. We call n its *length*. We say such a sequence has *height* at most r if each term has height at most r . If $\mathcal{M} = (\mathcal{L}_1, \dots, \mathcal{L}_n)$ is a covering sequence, we define $V(\mathcal{M})$ to be the union of the sets $V(\mathcal{L}_i)$ for $1 \leq i \leq n$. A covering sequence $(\mathcal{L}_1, \dots, \mathcal{L}_n)$ is a *multicovering* if $\mathcal{L}_1, \dots, \mathcal{L}_n$ all have the same base, and then this common base is called the *base* of the multicovering.

The main result of this section says that a graph with the usual properties (suitably coherent, suitable sparse) contains a multicover of length any specified constant, with height at most about $1/c$ and with base of linear cardinality. We prove this in several steps. We begin with:

5.1 *Let $n \geq 0$ be an integer. Let $c > 0$ such that $1/c$ is an integer; let $\varepsilon > 0$ with $\varepsilon \leq 2^{-n-2}$; and let G be an ε -sparse $(\varepsilon|G|^{1-c}, \varepsilon|G|)$ -coherent graph. Then there is a covering sequence $(\mathcal{L}_1, \dots, \mathcal{L}_n)$ in G , where $\mathcal{L}_i = (a_i, H_i, B_i)$ for $1 \leq i \leq n$, such that:*

- for $1 \leq i < j \leq n$, H_i is anticomplete to B_j ;
- for $1 \leq i \leq n$, \mathcal{L}_i has height at most $1/c$; and
- for $1 \leq i \leq n$, $|B_i| \geq 2^{-i-1}|G|$.

Proof. We proceed by induction on n . If $n = 0$ the result is trivial, so we assume that $n \geq 1$ and the result holds for $n - 1$. By 4.2, there exists $u \in V(G)$ and an integer $k < 1 + 1/c$ (and hence $k \leq 1/c$, since $1/c$ is an integer), such that:

- at most $|G|/2$ vertices of G have distance less than k from u ; and
- at least $|G|/4$ vertices of G have distance exactly k from u .

For $0 \leq i \leq k$ let L_i be the set of all vertices of G with distance exactly i from u . Then (L_0, \dots, L_k) is a levelling, with height at most $1/c$; and $|L_k| \geq |G|/4$, so the theorem holds for $n = 1$. Choose $L'_{k-1} \subseteq L_{k-1}$ minimal such that at least $|G|/4$ vertices in L_k have a neighbour in L'_{k-1} , and let L'_k be the set of vertices in L_k that have a neighbour in L'_{k-1} . Thus $|L'_k| \leq (1/4 + \varepsilon)|G|$ since G is ε -sparse. Let \mathcal{L}_1 be the levelling $(L_0, \dots, L_{k-2}, L'_{k-1}, L'_k)$, and let H_1 be its heart. Thus $|V(\mathcal{L}_1)| \leq (3/4 + \varepsilon)|G|$. Let W be the set of vertices of G not in $V(\mathcal{L}_1)$; so $|W| \geq (1/4 - \varepsilon)|G|$. Since W is anticomplete to H_1 , and $1/4 - \varepsilon \geq \varepsilon$ and G is $(\varepsilon|G|^{1-c}, \varepsilon|G|)$ -coherent, it follows that $|H_1| \leq \varepsilon|G|^{1-c}$, and so $|W| \geq (3/4 - \varepsilon)|G| - \varepsilon|G|^{1-c} \geq |G|/2$. Hence $G[W]$ is (2ε) -sparse and $((2\varepsilon)|W|^{1-c}, (2\varepsilon)|W|)$ -coherent. From the inductive hypothesis applied to $G[W]$, there is a covering sequence $(\mathcal{L}_2, \dots, \mathcal{L}_n)$ in $G[W]$, where $\mathcal{L}_i = (a_i, H_i, B_i)$ for $2 \leq i \leq n$, such that:

- for $2 \leq i < j \leq n$, H_i is anticomplete to B_j ;
- for $2 \leq i \leq n$, \mathcal{L}_i has height at most $1/c$; and
- for $2 \leq i \leq n$, $|B_i| \geq 2^{-i}|W| \geq 2^{-i-1}|G|$.

But then $(\mathcal{L}_1, \dots, \mathcal{L}_n)$ satisfies the theorem. This proves 5.1. ▀

5.2 Let $n \geq 0$ be an integer, and let $m = 2^{2n}$. Let $c > 0$ such that $1/c$ is an integer, let $\varepsilon > 0$ with $\varepsilon \leq 2^{-m-2}$, and let G be an ε -sparse $(\varepsilon|G|^{1-c}, \varepsilon|G|)$ -coherent graph. Then there is a covering sequence $(\mathcal{L}_1, \dots, \mathcal{L}_n)$ in G , where $\mathcal{L}_i = (a_i, H_i, B_i)$ for $1 \leq i \leq n$, such that:

- for $1 \leq i \leq n$, \mathcal{L}_i has height at most $1/c$;
- for $1 \leq i \leq n$, $|B_i| \geq 2^{-m-1}|G|$;
- either B_1, \dots, B_n are pairwise disjoint and H_i is anticomplete to B_j for all distinct $i, j \in \{1, \dots, n\}$, or $B_1 = B_2 = \dots = B_n$.

Proof. Choose $\mathcal{L}_1, \dots, \mathcal{L}_m$ as in 5.1, where each \mathcal{L}_i has base of cardinality at least $2^{-i-1}|G|$, where $\mathcal{L}_i = (a_i, H_i, B_i)$ for $1 \leq i \leq m$. Each has height at most $1/c$. For $1 \leq i \leq m$, H_1, \dots, H_{i-1} are anticomplete to B_i , but H_{i+1}, \dots, H_m might have neighbours in B_i . Choose $B'_i \subseteq B_i$ of cardinality at least $|B_i|/2^{m-i} \geq |G|/2^{m+1}$, such that for $i + 1 \leq j \leq m$, either every vertex in B'_i has a

neighbour in H_j , or none do. Let \mathcal{L}'_i be the covering obtained from \mathcal{L}_i by replacing its base by B'_i . Then $(\mathcal{L}'_1, \dots, \mathcal{L}'_m)$ is a covering sequence, and for $1 \leq i < j \leq m$, H_i is anticomplete to B_j , and either H_j is anticomplete to B_i or H_j covers B_i . By Ramsey's theorem, since $m = 2^{2^n}$, there exists $I \subseteq \{1, \dots, m\}$ with $|I| = n$ such that either

- for all distinct $i, j \in m$, H_i is anticomplete to B_j (and hence $B_i \cap B_j = \emptyset$), or
- for all $i, j \in I$ with $i < j$, H_j is complete to B_i .

In both cases the theorem holds. This proves 5.2. ■

Now we prove the main result of this section. Its proof is closely related to the proof of the main theorem of [3].

5.3 *Let $c > 0$ such that $1/c$ is an integer, and let $n \geq 0$ be an integer. Then there exist $\varepsilon, d > 0$ with the following property. If G is an ε -sparse $(\varepsilon|G|^{1-c}, \varepsilon|G|)$ -coherent graph then G contains a multicovering of length n and height at most $1 + c^{-1}$, and with base of cardinality at least $d|G|$.*

Proof. Define $m = 3^n$, and let $N = 2^{2^m}$. Let $x = 2^{-N-1}$. Choose $\varepsilon > 0$ such that $\varepsilon, d \leq x3^{-n}$. It follows that $\varepsilon \leq 2^{-N-2}$.

From 5.2, we may assume that there is a covering sequence $(\mathcal{L}_1, \dots, \mathcal{L}_m)$ in G , such that:

- $V(\mathcal{L}_1), \dots, V(\mathcal{L}_m)$ are pairwise disjoint;
- for $1 \leq i \leq m$, \mathcal{L}_i has height at most $1/c$;
- for $1 \leq i \leq m$, the base of \mathcal{L}_i has cardinality at least $x|G|$; and
- for all distinct $i, j \in \{1, \dots, m\}$, every edge between $V(\mathcal{L}_i)$ and $V(\mathcal{L}_j)$ is between the base of $V(\mathcal{L}_i)$ and the base of $V(\mathcal{L}_j)$.

Let $t, d_1, \dots, d_t > 0$ be integers, where $d_1, \dots, d_t \leq n$. Let us say a *battery of type (d_1, \dots, d_t)* is a sequence of t multicoverings $(\mathcal{M}_1, \dots, \mathcal{M}_t)$ in G , such that:

- $V(\mathcal{M}_1), \dots, V(\mathcal{M}_t)$ are pairwise disjoint;
- for $1 \leq i \leq t$, \mathcal{M}_i has length d_i , and height at most $1 + 1/c$, and the first term of \mathcal{M}_i has height at most $1/c$;
- for $1 \leq i \leq t$, the base of \mathcal{M}_i has cardinality at least $x3^{1-d_i}|G|$;
- for all distinct $i, j \in \{1, \dots, m\}$, every edge between $V(\mathcal{M}_i)$ and $V(\mathcal{M}_j)$ is between the base of $V(\mathcal{M}_i)$ and the base of $V(\mathcal{M}_j)$.

Thus G contains a battery of type $(1, \dots, 1)$, of length m . Choose a battery \mathcal{B} of type (d_1, \dots, d_t) with t minimum such that $2^{d_1} + \dots + 2^{d_t} \geq m$. Let $\mathcal{B} = (\mathcal{M}_1, \dots, \mathcal{M}_t)$. For $1 \leq i \leq t$, let the base of \mathcal{M}_i be B_i . If some $d_i = n$, then the i th term of \mathcal{B} is a multicovering satisfying the theorem, so we assume not. If $t = 1$ then $2^{d_1} \geq m = 2^n$, and so $d_1 \geq n$, a contradiction; so $t \geq 2$, and $d_1, \dots, d_t < n$. Consequently each $|B_i| \geq x3^{1-(n-1)}|G| \geq \varepsilon|G|$. By reordering the terms of the battery, we may

assume that $d_t \leq d_1, \dots, d_{t-1}$. Since G is $(\varepsilon|G|^{1-c}, \varepsilon|G|)$ -coherent, and $|B_t| \geq \varepsilon|G|$, for $1 \leq i < t$ there are fewer than $\varepsilon|G|^{1-c} \leq 2|B_i|/3$ vertices in B_i that have no neighbour in B_t . Hence we may choose $X \subseteq B_t$ minimal such that for some $i \in \{1, \dots, t-1\}$, at least $|B_i|/3$ vertices in B_i have a neighbour in X . For $1 \leq i < t$, let Y_i be the set of vertices in B_i that have a neighbour in X , and $Z_i = B_i \setminus Y_i$. By reordering, we may assume that $|Y_1| \geq |B_1|/3$. From the minimality of X , $|Y_i| \leq |B_i|/3 + \varepsilon|G|$ for $2 \leq i \leq t-1$, and so $|Z_i| \geq 2|B_i|/3 - \varepsilon|G| \geq |B_i|/3$. Let $\mathcal{M}_1 = (\mathcal{L}_1, \dots, \mathcal{L}_{d_1})$, and let the first term of \mathcal{M}_t be $\mathcal{L} = (a, H, B_t)$. Let \mathcal{L}' be the covering $(a, H \cup X, Y_1)$, which therefore has height at most $1 + 1/c$. Let \mathcal{M}'_1 be obtained from \mathcal{M}_1 by replacing its base by Y_1 and adding a new final term \mathcal{L}'_1 ; so \mathcal{M}'_1 has length $d_1 + 1$. For $2 \leq i \leq t-1$, let \mathcal{M}'_i be obtained from \mathcal{M}_i by replacing its base by Z_i . Then $(\mathcal{M}'_1, \dots, \mathcal{M}'_{t-1})$ is a battery of type $(d_1 + 1, d_2, \dots, d_{t-1})$. Since $d_1 \geq d_t$, it follows that

$$2^{d_1+1} + \dots + 2^{d_{t-1}} \geq 2^{d_1} + \dots + 2^{d_t} \geq 2^m,$$

a contradiction to the choice of \mathcal{B} . This proves 5.3. ▀

6 Making spiders

Let $\mathcal{L}_1, \dots, \mathcal{L}_n$ be coverings in G , such that

- $\mathcal{L}_1, \dots, \mathcal{L}_n$ all have the same apex a ;
- for $1 \leq i \leq n$ let \mathcal{L}_i have heart H_i ; then for $1 \leq i < j \leq n$, $H_i \setminus \{a\}$ is disjoint from and anticomplete to $H_j \setminus \{a\}$.

We call $(a, \mathcal{L}_1, \dots, \mathcal{L}_n)$ a *spider* in G , and a is its *apex*. Its *height* is the maximum of the heights of $\mathcal{L}_1, \dots, \mathcal{L}_n$, and its *length* is n . It has *mass* b where b is the minimum cardinality of the bases of $\mathcal{L}_1, \dots, \mathcal{L}_n$. The union of the hearts of $\mathcal{L}_1, \dots, \mathcal{L}_n$ is called the *heart* of the spider. We call $\mathcal{L}_1, \dots, \mathcal{L}_n$ the *members* of the spider.

6.1 *Let $c > 0$ such that $1/c$ is an integer, and let $n \geq 1$ be an integer. Then there exists $\varepsilon, d > 0$ with the following property. If G is an ε -sparse $(\varepsilon|G|^{1-c}, \varepsilon|G|)$ -coherent graph with $|G| \geq 2$ then G contains a spider of length n and height at most $2 + 2c^{-1}$, and with mass at least $d|G|$.*

Proof. Choose ε, d' as in 5.3 (with d' replacing d). By reducing ε we may assume that $\varepsilon \leq d'/2$, and $\varepsilon < 1/3$. Let $d = d'/2$. We claim that ε, d satisfy the theorem. Let G be as in the theorem; then 5.3 implies that G contains a multicovering $(\mathcal{L}_1, \dots, \mathcal{L}_n)$ of length n and height at most $1 + c^{-1}$, and with base B of cardinality at least $d'|G|$.

Choose $a \in B$. Let $1 \leq i \leq n$, and let H_i be the heart of \mathcal{L}_i . Then $G[H_i \cup \{a\}]$ is connected, and every vertex of $H_i \cup \{a\}$ can be joined to a by a path of $G[H_i \cup \{a\}]$ with length at most $1 + 2/c$. Hence $(a, H_i \cup \{a\}, B \setminus \{a\})$ is a covering of height at most $2 + 2/c$, say \mathcal{L}'_i . Consequently $(\{a\}, \mathcal{L}'_1, \dots, \mathcal{L}'_n)$ is a spider of length n and height at most $2 + 2c^{-1}$, and mass $|B| - 1 \geq d'|G| - 1$. By 2.1, $|G| \geq 1/\varepsilon \geq 2/d'$, and so $d'|G| - 1 \geq d|G|$. This proves 6.1. ▀

A *troupe* of spiders is a set of spiders such that their hearts are pairwise anticomplete.

6.2 Let $c > 0$ such that $1/c$ is an integer, and let $m \geq 0, n \geq 1$ be integers. Then there exist $\varepsilon, d > 0$ with the following property. If G is an ε -sparse $(\varepsilon|G|^{1-c}, \varepsilon|G|)$ -coherent graph, then G contains a troupe of m spiders, each of length n and height at most $2 + 2/c$, and with mass at least $d|G|$.

Proof. Let ε', d' satisfy 6.1 (with ε, d replaced by ε', d' respectively). Define $\varepsilon = \varepsilon'(d'/2)^m$ and $d = d'(d'/2)^m$. We will show that ε, d satisfy the theorem.

We proceed by induction on m . For $m = 0$ the result is trivial, so we assume that $m \geq 1$ and the result holds for $m - 1$. By 6.1 G contains a spider of length n and height at most $2 + 2c^{-1}$, and with mass at least $d'|G|$; say $(a_1, \mathcal{L}_1, \dots, \mathcal{L}_n)$. For $1 \leq j \leq n$ let $\mathcal{L}_j = (a_1, H_j, B_j)$. We may assume that every vertex of H_j has $G[H_j]$ -distance from a_1 at most $1 + 2c^{-1}$ (because any other vertices can be deleted). Since H is connected, we can find a sequence of induced subgraphs of H , starting with the subgraph with one vertex a_1 , and adding vertices one by one, in such a way that each of these graphs is connected and every vertex is joined to a_1 by a path of length at most $1 + 2c^{-1}$ within the subgraph. Choose one of these subgraphs, say H'_j , the first such that at least $d|G|$ vertices in B_j have a neighbour in H'_j . Let B'_j be the set of vertices in B_j with a neighbour in H'_j . Thus $d|G| \leq |B'_j| \leq (d + \varepsilon)|G|$ since G is ε -sparse. Let \mathcal{L}'_j be the covering (a_1, H'_j, B'_j) .

The union of B'_1, \dots, B'_n has cardinality at most $n(d + \varepsilon)|G| \leq d'|G|/2$ and so there is a subset $X \subseteq B_1$ of cardinality at least $d'|G|/2$, anticomplete to H'_1, \dots, H'_n . Then $\mathcal{S}_1 = (a, \mathcal{L}'_1, \dots, \mathcal{L}'_n)$ is a spider of length n and height at most $2 + 2c^{-1}$, and with mass at least $d|G|$; and X is anticomplete to the heart of this spider.

Let $\varepsilon'' = 2\varepsilon/d'$, and $d'' = 2d/d'$. Since $|X| \geq d'|G|/2$, it follows that $G[X]$ is ε'' -sparse and $(\varepsilon''|G|^{1-c}, \varepsilon''|G|)$ -coherent. Since $\varepsilon'' \leq \varepsilon'(d'/2)^{m-1}$ and so on, we can apply the inductive hypothesis to $G[X]$, and deduce that there is a troupe of $m - 1$ spiders in $G[X]$, each of length n and height at most $2 + 2c^{-1}$, and with mass at least $d''|X| = (2d/d')|X| \geq d|G|$. But then adding \mathcal{S}_1 to this troupe gives a troupe of m spiders satisfying the theorem. This proves 6.2. \blacksquare

So, our graph contains a troupe of spiders, of arbitrarily large cardinality, and each with arbitrarily large length, all of height at most $2 + 2/c$, and with bases of linear cardinality. The next result converts the members of these spiders to levellings, but we need to be careful exactly what we mean. In a levelling, all edges from heart to base start from the penultimate term of the levelling. We need more than this: we need that for every two levellings involved as members of spiders in the troupe, every edge from the heart of one to the base of the other leaves from the penultimate level of the first, and this is more tricky to arrange.

Let us first state the definition formally. Let $n \geq 1$ and let $\mathcal{L}_1, \dots, \mathcal{L}_n$ be levellings in a graph G , all with the same apex a , such that for $1 \leq i < j \leq n$, every edge of G between $V(\mathcal{L}_i) \setminus \{a\}$ and $V(\mathcal{L}_j) \setminus \{a\}$ is between the base of \mathcal{L}_i and the base of \mathcal{L}_j . We call $(a, \mathcal{L}_1, \dots, \mathcal{L}_n)$ a *lobster* in G , and a is its *apex*. Its *height* is the maximum height of $\mathcal{L}_1, \dots, \mathcal{L}_n$, and its *length* is n . It has *mass* b where b is the minimum cardinality of the bases of $\mathcal{L}_1, \dots, \mathcal{L}_n$. Its *heart* is the union of the hearts of $\mathcal{L}_1, \dots, \mathcal{L}_n$. We call $\mathcal{L}_1, \dots, \mathcal{L}_n$ the *members* of the lobster.

A *troupe* of lobsters is a set $\{\mathcal{T}_1, \dots, \mathcal{T}_m\}$ of lobsters, such that for all $i, j \in \{1, \dots, m\}$:

- for $1 \leq i < j \leq m$, the heart of \mathcal{T}_i is disjoint from and anticomplete to the heart of \mathcal{T}_j ;
- let \mathcal{L}, \mathcal{M} each be a member of one of $\mathcal{T}_1, \dots, \mathcal{T}_m$, with $\mathcal{L} \neq \mathcal{M}$, and let $\mathcal{L} = \{L_0, \dots, L_k\}$; then there is no edge between $L_0 \cup \dots \cup L_{k-2}$ and the base of \mathcal{M} .

6.3 Let $c > 0$ such that $1/c$ is an integer, and let $m \geq 0, n \geq 1$ be integers. Then there exists $\varepsilon, d > 0$ with the following property. If G is an ε -sparse $(\varepsilon|G|^{1-c}, \varepsilon|G|)$ -coherent graph, then G contains a troupe of m lobsters, each of length n and height at most $2 + 2/c$, and with mass at least $d|G|$.

Proof. Let ε, d' satisfy 6.2 with d replaced by d' . Define $w(h) = d'(1 + 2/c)^{-h}$ for $0 \leq h \leq n$, and define $d = w(n)$. We claim that ε, d satisfy the theorem.

Let G be an ε -sparse $(\varepsilon|G|^{1-c}, \varepsilon|G|)$ -coherent graph. By 6.2 there is a troupe of spiders $\{\mathcal{S}_1, \dots, \mathcal{S}_m\}$ in G , each of length n and height at most $2 + 2/c$, and with mass at least $d|G|$. Let the members of these spiders (in some order) be $\mathcal{L}_1, \dots, \mathcal{L}_n$, and for $1 \leq i \leq n$ let $\mathcal{L}_i = \{a_i, H_i, B_i\}$. We shall convert them one by one to levellings, at each step shrinking all the bases. Let $X^0 = B_1 \cup \dots \cup B_n$, and for $1 \leq i \leq n$ let X_i^0 be the set of all vertices in X^0 with a neighbour in H_i (thus $B_i \subseteq X_i^0$). Inductively, let $1 \leq h \leq n$, and suppose that X^{h-1} and $\mathcal{L}'_1, \dots, \mathcal{L}'_{h-1}$ are defined, and for $1 \leq i \leq n$ X_i^{h-1} is defined,

- for $1 \leq i \leq h-1$, \mathcal{L}'_i is a levelling; its heart is a subset of H_i , and a_i is its apex; its height is at most $2 + 2/c$;
- for $1 \leq i \leq h-1$, X_i^{h-1} is the set of all vertices in X^{h-1} with a neighbour in the heart of \mathcal{L}'_i , and for $h \leq i \leq m$, X_i^{h-1} is the set of all vertices in X^{h-1} with a neighbour in the heart of \mathcal{L}_i ;
- for $1 \leq i \leq h-1$, every edge between the heart of \mathcal{L}'_i and X^{h-1} has an end in the penultimate term of \mathcal{L}'_i ; and
- for $1 \leq i \leq n$, $|X_i^{h-1}| \geq w(h-1)|G|$.

For $0 \leq j \leq 1 + 2/c$, let L_j be the set of vertices in H_h with $G[H_h]$ -distance to a_h exactly j . Thus every vertex $v \in X_h^{h-1}$ has a neighbour in one of L_0, \dots, L_j where $j \leq 1 + 2/c$, and the smallest such j is called the *type* of v . There are only $1 + 2/c$ possible types, and so there exists $k \in \{1 + 2/c\}$ such that at least $|X_h^{h-1}|/(1 + 2/c)$ vertices in X_h^{h-1} have type k . Consequently, since

$$|X_h^{h-1}|/(1 + 2/c) \geq w(h-1)|G|/(1 + 2/c) = w(h)|G|,$$

there exists $k \in \{1 + 2/c\}$ minimum such that at least $w(h)|G|$ vertices in X_h^{h-1} have type k . Let X_h^h be the set of all vertices in X_h^{h-1} that have type k . Let $\mathcal{L}'_h = (L_0, \dots, L_k, X_h^h)$. Thus \mathcal{L}'_h is a levelling with height $k + 1 \leq 2 + 2/c$.

Let Z^h be the set of vertices in X_h^{h-1} with type less than k . Thus $|Z^h| \leq (2/c)w(h)|G|$. For $1 \leq i \leq n$ with $i \neq h$, define $X_i^h = X_i^{h-1} \setminus Z^h$. Thus $|X_i^h| \geq |X_i^{h-1}| - |Z^h|$, and so $|X_i^h| \geq w(h-1)|G| - (2/c)w(h)|G| \geq w(h)|G|$. Let X^h be the union of the sets X_i^h ($1 \leq i \leq n$). This completes the inductive definition.

For $1 \leq i \leq m$, let \mathcal{T}_i be the lobster obtained from \mathcal{S}_i by replacing each member \mathcal{L}_j by \mathcal{L}'_j . This makes a troupe of lobsters satisfying the theorem, and so proves 6.3. ■

7 Part assembly

Now we put these several pieces together to prove 1.8, which we restate:

7.1 Let $c > 0$ with $1/c$ an integer, and let H_1, H_2 be graphs with branch-length at least $4c^{-1} + 5$. Then there exists $\varepsilon > 0$ such that if G is a graph with $|G| > 1$ that is H_1 -free and $\overline{H_2}$ -free, then there is a pure pair A, B in G with $|A| \geq \varepsilon|G|$ and $|B| \geq \varepsilon|G|^{1-c}$.

We saw in section 2 that to prove 7.1, it suffices to show:

7.2 Let $c > 0$ with $1/c$ an integer, and let H be a graph with branch-length at least $4c^{-1} + 5$. Then there exists $\varepsilon > 0$ such that if G is an H -free graph with $|G| > 1$ and with maximum degree at most $\varepsilon|G|$, then there is an anticomplete pair A, B in G with $|A| \geq \varepsilon|G|$ and $|B| \geq \varepsilon|G|^{1-c}$.

Proof. By adding more vertices to H , we may assume that if X denotes the set of vertices of H that have degree different from two, then every cycle of H contains at least one vertex in X , and every path in H with both ends in X has length at least $4c^{-1} + 5$, and every cycle of H has length at least $4c^{-1} + 5$. Let $X = \{x_1, \dots, x_m\}$. Consequently H can be obtained from the set X of m isolated vertices by adding

- paths with ends in X and each of length at least $4/c + 5$, and
- cycles with exactly one vertices in X , of length at least $4/c + 5$

where every vertex of $V(H) \setminus X$ belongs to exactly one of these paths and cycles, and has degree exactly two in H . Let the paths be R_i ($i \in I_1$), and let the cycles be R_i ($i \in I_2$), where $I_1 \cup I_2 = \emptyset$. For $i \in I_1$, let R_i have ends (u_i, v_i) (ordered arbitrarily) and have length ℓ_i , and for $i \in I_2$, let $u_i = v_i$ be the unique vertex of R_i in X , and let R_i have length ℓ_i . Thus H is determined up to isomorphism by a knowledge of X , the pairs (u_i, v_i) ($i \in I_1 \cup I_2$), and the numbers ℓ_i ($i \in I_1 \cup I_2$). Let $I = I_1 \cup I_2$, and for each $i \in I$ let $\alpha_i \in \{1, \dots, m\}$ such that $x_{\alpha_i} = u_i$ and let $\beta_i \in \{1, \dots, m\}$ such that $x_{\beta_i} = v_i$.

Let n be the maximum degree of H . Choose ε, d as in 6.3. By reducing ε , we may assume that $\varepsilon \leq d/|H|$, and that ε satisfies 3.4 for all choices of integers $s, t > 0$ satisfying $s, t \leq 2 + 2c^{-1}$. We claim that ε satisfies the theorem. Let G be an ε -sparse $(\varepsilon|G|^{1-c}, \varepsilon|G|)$ -coherent graph. We must show that G contains H . By 6.3, G contains a troupe $\{\mathcal{S}_1, \dots, \mathcal{S}_m\}$ of m lobsters, each of length n and height at most $2 + 2c^{-1}$, and with mass at least $d|G|$. Let $I = \{1, \dots, p\}$. For $1 \leq h \leq p$, choose a member \mathcal{L}_{2i-1} of \mathcal{S}_{α_i} and a member \mathcal{L}_{2i} of \mathcal{S}_{β_i} , in such a way that the levellings $\mathcal{L}_i, \mathcal{M}_i$ ($i \in I$) are all different. (This is possible from the definition of n).

We will prove that for all $h \in I$, there is a path P_h (or cycle, if the two apexes are equal) between the apex of \mathcal{L}_{2h-1} and the apex of \mathcal{L}_{2h} of length ℓ_h , such that the union of P_1, \dots, P_p makes an induced subgraph of G isomorphic to H . We will choose these paths and cycles in order. Also for $1 \leq h \leq p$ we need to choose a subset of the base of each \mathcal{L}_k for $2h < k \leq 2p$, and a subset of the penultimate term of \mathcal{L}_k ; these are denoted by X_k^h and Y_k^h . We denote by P_h^* the set of vertices of P_h different from its ends (if it is a path) or different from the apex of A_h (if it is a cycle). In either case $|P_h^*| = \ell_h - 1$.

For $0 \leq h \leq p$ let $w_h = (4p)^{p-h}d$. For $1 \leq i \leq 2p$, let X_i^0 be the base of \mathcal{L}_i , and let Y_i^0 be the penultimate term of \mathcal{L}_i . Let a_i be its apex. Let B be the union of the sets X_i^0 for $1 \leq i \leq 2p$. Now inductively, suppose we have chosen the first $h - 1$ paths or cycles, say P_1, \dots, P_{h-1} , where $1 \leq h \leq p$, satisfying:

- for $1 \leq g \leq h - 1$, if $a_{2g-1} \neq a_{2g}$, then P_g is an induced path joining these apexes, of length ℓ_g ; and if the apexes are equal then P_g is a cycle of length ℓ_g containing this apex;

- for $1 \leq g \leq h-1$, P_g^* is anticomplete to the hearts of $L_{2h+1}, \dots, \in \surd$.

Suppose moreover that for $2h+1 \leq i \leq 2p$ we have chosen $X_i^{h-1} \subseteq X_i^0$ and $Y_i^{h-1} \subseteq Y_i^0$, such that for all $i \in \{2h+1, \dots, 2p\}$:

- X_i^{h-1} is the set of all vertices in B with a neighbour in Y_i^{h-1} ;
- $X_i^{h-1} \cup Y_i^{h-1}$ is anticomplete to P_1^*, \dots, P_{h-1}^* ;
- $|X_{i,j}^{h-1}| \geq w_{h-1}|G|$.

We choose P_h as follows. For $2h+1 \leq i \leq 2p$, choose $Y_i \subseteq Y_i^{h-1}$ minimal such that at least $(w_h + \varepsilon(|H| - 1))|G|$ vertices in B (necessarily all in X_i^{h-1}) have a neighbour in Y_i , and let X_i be the set of vertices in B_i with a neighbour in Y_i . From the minimality of Y_i ,

$$(w_h + \varepsilon(|H| - 1))|G| \leq |X_i| \leq (w_h + \varepsilon|H|)|G|.$$

Let $Z = X_{2h+1} \cup \dots \cup X_{2p}$. Thus $|Z| \leq (2p-2)(w_h + \varepsilon\ell_h)|G|$. For $i = 2h-1, 2h$ let $X_i = X_i^{h-1} \setminus Z$. Thus

$$|X_i| \geq |X_i^{h-1}| - |Z| \geq (w_{h-1} - 2p(w_h + \varepsilon\ell_h))|G|$$

for $i = 2h-1, 2h$. Now $\ell_h \leq |H|$ and $\varepsilon|H| \leq d \leq w_h$, so $w_h + \varepsilon\ell_h \leq 2w_h$; and hence

$$|X_i| \geq (w_{h-1} - 2(2p-2)w_h)|G| \geq w_h|G| \geq d|G|$$

since $w_{h-1} = 4pw_h$. For $i = 2h-1, 2h$ let \mathcal{L}'_i be the levelling obtained from \mathcal{L}_i by replacing its base by X_i .

Now $\mathcal{L}'_{2h-1}, \mathcal{L}'_{2h}$ both have height at most $2 + 2/c$, and $\ell_h \geq 5 + 4/c$. By 3.4 applied to the levellings $\mathcal{L}'_{2h-1}, \mathcal{L}'_{2h}$, there is an induced path P_h between a_{2h-1}, a_{2h} (or a cycle, if $a_{2h-1} = a_{2h}$) of length ℓ_h , with vertex set included in $V(\mathcal{L}'_{2h-1}) \cup V(\mathcal{L}'_{2h})$. Consequently P_h^* is anticomplete to Y_i for $2h+1 \leq i \leq 2p$, and to P_1^*, \dots, P_{h-1}^* . It might have neighbours in X_i for $2h+1 \leq i \leq 2p$, but since $|P_h^*| \geq |H|$, there are at most $\varepsilon|H|$ such vertices. For $2h+1 \leq i \leq 2p$, let X_i^h be the set of vertices in X_i with no neighbour in P_h^* . Thus $|X_i^h| \geq |X_i| - \varepsilon|H| \geq w(h)|G|$. This completes the inductive definition.

But then the union of P_1, \dots, P_p forms an induced subgraph isomorphic to H . This proves 7.2, and hence completes the proof of 1.8. ■

References

- [1] J. Fox, “A bipartite analogue of Dilworth’s theorem”, *Order* **23** (2006), 197–209.
- [2] J. Fox, J. Pach and C. Tóth, “Turán-type results for partial orders and intersection graphs of convex sets”, *Israel J. Math.* **178** (2010), 29–50.
- [3] A. Liebenau, M. Pilipczuk, P. Seymour and S. Spirkl, “Caterpillars in Erdős-Hajnal”, *J. Combinatorial Theory, Ser. B*, **136** (2019), 33–43, [arXiv:1810.00811](https://arxiv.org/abs/1810.00811).

- [4] V. Rödl, “On universality of graphs with uniformly distributed edges”, *Discrete Math.* **59** (1986), 125–134.
- [5] A. Scott, P. Seymour and S. Spirkl, “Pure pairs. VIII. Excluding a sparse graph”, in preparation.