

Subdivisions and near-linear stable sets

Tung Nguyen¹
Princeton University,
Princeton, NJ 08544, USA

Alex Scott²
Oxford University,
Oxford, UK

Paul Seymour³
Princeton University,
Princeton, NJ 08544, USA

May 10, 2024; revised March 17, 2025

¹Supported by AFOSR grant FA9550-22-1-0234, and by NSF grant DMS-2154169, and a Porter Ogden Jacobus Fellowship.

²Supported by EPSRC grant EP/X013642/1

³Supported by AFOSR grant FA9550-22-1-0234, and by NSF grant DMS-2154169.

Abstract

We prove that for every complete graph K_t , all graphs G with no induced subgraph isomorphic to a subdivision of K_t have a stable subset of size at least $|G|/\text{polylog } |G|$. This is close to best possible, because for $t \geq 7$, not all such graphs G have a stable set of linear size, even if G is triangle-free.

1 Introduction

Graphs in this paper are finite, and have no loops or parallel edges. For a graph G , $|G|$ denotes the number of vertices of G , $\chi(G)$ is its chromatic number, and $\omega(G)$ and $\alpha(G)$ denote the sizes of its largest clique and stable set respectively. If G, H are graphs, G is H -free if no induced subgraph of G is isomorphic to H , and if \mathcal{H} is a set of graphs, G is \mathcal{H} -free if G is H -free for each $H \in \mathcal{H}$. We say G is H -subdivision-free if no induced subgraph of G is isomorphic to a subdivision of H . All logarithms are to base two.

We will prove:

1.1 *Let $t \geq 3$ be an integer, and let $c_t = (2t)^{-4(t-2)}$. Then every K_t -subdivision-free graph G with $|G| \geq 2$ satisfies*

$$\alpha(G) \geq \frac{c_t |G|}{(\log |G|)^{3t-5}}.$$

Let us put this in some context. First, a *string graph* is the intersection graph of curves in the plane. String graphs with clique number at most k are K_t -subdivision-free, where t depends only on k , and so our result extends a result of Fox and Pach [7] (and see also [8]):

1.2 *For every integer $k \geq 1$, there exists $c > 0$ such that every string graph G with clique number at most k has a stable set of size at least $|G|/(\log |G|)^c$.*

Our result is numerically weaker: c in the Fox-Pach theorem is $O(\log k)$, while for us it is $O(k)$ (applying 3.1). Nevertheless, it is pleasing to extend a geometric result to a much broader class of graphs.

Second, the Gyárfás-Sumner conjecture [10, 20] says:

1.3 Conjecture: *For every forest H and every integer $k \geq 1$, there exists $c \geq 1$ such that $\chi(G) \leq c$ for every $\{H, K_{k+1}\}$ -free graph G .*

In particular, if the Gyárfás-Sumner conjecture is true, then every $\{H, K_{k+1}\}$ -free graph G has a stable set of size at least $|G|/c$. This is still open, but in [16] we proved that it is “nearly” true, in the following sense:

1.4 *For every forest H and integer k , every $\{H, K_{k+1}\}$ -free graph G satisfies $\alpha(G) \geq |G|^{1-o(1)}$, and hence has chromatic number at most $|G|^{o(1)}$.*

One might regard this as evidence in support of the Gyárfás-Sumner conjecture, but one should not place much faith in such evidence, because the result of this paper provides just as much evidence to support a conjecture that turned out to be false. Scott [18] suggested an analogue of 1.3, that we could replace excluding a forest with excluding all subdivisions of any given graph:

1.5 False conjecture: *For every graph H , and every integer $k \geq 1$, there exists c such that every K_{k+1} -free, H -subdivision-free graph G satisfies $\chi(G) \leq c$.*

Scott proved that this is true if H is a forest, but it was shown to be false in general by Pawlik, Kozik, Krawczyk, Lasoń, Micek, Trotter and Walczak [17]. Results of Chalopin, Esperet, Li and Ossona de Mendez [2] and Walczak [21] together gave a strengthening:

1.6 *Let H be obtained from K_4 by subdividing once every edge in a cycle of length four. There are infinitely many K_3 -free and H -subdivision-free graphs G such that $\alpha(G) < |G|/\log \log |G|$.*

Every subdivision of K_7 contains a triangle or a subdivision of the graph H of 1.6, so the graphs G of 1.6 are K_7 -subdivision-free, as mentioned in the abstract.

However, we can prove that all H -subdivision-free graphs G with bounded clique number have a stable set of size $|G|/\text{polylog } |G|$, and so 1.5 is “nearly” true in the same sense as 1.3, despite 1.6. We will show:

1.7 *Let G be a graph with $|G| \geq 2$, let $k \geq 1$ be an integer with $\omega(G) \leq k$, and let H be a graph such that no induced subgraph of G is a subdivision of H . Then*

$$\alpha(G) \geq \frac{|G|}{(2|H|)^{4(k-1)}(\log |G|)^{3k-2}}.$$

This immediately implies a polylogarithmic bound $O((\log |G|)^{3k-1})$ on the chromatic number of such graphs G .

We remark that if we work with induced minors rather than induced subdivisions, there is a related result, proved by T. Korhonen and D. Lokshtanov [13]:

1.8 *If G is a graph with m edges, and does not contain a graph H as an induced minor, then G has a separator of cardinality $O(\sqrt{m})$.*

(We omit the definitions of “induced minor” and “separator”, since we will not need them any more.) By an argument of Fox and Pach, this implies that if in addition G satisfies $\omega(G) \leq k$, then $\alpha(G) \geq |G|(\log |G|)^{-O(\log k)}$.

2 A sketch of the proof

To prove 1.1, for inductive purposes we will prove a stronger result, proving a bound on $\alpha(G)$ that depends not only on t but also on $\omega(G)$ and on the maximum degree $\Delta(G)$ of G . In addition, we find that the same bound holds even if we only exclude subdivisions of K_t in which each edge is subdivided at least twice and at most $(\log |G|)^2 - 1$ times. Let us say a subdivision of H has *lengths between p and q* if each edge of H is replaced by a path of length between p and q . So our main result is:

2.1 *Let G be a graph. Let $d \geq 2$ be an integer with $\Delta(G) \leq d$; let $k \geq 1$ be an integer with $\omega(G) \leq k$; and let $t \geq 3$ be an integer such that no induced subgraph of G is a subdivision of K_t with lengths between 3 and $(\log |G|)^2$. Then*

$$\alpha(G) \geq \frac{|G|}{(2t)^{4(k-1)}(\log |G|)^{3(k-1)} \log d}.$$

We give the proof in the next section, but let us sketch it first. There is an initial step, that we can assume $\Delta(G) \leq |G|/(\text{polylog } |G|)$, because if there is a vertex of sufficiently large degree we just apply induction on the clique number to its set of neighbours and win.

The remainder of the proof breaks into two parts. In the first part we find t disjoint “stars”: t suitable vertices a_1, \dots, a_t , that we intend to be the high-degree vertices of a subdivision of K_t that we will try to construct. They need to be nonadjacent, and for each to have a large stable set B_i of neighbours nonadjacent to the other a_j ’s; and we need these sets B_i to be sparse to one another. We try to construct a_1, \dots, a_t greedily. Say we have selected a_1, \dots, a_i with corresponding sets B_1, \dots, B_i . Let G' be the set of vertices that nonadjacent to a_1, \dots, a_i , and sparse to each of B_1, \dots, B_i (the sparsity we need here depends just on t). Then G' still contains nearly all the vertices of G , and if it has maximum degree at most $\Delta(G)/2$, we win by inductively applying the theorem to G' . This is why we need the $\log d$ term in 2.1, and this is the only place we use that.

If there is a vertex a_{i+1} in G' with degree at least $\Delta(G)/2$, we apply the theorem inductively to its set X of neighbours, and find a large stable subset B_{i+1} , and they form the next star. It is critical here that the clique number of $G[X]$ is less than that of G , and since the bound we are proving depends on clique number, the large stable subset we find is bigger (by a polylog $|G|$ factor) than we could guarantee in a general set of the same size as X .

When we have found suitable a_1, \dots, a_t and B_1, \dots, B_t , we go to the second part of the proof. Each pair of B_i ’s needs to be joined by a path, and we need to choose these paths so that their union is an induced subgraph. So, suppose we have chosen some of the paths, and let us try to choose the next one. We need to put aside all vertices with neighbours in the paths that are already chosen, and route the next path through the remaining vertices. For this we must arrange that the number of vertices put aside is not too big. Each vertex only disqualifies at most $\Delta(G)$ other vertices, but we must arrange that the total number of vertices in the paths already selected is under control; and to do this, we insist that each path must have length at most some fixed power of $\log |G|$, and we find $(\log |G|)^2$ works.

There is another disqualification issue as well: some vertices in the B_i ’s cannot be used because they have neighbours in already-selected paths. So to keep this number small, we will take care to route all paths through the set of vertices that are reasonably sparse to each B_i . Since we will have to disqualify the neighbours in B_i of about $t^2(\log |G|)^2$ vertices, we need “reasonably sparse” to mean having $O(|B_i|/(\log |G|)^2)$ neighbours in each B_i . The number of vertices disqualified for this reason turns out to be $O((\log |G|)^2 \Delta(G))$, about the same as the number disqualified for the first reason. So there is a set S say, of size about $(\log |G|)^2 \Delta(G)$, of disqualified vertices, and we have to route the next path between B_1 and B_2 (say) without using any vertex in S .

To do this, we look at set expansion. Start with $N_1^0 = B_1$, and define N_1^i to be the set of vertices that belong to or have a neighbour in N_1^{i-1} . Suppose we could prove that $|N_1^i| \geq (1+2/\log |G|)|N_1^{i-1}|$ for each i ; then when i is about $(\log |G|)^2/2$, N_1^i contains more than half the vertices of the graph. Do the same for B_2 , and so N_1^i and N_2^i intersect, and we can pick out a short path between B_1, B_2 in their union. This is the plan, but to make it work there are several things to worry about.

First, how can we prove that subsets have large expansion in this sense? If there is a set X , of size less than $|G|/2$ and such that the set N of vertices with neighbours in X (including X itself) is not very big, then we can apply the inductive hypothesis to X and to $G \setminus N$, find large stable sets in each, and take their union to find a really large stable set in G and win. So this gives a mechanism to show that N is large, for any set X .

Second, we have to avoid the disqualified vertices S ; so we should redefine N_1^i to be the set of vertices, not in S , that have a neighbour in N_1^{i-1} , and then that issue is resolved. But to prove that $|N_1^i| \geq (1 + 2/\log |G|)|N_1^{i-1}|$ with the new meaning of N_1^i , we would need to prove that the

set of *all* neighbours of N_1^{i-1} in G is somewhat larger, so that even when we remove those in S , what remains is still large. So we need the set of neighbours of N_1^{i-1} to be something like at least $|S| + (1 + 2/\log |G|)|N_1^{i-1}|$. This is not a problem for the larger values of i , because then $|S|$ is fixed and all the other sets are getting big. It would be a problem when $i = 1$, if all we knew about the B_i 's was that their size was something like $\Delta(G)$, because S is much larger than this and the expansion argument would not work. But when $i = 1$ we have an advantage, because B_i is already stable, and we found it by using the induction on clique number, as explained earlier. That gain is enough to make the expansion argument work again; indeed, the expansion when $i = 1$ is big enough to ensure that all the subsequent sets N_1^j are comfortably large and consequently they expand nicely.

3 The main proof

In this section we prove our main result, which we restate:

3.1 *Let G be a graph. Let $d \geq 2$ be an integer with $\Delta(G) \leq d$; let $k \geq 1$ be an integer with $\omega(G) \leq k$; and let $t \geq 3$ be an integer such that no induced subgraph of G is a subdivision of K_t with lengths between 3 and $(\log |G|)^2$. Then*

$$\alpha(G) \geq \frac{|G|}{(2t)^{4(k-1)}(\log |G|)^{3(k-1)} \log d}.$$

Proof. We assume inductively that the result holds for all proper induced subgraphs of G (for all choices of $k \geq 1$ and $t \geq 3$). If $k = 1$, the result is trivial, so we may assume that $k \geq 2$. If $\Delta(G) \leq 1$, then $\alpha(G) \geq |G|/2$ and the result holds since $(2t)^{4(k-1)} \geq 6^4 \geq 2$; so we may assume that $\Delta(G) \geq 2$. Hence, by reducing d if possible, we may assume that $\Delta(G) = d$. To save writing, let us define $T = (2t)^4$, $L = \log |G|$, and $D = \log d$. Then we must show that $\alpha(G) \geq \frac{|G|}{T^{k-1}L^{3(k-1)}D}$. We suppose this is false. We claim:

$$(1) \quad dD^3 \leq |G|/T \text{ and } d+1 \geq T^{k-1}L^{3(k-1)}D, \text{ and consequently } d \geq 10^9 \geq 2^{29}.$$

Let v be a vertex with degree $\Delta(G) = d$, and let X be the set of all neighbours of v ; so $|X| = d$, and $\omega(G[X]) \leq k-1$. From the inductive hypothesis,

$$\alpha(G[X]) \geq \frac{d}{T^{k-2}D^{3(k-2)}D}$$

and therefore this is smaller than

$$\frac{|G|}{T^{k-1}L^{3(k-1)}D},$$

because otherwise the result would hold. This proves the first inequality of (1). For the second, since G has maximum degree d , and hence has a stable set of size at least $|G|/(d+1)$, it follows that

$$\frac{|G|}{d+1} < \frac{|G|}{T^{k-1}L^{3(k-1)}D},$$

and so $d+1 \geq T^{k-1}L^{3(k-1)}D$. This proves the second inequality. Since $|G| \geq d$, and $t \geq 3$ and $k \geq 2$, it follows that $T \geq 6^4$; since $L^{3(k-1)}D \geq D^4$, it follows that $d+1 \geq 6^4(\log d)^4$ and therefore

$d > 10^9$. This proves (1).

(2) If $X \subseteq V(G)$ and $|X| \leq |G|/D$, then $\Delta(G \setminus X) \geq d/2$.

Let $G' = G \setminus X$, and let $d' = \lfloor d/2 \rfloor$, and suppose that $\Delta(G') \leq d/2$. Thus $\Delta(G') \leq d'$, and $d' \geq 2$, so from the inductive hypothesis,

$$T^{k-1}\alpha(G') \geq \frac{|G'|}{(\log |G'|)^{3(k-1)} \log d'} \geq \frac{(1-1/D)|G|}{L^{3(k-1)}(D-1)} = \frac{|G|}{L^{3(k-1)}D},$$

and so the result holds, a contradiction. This proves (2).

If $A, B \subseteq V(G)$ are disjoint, we say A is x -sparse to B if each vertex in A has at most $x|B|$ neighbours in B , and the smallest such x is the *sparcity* of A to B . Let us say a *star system* in G is a pair of sequences (a_1, \dots, a_n) and (B_1, \dots, B_n) of the same length, with the following properties:

- $a_1, \dots, a_n \in V(G)$ are distinct and pairwise nonadjacent;
- B_1, \dots, B_n are pairwise disjoint stable subsets of $V(G) \setminus \{a_1, \dots, a_n\}$; and
- for $1 \leq i \leq n$, a_i is adjacent to each vertex in B_i and has no other neighbour in $B_1 \cup \dots \cup B_n$.

We call n the *length* of the star system, and its *size* is $\min(|B_i| : 1 \leq i \leq n)$ (or $|G|$, if $n = 0$). Its *semi-sparcity* is the maximum over all i, j with $1 \leq i < j \leq n$ of the sparsity of B_j to B_i (or 0, if $n \leq 1$). Its *sparcity* is the maximum over all distinct $i, j \in \{1, \dots, n\}$ of the sparsity of B_i to B_j (or 0, if $n \leq 1$).

(3) Let p be a nonnegative integer, and let $0 < q \leq 1$, such that $p/q \leq 400T$. Then G contains a star system of length p , size at least $\frac{d}{2T^{k-2}D^{3k-5}}$, and semi-sparcity at most q .

We proceed by induction on p , and so we may assume that $p \geq 1$ and G contains a star system $(a_1, \dots, a_{p-1}), (B_1, \dots, B_{p-1})$ of length $p-1$, size at least $\frac{d}{2T^{k-2}D^{3k-5}}$, and semi-sparcity at most q . Let X_1 be the set of vertices of G that are adjacent to one of a_1, \dots, a_{p-1} . Let X_2 be the set of vertices v of $G \setminus X_1$ such that for some $i \in \{1, \dots, p-1\}$, v has at least $q|B_i|$ neighbours in B_i .

Thus $|X_1| \leq (p-1)d \leq pd/q$, and $|X_2| \leq (p-1)d/q \leq pd/q$ (because there are at most $|B_i|d$ edges incident with B_i); and so $X = X_1 \cup X_2$ has cardinality at most $(2p/q)d$. Since $D^2 \geq 800$ by the last statement of (1), it follows (from the first statement of (1)) that

$$(2p/q)dD \leq (p/q)dD^3/400 \leq (p/q)|G|/(400T) \leq |G|.$$

Consequently $|X| \leq |G|/D$, and so by (2), $\Delta(G \setminus X) \geq d/2$. Hence there exists $a_p \in V(G) \setminus X$ with degree at least $d/2$ in $G \setminus X$. Let C be the set of its neighbours in $G \setminus X$; so $|C| \geq d/2$. Now $G[C]$ has clique number at most $k-1$, so from the inductive hypothesis, there exists a stable subset $B_p \subseteq C$ with

$$|B_p| \geq \frac{|C|}{T^{k-2}(\log |C|)^{3(k-2)}D} \geq \frac{d}{2T^{k-2}D^{3k-5}}.$$

But then $(a_1, \dots, a_{p-1}, a_p)$ and $(B_1, \dots, B_{p-1}, B_p)$ form the desired star system. This proves (3).

(4) Let p be a nonnegative integer, and let $0 < q \leq 1$, such that $p^2/q \leq 200T$. Then G contains a star system of length p , size at least $\frac{d}{4T^{k-2}D^{3k-5}}$, and sparsity at most q .

Let $q' = q/(2p)$; then by (3), G contains a star system $(a_1, \dots, a_p), (B_1, \dots, B_p)$ of length p , size at least $\frac{d}{2T^{k-2}D^{3k-5}}$, and semi-sparsity at most q' . Inductively, for $i = p, p-1, \dots, 1$ in turn, we define $C_i \subseteq B_i$ as follows. For $i+1 \leq j \leq p$, since C_j is q' -sparse to B_i , at most $|B_i|/(2p)$ vertices in B_i have more than $2pq'|C_j|$ neighbours in C_j . Hence there exists $C_i \subseteq B_i$ with $|C_i| \geq |B_i|/2$ that is $2pq'$ -sparse and hence q -sparse to each of C_{i+1}, \dots, C_p . Moreover, for $i < j \leq p$, C_j is $2q'$ -sparse and hence q -sparse to C_i , since $|C_i| \geq |B_i|/2$. This completes the inductive definition of C_p, \dots, C_1 . Then $(a_1, \dots, a_{p-1}, a_p), (C_1, \dots, C_{p-1}, C_p)$ satisfies (4). Now we show a local expansion property:

(5) If $A \subseteq V(G)$ is stable, then there are at least $T^{k-1}L^{3(k-1)}D|A|$ vertices that belong to or have a neighbour in A .

Let S be the set of all vertices that belong to or have a neighbour in A . From the inductive hypothesis applied to $G \setminus S$, we obtain a stable set, and its union with A is stable; so

$$\frac{|G| - |S|}{T^{k-1}(\log(|G| - |S|))^{3(k-1)}D} + |A| < \frac{|G|}{T^{k-1}L^{3(k-1)}D}.$$

Since $\log(|G| - |S|) \leq L$, this implies

$$\frac{|G| - |S|}{T^{k-1}L^{3(k-1)}D} + |A| < \frac{|G|}{T^{k-1}L^{3(k-1)}D},$$

which simplifies to the claim. This proves (5).

From (4), taking $p = t$ and $q = 1/(4t^2)$, we deduce that, since $4t^4 \leq 3200t^4 = 200T$, G contains a star system of length t , size at least $\frac{d}{4T^{k-2}D^{3k-5}}$, and sparsity at most $1/(4t^2)$. Let $(a_1, \dots, a_t), (B_1, \dots, B_t)$ be such a star system. Thus, from the second statement of (1), each B_i has cardinality at least

$$\frac{d}{4T^{k-2}D^{3k-5}} \geq \frac{dT D^3}{4T^{k-1}L^{3(k-1)}D} \geq \frac{dT D^3}{d+1} \geq 8t^4 D^3.$$

Let X_1 be the set of vertices adjacent to one of a_1, \dots, a_t (so $|X_1| \leq td$), and let X_2 be the set of vertices $v \in V(G) \setminus X_1$ such that for some $i \in \{1, \dots, t\}$, v has at least $|B_i|/(2t^2L^2)$ neighbours in B_i . Thus (counting edges incident with $V(G) \setminus X_1$), we have $|X_2| \leq 2t^3L^2d$. Let $Y \subseteq V(G)$ be some subset of $V(G)$ such that:

- $Y \cap X_1 \subseteq B_1 \cup \dots \cup B_t$ and $Y \cap X_2 = \emptyset$;
- $|Y \cap X_1| \leq t^2$, and $|Y| \leq \frac{1}{2}t^2L^2$.

Let us call such a set Y an *obstruction set*. For the moment, let us fix an obstruction set Y , and let X_3 be the set of vertices of G that are equal or adjacent to a vertex in Y .

(6) $|X_3| \leq \frac{1}{2}t^2L^2d$, and for $1 \leq i \leq t$, $|X_3 \cap B_i| \leq |B_i|/2$.

The first statement is clear, since $\Delta(G) \leq d$. The vertices of $X_3 \cap B_i$ are of three types: those in $Y \cap B_i$, those with a neighbour in $(Y \setminus B_i) \cap (B_1 \cup \dots \cup B_t)$, and those with a neighbour in $Y \setminus (B_1 \cup \dots \cup B_t)$. Let $|Y \cap B_i| = n$; then the number of the second type is at most $(t^2 - n)|B_i|/(4t^2)$ (since B_i is stable and the star system is $1/(4t^2)$ -sparse). So the number of vertices of the first or second type is at most $n + (t^2 - n)|B_i|/(4t^2) \leq |B_i|/4$ (since $|B_i| \geq 8t^4 D^3 \geq 4t^2$). Since Y is disjoint from X_2 , there are at most $|Y| \cdot |B_i|/(2t^2 L^2) \leq |B_i|/4$ vertices of the third type. Consequently $|X_3 \cap B_i| \leq |B_i|/2$. This proves (6).

For $1 \leq i \leq t$ and each integer $r \geq 0$, let N_i^r be the union of the vertex sets of all paths of G with length at most r that have one end in $B_i \setminus X_3$ and have no other vertices in $X_1 \cup X_2 \cup X_3$.

(7) For $1 \leq i \leq t$, $|N_i^1| \geq 3t^3 L^3 d$.

Let $S = N_i^1 \cup X_1 \cup X_2 \cup X_3$. Since

$$|B_i \setminus X_3| \geq |B_i|/2 \geq \frac{d}{8T^{k-2}D^{3k-5}}$$

by (6), and since S contains all vertices that belong to or have a neighbour in $B_i \setminus X_3$, (5) implies that

$$|S| \geq T^{k-1} L^{3(k-1)} D |B_i \setminus X_3| \geq \frac{dT^{k-1} L^{3(k-1)} D}{8T^{k-2} D^{3k-5}} = \frac{dT L^{3(k-1)}}{8D^{3k-6}} \geq 2t^4 L^3 d.$$

Since

$$|X_1 \cup X_2 \cup X_3| \leq td + 2t^3 L^2 d + \frac{1}{2} t^2 L^2 d \leq 3t^3 L^2 d,$$

it follows that

$$|N_i^1| \geq 2t^4 L^3 d - 3t^3 L^2 d \geq 3t^3 L^3 d.$$

This proves (7).

(8) For $1 \leq i \leq t$ and each integer $r \geq 2$, if $|N_i^{r-1}| \leq |G|/2$ then

$$|N_i^r| \geq (1 + 2/L) |N_i^{r-1}|.$$

Let $|N_i^{r-1}| = m$ and let $S = N_i^r \cup X_1 \cup X_2 \cup X_3$. Let $A \subseteq N_i^{r-1}$ be stable with size at least $\frac{m}{T^{k-1}(\log m)^{3(k-1)}D}$. Since S contains all vertices that belong to or have a neighbour in A , (5) implies that

$$|S| \geq T^{k-1} L^{3(k-1)} D |A| \geq \frac{L^{3(k-1)}}{(\log m)^{3(k-1)}} m.$$

Since $\log m \leq L - 1$, we deduce that

$$|S| \geq (1 + 1/L)^{3(k-1)} m \geq (1 + 3/L)m.$$

As before, $|X_1 \cup X_2 \cup X_3| \leq 3t^3 L^2 d \leq m/L$, since $m \geq 3t^3 L^3 d$ (because $N_i^1 \subseteq N_i^{r-1}$ and $|N_i^1| \geq 3t^3 L^3 d$ by (7)). Since $N_i^r = S \setminus (X_1 \cup X_2 \cup X_3)$, it follows that

$$|N_i^r| \geq (1 + 3/L)m - m/L = (1 + 2/L)m.$$

This proves (8).

We recall that earlier we fixed some obstruction set Y , and used it to define X_3 . We deduce that:

(9) *For every obstruction set Y , and for all distinct $i, j \in \{1, \dots, t\}$, there is an induced path of length at most $L^2 - 2$ between B_i, B_j with no vertices in X_2 and with no internal vertices in X_1 , such that none of its vertices belong to or have a neighbour in Y .*

Define X_3 as before. Since $(1 + 2/L)^{L/2} \geq 2$, it follows that $(1 + 2/L)^{L^2/2} \geq 2^L = |G|$. Let $r = \lfloor L^2/2 \rfloor - 1$. Then $r \geq L^2/2 - 2$, and so $(1 + 2/L)^{r+2} \geq |G|$. From the last statement of (1), $(1 + 2/L)^2 < 1/2$, and so $(1 + 2/L)^r \geq |G|/2$. From (8), it follows that $|N_i^r| > |G|/2$, and similarly $|N_j^r| > |G|/2$, and so $N_i^r \cap N_j^r \neq \emptyset$. This proves (9).

Let $[t]^{(2)}$ be the set of all two-element subsets of $\{1, \dots, t\}$. Choose $I \subseteq [t]^{(2)}$ maximal such that for each $\{i, j\} \in I$, there is a path $P_{ij} = P_{ji}$ of G with the following properties:

- for each $\{i, j\} \in I$, P_{ij} has one end in B_i and the other in B_j , and has no vertices in X_2 and no internal vertices in X_1 ;
- each $P_{i,j}$ is induced and has length at most $L^2 - 2$; and
- for all distinct $\{i, j\}, \{i', j'\} \in I$, the paths P_{ij} and $P_{i'j'}$ are vertex-disjoint and there are no edges between their vertex sets.

Let Y be the union of the vertex sets of all the paths P_{ij} ; then Y is an obstruction set. From (9) and the maximality of I , it follows that $I = [t]^{(2)}$, and so G has an induced subgraph that is a subdivision of K_t with lengths at least three and at most $(\log |G|)^2$, a contradiction. This proves 3.1. \blacksquare

Since $\Delta(G) \leq |G|$, we may replace d by $|G|$, and deduce:

3.2 *Let G be a graph with $|G| \geq 2$, let $k \geq 1$ be an integer with $\omega(G) \leq k$, and let $t \geq 3$ be an integer such that no induced subgraph of G is a subdivision of K_t with lengths between 3 and $(\log |G|)^2$. Then*

$$\alpha(G) \geq \frac{|G|}{(2t)^{4(k-1)}(\log |G|)^{3k-2}}.$$

Since a subdivision of K_t with lengths at least two contains a subdivision of any t -vertex graph, we could replace K_t by a general graph H , and deduce a strengthened version of 1.7:

3.3 *Let G be a graph with $|G| \geq 2$, let $k \geq 1$ be an integer with $\omega(G) \leq k$, and let H be a graph with t vertices, such that no induced subgraph of G is a subdivision of H with lengths between 3 and $(\log |G|)^2$. Then*

$$\alpha(G) \geq \frac{|G|}{(2t)^{4(k-1)}(\log |G|)^{3k-2}}.$$

If we drop the lower bound on the lengths of the subdivision, then this automatically excludes large cliques, so we can omit the condition that $\omega(G) \leq k$. We obtain a strengthened form of 1.1:

3.4 *Let G be a graph with $|G| \geq 2$, and let $t \geq 3$ be an integer such that no induced subgraph of G is a subdivision of K_t with length at most $(\log |G|)^2$. Then*

$$\alpha(G) \geq \frac{|G|}{(2t)^{4(t-2)}(\log |G|)^{3t-5}}.$$

4 Excluding trees and excluding subdivisions

So we have a pair of theorems: for every forest H , all H -free graphs with bounded clique number have a nearly-linear stable set (proved in [16]), and for every graph H , all H -subdivision-free graphs with bounded clique number have a nearly-linear stable set (proved in this paper). There have been previous examples of such pairings between a theorem about excluding a forest, and a theorem about excluding subdivisions. Here is one, between results concerning the “strong Erdős-Hajnal property”:

4.1 [3] *For every forest H there exists $\varepsilon > 0$ such that in every H -free graph G with $|G| \geq 2$ and maximum degree at most $\varepsilon|G|$, there exist disjoint subsets A, B of $V(G)$ with $|A|, |B| \geq \varepsilon|G|$ such that there are no edges between A, B .*

4.2 [4] *For every graph H there exists $\varepsilon > 0$ such that in every H -subdivision-free graph G with $|G| \geq 2$ and maximum degree at most $\varepsilon|G|$, there exist disjoint subsets A, B of $V(G)$ with $|A|, |B| \geq \varepsilon|G|$ such that there are no edges between A, B .*

Let $\tau(G)$ denote the maximum t such that G contains a subgraph isomorphic to $K_{t,t}$. Here is a second pairing:

4.3 [12] *For every forest H , there is a function f such that every H -free graph G has degeneracy at most $f(\tau(G))$.*

4.4 [14] *For every graph H , there is a function f such that every H -subdivision-free graph G has degeneracy at most $f(\tau(G))$.*

And a third pairing, between polynomial versions of the two previous results:

4.5 [19, 11] *For every forest H , there is a polynomial f such that every H -free graph G has degeneracy at most $f(\tau(G))$.*

4.6 [1, 9, 11] *For every graph H , there is a polynomial f such that every H -subdivision-free graph G has degeneracy at most $f(\tau(G))$.*

A fourth pairing is between the Gyárfás-Sumner conjecture 1.3 and Scott’s false conjecture 1.5 (somewhat worrying for those who believe in the predictive value of such pairings and in the truth of the Gyárfás-Sumner conjecture.)

In view of this, one would ask whether any of these four pairings can be unified. Let H be a graph and let T be a forest of H . We say that a graph G is $H(T)$ -subdivision-free if no induced subgraph of G is isomorphic to a subdivision of H in which the edges of T are not subdivided. This provides a common weakening of the two previous hypotheses (being T -free and being H -subdivision-free), and one might hope to unify the pairs on these lines. For three of the four pairings above this is not known. (In [4] a version of 4.2 was proved for $H(T)$ -subdivision-free graphs, but only when T is a path of H , not a general forest.) But we have been able to prove a (numerically somewhat weaker) unification of 1.1 and 1.4:

4.7 For every graph H , and every tree T of H with radius at most $r \geq 0$, and every $k \geq 2$, let $q = (r + 1)(k - 1)$; then there exists $b > 0$ such that every $H(T)$ -subdivision-free graph G with clique number at most k and maximum degree at most d satisfies

$$\alpha(G) \geq \frac{|G|}{d^{b(\log d)^{-\frac{1}{q}}} (\log |G|)^b}.$$

The proof contains no new ideas, and is a (rather messy) combination of the proofs of 1.1 and 1.4, so we will skip it. It will appear in detail in [15].

Finally, there is an intriguing conjecture of Du and McCarty [5].

4.8 Conjecture: For every class \mathcal{H} of graphs closed under taking induced subgraphs, if there is a function f such that every $G \in \mathcal{H}$ has degeneracy at most $f(\tau(G))$, then $\chi(G) \leq |G|^{o(1)}$ for every triangle-free $G \in \mathcal{H}$.

Our theorems 1.4 and 1.1 both imply special cases of this conjecture, since the existence of functions f as in 4.8 is proved in [19, 14] respectively.

References

- [1] R. Bourneuf, M. Bucić, L. Cook and J. Davies, “On polynomial degree-boundedness”, *Advances in Combinatorics* **2024:5**, 16pp, <https://arxiv.org/abs/2311.03341>.
- [2] J. Chalopin, L. Esperet, Z. Li and P. Ossona de Mendez, “Restricted frame graphs and a conjecture of Scott”, *Electronic J. Combinatorics* **23** (2016), #P1.30, [arXiv:1406.0338](https://arxiv.org/abs/1406.0338).
- [3] M. Chudnovsky, A. Scott, P. Seymour and S. Spirkl, “Pure pairs. I. Trees and linear anticomplete pairs”, *Advances in Math.*, **375** (2020), 107396, [arXiv:1809.00919](https://arxiv.org/abs/1809.00919), <https://doi.org/10.1016/j.aim.2020.107396>.
- [4] M. Chudnovsky, A. Scott, P. Seymour and S. Spirkl, “Pure pairs. II. Excluding all subdivisions of a graph”, *Combinatorica* **41** (2021), 279–405, [arXiv:1804.01060](https://arxiv.org/abs/1804.01060).
- [5] Xiyang Du and Rose McCarty, “A survey of degree-boundedness”, *European Journal of Combinatorics* (2024), p. 104092, [arXiv:2403.05737](https://arxiv.org/abs/2403.05737).
- [6] P. Erdős, “Graph theory and probability”, *Canad. J. Math.* **11** (1959), 34–38.
- [7] J. Fox and J. Pach, “Applications of a new separator theorem for string graphs”, *Combinatorics, Probability and Computing* **23** (2014), 66–74.
- [8] J. Fox, J. Pach and A. Suk, “Quasiplanar graphs, string graphs, and the Erdős-Gallai problem”, *European J. Comb.* **119** (2024), 103811.
- [9] A. Girão and Z. Hunter, “Induced subdivisions in $K_{s,s}$ -free graphs with polynomial average degree”, [arXiv:2310.18452](https://arxiv.org/abs/2310.18452).
- [10] A. Gyárfás, “On Ramsey covering-numbers”, *Coll. Math. Soc. János Bolyai*, in *Infinite and Finite Sets*, North Holland/American Elsevier, New York (1975), 10.

- [11] Z. Hunter, A. Milojević, B. Sudakov and I. Tomon, “Kővári-Sós-Turán theorem for hereditary families”, (2024), *J. Combinatorial Theory Ser. B* **172** (2025), 168–197, [arxiv:2401.10853](https://arxiv.org/abs/2401.10853).
- [12] H. A. Kierstead and S. G. Penrice, “Radius two trees specify χ -bounded classes”, *J. Graph Theory* **18** (1994), 119–129.
- [13] T. Korhonen and D. Lokshtanov, “Induced-minor-free graphs: Separator theorem, subexponential algorithms, and improved hardness of recognition”, *Proceedings of the 2024 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA)*, Society for Industrial and Applied Mathematics, 2024, [arXiv:2308.04795](https://arxiv.org/abs/2308.04795).
- [14] D. Kühn and D. Osthus. “Induced subdivisions in $K_{s,s}$ -free graphs of large average degree”, *Combinatorica* **24** (2004), 287–304.
- [15] T. Nguyen, *Induced Subgraph Density*, Ph.D. thesis, Princeton University, May 2025, in preparation.
- [16] T. Nguyen, A. Scott and P. Seymour, “Trees and near-linear stable sets”, manuscript May 2024, [arXiv:2409.09397](https://arxiv.org/abs/2409.09397).
- [17] A. Pawlik, J. Kozik, T. Krawczyk, M. Lasoń, P. Micek, W. T. Trotter and B. Walczak, “Triangle-free intersection graphs of line segments with large chromatic number”, *J. Combinatorial Theory, Ser. B* **105** (2014), 6–10.
- [18] A. Scott, “Induced trees in graphs of large chromatic number”, *J. Graph Theory* **24** (1997), 297–311.
- [19] A. Scott, P. Seymour and S. Spirkl, “Polynomial bounds for chromatic number. I. Excluding a biclique and an induced tree”, *J. Graph Theory* **102** (2023), 458–471. [arXiv:2104.07927](https://arxiv.org/abs/2104.07927), <https://doi.org/10.1002/jgt.22880>.
- [20] D.P. Sumner, “Subtrees of a graph and chromatic number”, in *The Theory and Applications of Graphs*, (G. Chartrand, ed.), John Wiley & Sons, New York (1981), 557–576.
- [21] B. Walczak, “Triangle-free geometric intersection graphs with no large independent set”, *Discrete & Combinatorial Geometry* **53** (2015), 221–225.