

Links from complex spikes to local plasticity and motor learning in the cerebellum of awake-behaving monkeys

Javier F Medina¹ & Stephen G Lisberger²

The hypothesis of cerebellar learning proposes that complex spikes in Purkinje cells engage mechanisms of plasticity in the cerebellar cortex; in turn, changes in the cerebellum depress the simple-spike response of Purkinje cells to a given stimulus and cause the adaptive modification of a motor behavior. Many elements of this hypothesis have been supported by prior experiments, and correlations have been found between complex spikes, simple-spike plasticity and behavior during the learning process. We carried out a trial-by-trial analysis of Purkinje cell responses in awake-behaving monkeys and found evidence for a causal role for complex spikes in the induction of cerebellar plasticity during a simple motor learning task. We found that the presence of a complex spike on one learning trial was linked to a substantial depression of simple-spike responses on the subsequent trial, at a time when behavioral learning was expressed.

The cerebellum possesses two input systems that are very different in both anatomy and physiology^{1,2}. The mossy fiber system originates in many brain regions and projects to Purkinje cells in the cerebellar cortex through granule cells and their parallel fibers, as well as through a number of different inhibitory interneurons. Each Purkinje cell receives inputs from tens of thousands of parallel fibers, resulting in traditional simple spikes with spontaneous rates of 100 s⁻¹ or higher. The climbing fiber system originates in the inferior olive. Each Purkinje cell receives massive inputs from just one climbing fiber, resulting in complex spikes that have an unusual waveform and occur infrequently, about 1 s⁻¹. The notable differences between the two systems led to the idea that their interplay may create a learning system for movement³⁻⁵.

The theory of cerebellar learning is comprised of three separate, but connected, hypotheses. First, climbing fiber inputs are activated when a movement is inaccurate or erroneous^{6,7}. Second, the activation of the climbing fiber input engages mechanisms of plasticity that cause changes in synaptic strength and alter the simple-spike responses of Purkinje cells^{8,9}. Third, these changes in cerebellar output lead to adaptive modification of the motor behavior¹⁰⁻¹⁴. Each of these three hypotheses has received considerable experimental support, but the theory of cerebellar learning has not been tested as a full entity by examining all three hypotheses simultaneously in the same experiment. In addition, it has been difficult to draw strong conclusions about cause and effect links between complex spikes, simple spikes and behavioral learning because previous work has treated plasticity in the cerebellar cortex as a static process by measuring the simple-spike response of Purkinje cells after learning, at a time when the climbing fiber inputs are no longer activated. One line of work has supported the theory by demonstrating that arm movement errors evoke complex spikes^{51,52}

and that subsequent learned changes in motor behavior are associated with suitable changes in simple spike responses⁵¹. This work demonstrates a strong correlation, but stops short of showing cause-and-effect links between individual complex spikes, changes in simple spikes and behavioral learning. A more rigorous test of the cerebellar learning theory demands that we examine the dynamic interaction between complex spikes and the induction of neural and behavioral learning on a trial-by-trial basis during the learning process itself.

We developed a new approach that tests directly for cause and effect links from complex spikes in single trials to changes in simple-spike firing, and in turn to behavioral learning in awake-behaving monkeys. While monkeys were actively learning smooth pursuit eye movements¹⁵, we recorded from single Purkinje cells in the floccular complex of the cerebellum, a structure that is only two synapses removed from motor neurons and is critical for both the initiation and maintenance of pursuit^{16,17}. We found that the presence or absence of a complex spike on one learning trial determines whether or not the simple-spike response of the Purkinje cell will be depressed on the subsequent trial, and we show with the same dataset how complex-spike responses during learning are related to changes in simple-spike responses that cause behavioral learning.

RESULTS

Even while simultaneously investigating the full set of linkages among complex spikes, learned changes in simple-spike responses and learning in pursuit behavior, we needed to separate the questions of how complex spikes during learning lead to changes in simple-spike responses and how the changes in simple-spike responses lead to changes in behavior. This is because the three variables that we measure

¹Department of Psychology, University of Pennsylvania, 3720 Walnut Street, Philadelphia, Pennsylvania 19104, USA. ²Howard Hughes Medical Institute, W. M. Keck Foundation Center for Integrative Neuroscience, and Department of Physiology, University of California San Francisco, 513 Parnassus Avenue, San Francisco, California 94143, USA. Correspondence should be addressed to S.G.L. (sgl@phy.ucsf.edu).

Received 6 June; accepted 31 July; published online 21 September 2008; corrected after print 15 January and 30 April 2009; doi:10.1038/nn.2197




Figure 1 Example trials showing the target motions used to instruct and probe directional learning in pursuit eye movements. (**a–d**) Positive numbers represent motion to the right (horizontal plots) or upwards (vertical plots). Negative numbers represent motion to the left or downwards. Learning trials are shown in **a** and **c**. Gray and black traces indicate data from representative trials early versus late in a learning block. Probe trials are shown in **b** and **d**. Gray and black traces indicate data from representative trials before learning and after at least 100 learning trials. Dashed traces show target motion. Insets in **a** and **b** indicate the target motion in polar coordinates. The arrow pointing down and right in **c** indicates the learned response. The arrow pointing up and left indicates the hardwired visual response to the change in target direction. Note that the scales on vertical eye position and velocity are different in the probe versus the learning trials.

are linked by mechanisms that operate independently, at different loci and at different times. We started by considering the linkage between changes in simple-spike responses and changes in eye velocity that are measured at the same time. Then we evaluated the relationship between the total number of complex spikes fired during learning and the changes in simple-spike response observed after learning for each Purkinje cell. Finally, we analyzed the link between a single complex spike on one learning trial and the changes that occurred in the simple-spike response on the next trial.

Simple spike correlates of directional learning in pursuit

In a typical learning experiment (Fig. 1)¹⁵, a monkey performed 200 learning trials (Fig. 1a,c), each one beginning with target motion to the right at 20 deg s⁻¹. After 250 ms, we introduced an instructive stimulus in the form of a vertical component of target motion at 30 deg s⁻¹ so that the target moved up and to the right for 450 ms before stopping. Starting at about 100 ms after the onset of upward target motion, the monkey began to move his eyes upwards (Fig. 1c) and then made a large saccade to catch up with the target. There is, in addition, a more subtle difference in the vertical eye velocity traces between the early and late learning trials (Fig. 1c). In the late learning trials, an upward vertical eye velocity appeared just before the onset of upward target motion. This early upward eye velocity was a learned response because it preceded the instructive stimulus and because it occurred after, but not before, learning (Fig. 1b,d) in occasional probe trials that present purely rightward target motion.

Consider a learning experiment on a Purkinje cell with a rightward on direction, defined by the direction of pursuit associated with the largest simple-spike response before learning; the off direction was leftward. In pre-learning probe trials (Fig. 2a), the Purkinje cell had a relatively small simple-spike response during pure downward pursuit and had larger and smaller simple-spike responses during pursuit with a small rightward or leftward component, respectively.

In the learning trials of learning block 1 (Fig. 2b), the target moved downward for 250 ms and then changed direction abruptly to include a leftward (off direction) component of motion at 30 deg s⁻¹. After 100 learning trials, occasional probe trials consisting of purely

downward target motion evoked a leftward component of smooth eye motion that began about 200 ms after the onset of target motion, anticipating the change of target direction that would have occurred 50 ms later in the learning trials (Fig. 2b). In the first learning block, the change in target direction was randomly selected to be either in the on or off direction of the Purkinje cell; we always proceeded to a second learning block that evoked learning in the opposite direction (Fig. 2c). For the example Purkinje cell, behavioral learning was modeled by a markedly similar, reciprocal change in simple-spike firing rate for both directions of learning (Fig. 2b,c). In subsequent experiments, we capture the learned eye velocity and simple-spike firing rate as the millisecond-by-millisecond difference between the average response in the learning block minus the pre-learning block.

A large majority of the Purkinje cells in our sample (as indicated in Fig. 3) showed learned changes in simple-spike firing that had the same shape and direction as the changes in eye velocity ($n = 49$, group 1; Fig. 3c,e). For group 1 Purkinje cells, the learned change in eye velocity lagged the learned change in simple-spike firing by a small time delay of 36 and 28 ms for on-direction and off-direction learning, respectively (Fig. 3c). For the group 1 Purkinje cells, the average peak-to-peak changes in simple-spike firing and eye velocity were approximately 30 spikes s⁻¹ and 12 deg s⁻¹, respectively, for a ratio of 2.5 spikes s⁻¹ per deg s⁻¹.

For a minority of our sample, the learned changes in simple-spike firing and eye velocity were not reciprocal ($n = 11$, group 2; Fig. 3d,f). The most notable deviation of the learned simple-spike responses of group 2 Purkinje cells was a delayed increase in firing for off-direction learning. The difference between the learned responses of group 1 and 2 Purkinje cells could not be attributed to differences in the learned eye velocities (Fig. 3e,f). The increases in simple-spike firing of group 2 Purkinje cells modeled the positive portion of the learned eye acceleration (Fig. 3d): that is, when the eyes were speeding up in the on direction or slowing down in the off direction. The learned simple-spike responses of group 1 and 2 Purkinje cells divide neatly in terms of relationships to learned eye velocity versus acceleration. However, the same clean division was not observed in the pre-learning block of trials used to measure the directional selectivity of each Purkinje cell; during pre-learning trials in either the on or off direction, all of the Purkinje cells in both groups showed an eye velocity component of firing during steady-state pursuit and many in each group showed a transient during eye acceleration at the initiation of pursuit (see also refs. 18,19).

From instructive complex spikes to simple-spike learning

Consistent with previous work, we found complex-spike responses to image motion in the off direction^{20,21}: that is, when the eyes are moving more slowly than the target in the off direction or faster than the target in the on direction. During off-direction learning trials, however, group 1 and group 2 Purkinje cells differed in the probability of emitting a




Figure 2 Effects of learning on responses of one Purkinje cell. (a–c) From top to bottom, the three rows show the control block and the learning blocks in the off and on direction of the Purkinje cell. Left, the colored arrows show different target motions and the black continuous curve shows the direction tuning for the Purkinje cell. Middle and right, eye velocity and simple-spike firing rate during probe trials. The colors of the traces refer to the different target trajectories indicated by the colored arrows in the polar plots. For ease of comparison, the light blue traces in **b** and **c** replot the pre-learning data for downward target motion from **a**. The ribbons show the mean \pm s.e.m. of eye velocity and firing rate.

complex spike (Fig. 3a,b). The probability of a complex spike in the interval 75–175 ms after the change in target direction increased to an average of 0.5 in group 1 versus 0.2 in group 2. As eye (and image) velocity was essentially the same on trials with and without a complex spike in our experiments (Supplementary Fig. 1 online), the occurrence of a complex spike on any particular learning trial reflects its stochastic nature and not a large trial-by-trial difference in the magnitude of image motion caused by the instructive change in target direction. During on-direction learning trials, complex-spike responses showed relatively little modulation (Figs. 3a,b), except for a prominent peak that will be considered later because it occurred 125 ms after the offset of target motion (more than 800 ms after the onset of target motion).

To quantitatively test the relationship between off-direction learning in the simple-spike firing of each Purkinje cell and the probability of emitting a complex spike in response to the learning stimulus, it is

necessary to appreciate the relative timings of the learned eye velocity and simple-spike firing in probe trials and the complex-spike responses in learning trials. Complex-spike responses were time-locked to the instructive visual stimulus. They occurred \sim 125 ms after the change in target direction^{20,21} and were measured from 75–175 ms after the change in target direction in learning trials (Fig. 3a,b). The learned eye velocity and simple-spike firing anticipate the change in target direction¹⁵. They started 50–100 ms before the time of the change in target direction in learning trials and could be measured in either learning or probe trials, in the interval from 50 ms before to 50 ms after the time when the change in target direction occurred in the learning trials (Fig. 3c,e).

Across Purkinje cells, the probability of a complex spike during learning was a good predictor of the size and direction of the learned change in simple-spike firing in each individual Purkinje cell (Fig. 4). For learning in the off direction, group 1 Purkinje cells had a high probability of emitting a complex spike in the learning trials and showed large learned decreases in simple-spike firing in probe trials. Group 2 Purkinje cells were less likely to emit a complex spike during learning and showed learned increases, rather than decreases, in




Figure 3 Relationship between complex-spike probability in learning trials and learned simple-spike response in probe trials for group 1 and group 2 Purkinje cells. (a,b) Probability of a complex spike during the first 50 learning trials in a learning block (measured in 100-ms bins). (c,d) Learning-induced change in simple-spike response, calculated as the mean firing rate in probe trials after learning minus the mean firing rate in pre-learning probe trials. In **c**, the black traces repeat the eye velocity averages from **e**. In **d**, the black traces show eye acceleration for the eye velocity traces in **f**, where the negative eye accelerations are shown as dashed traces. (e,f) Learned eye movement, measured as the component of the eye velocity orthogonal to the direction of the target in probe trials. Blue and red traces indicate data for on-direction and off-direction learning, respectively. The ribbons indicate mean \pm s.e.m. Vertical dashed lines show the time when the target changed direction in learning trials. Shaded areas show the analysis intervals for complex-spike responses (a,b) and for simple-spike responses and eye velocity (c,e). Data were averaged across 49 group 1 and 11 group 2 Purkinje cells.




Figure 4 Quantitative relationship between complex-spike probability in learning trials and learned changes in simple-spike activity in probe trials. Each point plots the results of one learning experiment on one Purkinje cell and shows averages across all of the trials in the learning block. Analysis intervals for the complex-spike probability and the learned change in simple-spike firing are indicated in **Figure 3**. Filled and open symbols indicate data from group 1 ($n = 11$) and group 2 ($n = 6$) Purkinje cells, respectively. Blue and red symbols indicate learning in the on-direction and off-direction of the Purkinje cell under study, respectively. Black symbols indicate mean and s.d. across Purkinje cells.

that the probability of a complex spike had to at least triple from baseline to be consistently associated with a learned decrease in simple-spike response.

If complex spike-instructed cellular changes were the only plasticity mechanism operating in the cerebellar cortex, then we would have expected zero learning in simple-spike firing when complex-spike probability was at baseline levels. Our data contradicts this prediction (**Fig. 4**). When the probability of a complex spike is at baseline levels, and thus cannot possibly be signaling 'errors', there are consistent learned increases in simple-spike responses for on-direction learning. Therefore, we suggest that complex spike-instructed cellular changes cannot be the only plasticity mechanism at work during pursuit learning.

The tight correlation between the probability of a complex spike and the amount of neural and behavioral learning is emphasized by comparison of the time courses of different response measures during off-direction learning for group 1 versus group 2 Purkinje cells. For all Purkinje cells, the learned eye velocity (**Fig. 5a**) changed quickly in the first three sets of ten learning trials and then reached an asymptote. Considering each set of learning trials separately, simple-spike firing (**Fig. 5b**) decreased the most at times when the complex-spike probability (**Fig. 5c**) was higher than threefold of the normal baseline level of about 0.1 in a 100-ms bin, for example in the first three sets of ten learning trials for group 1 Purkinje cells. When the complex-spike probability was lower than 0.3 during the learning trials, including all times for group 2 Purkinje cells, we observed learned increases in simple-spike responses.

Simple-spike learning after single complex spikes

The results presented so far demonstrate impressive correlations between the learned

changes in simple-spike firing and eye velocity and between the complex-spike activity in learning trials and the subsequent changes in the simple-spike firing of each Purkinje cell. Still, these data treat learning as a static process, comparing average responses across multiple trials that occurred in different phases of the learning process. To move from correlation to cause and effect using the same dataset, we examined changes in simple-spike responses between two successive learning trials. If the presence (but not the absence) of a complex-spike response to the instructive stimulus on one trial were associated with a reduction in simple-spike firing rate at the correct time on the subsequent trial, then we could conclude that there is a tight link between complex-spike responses and depression of simple-spike responses during behavioral learning.

We looked for complex spike-linked trial-over-trial changes in simple-spike responses in the first 50 trials of off-direction learning, where most of the learning takes place and where the learning stimulus evokes complex-spike responses with a high probability, at least for group 1 Purkinje cells (**Figs. 3–5**). We evaluated pairs of successive learning trials (see Methods), characterizing each according to the presence or absence of a complex spike in the interval of 75–175 ms after the change in target direction. We retained pairs that lacked a complex spike in both trials of the pair ('0-0') or that had a complex spike only in the first trial of the pair ('1-0') and computed the difference in simple-spike firing between the first and second trial of the pair (**Fig. 6**). We looked for depression of simple-spike responses at the same time when learning normally is expressed, which is just before the time of the change in target direction that serves as an instructive stimulus. When there was a complex spike in the first trial (1-0 pairs), the trial-over-trial change in firing in the simple-spike analysis interval was negative, relative to both the 0-0 pairs and the average of 200 random draws of consecutive learning trials (**Fig. 6b**). The reduction in simple-spike firing for 1-0 pairs lay outside of the ± 2 s.d. envelope from the random-draw analysis for almost the entire analysis window and almost exclusively in the analysis window. In contrast, the trial-over-trial change in simple-spike firing for the 0-0 trials was very similar to the random-draw average. The depression of simple-spike firing between the first and second trials of 1-0 pairs was entirely the result of lower than normal simple-spike activity on the second trial of the pair (**Fig. 6a**). Note that the trial-over-trial depression of simple-spike activity is not related to the well-known post-complex spike pause in simple-spike firing^{6,22}; the latter occurred after the complex spike on the same trial, whereas the former occurred before the time of the complex spike, and on the subsequent trial.

The presence of a complex spike on one trial also affected the learned eye velocity on the subsequent trial, although the effect was small; the change in eye velocity in the 1-0 pairs stayed in the ± 2 s.d. envelope





Figure 5 Time courses of complex spike-related instructive signals and learned changes in eye velocity and simple-spike responses. Each graph divides the first 50 learning trials for off-direction learning in sets of ten and plots the average response measure as a function of the set. (**a,b**) Learned eye velocity and simple-spike firing in probe trials. (**c**) Probability of a complex spike in learning trials. Open and filled symbols show data from group 1 ($n = 11$) and group 2 ($n = 6$) Purkinje cells, respectively.



from the random-draw analysis (Fig. 6c). The changes in simple-spike firing and eye velocity for the 1-0 pairs were approximately 12 spikes s^{-1} and 0.8 deg s^{-1} , respectively, yielding a ratio of 15 spikes s^{-1} per deg s^{-1} (the meaning of this eye-velocity sensitivity ratio, which was sixfold larger than the one we measured earlier (Fig. 3), is treated quantitatively in the **Supplementary Appendix** online). Note that these data represent analyses performed on individual Purkinje cells and then averaged across a population of 13 Purkinje cells (Fig. 6). The trial-over-trial reduction of simple-spike firing in the 1-0 pairs was a feature of most individual Purkinje cells and indeed of most individual pairs of trials (**Supplementary Fig. 2** online).

This data (Fig. 6) verifies a key prediction of the cerebellar learning theory on data obtained during motor learning in a behaving primate: a complex spike in one trial is linked to a measurable and properly

Figure 6 Neural and behavioral learning after single complex spikes emitted during off-direction learning trials. All traces are plotted as a function of time, where the target started to move at $t = 0$, the target changed direction at $t = 250$, and the interval when learning was analyzed is bounded by the two vertical dashed lines. (a) Actual simple-spike firing rate on first and second trials of each pair of successive learning trials computed separately for each of 13 Purkinje cells in our sample and then averaged together. (b,c) Average trial-over-trial change in simple-spike firing (b) and eye velocity (c) for the 0-0 sequences (blue) and the 1-0 sequences (red). The black traces show the average change obtained by performing the trial-over-trial subtraction on 200 randomly chosen sequences of two consecutive learning trials and the gray envelope indicates ± 2 s.d.


timed depression of the simple-spike response on the subsequent trial. The depression of simple-spike activity peaked approximately 100 ms before the time of the complex-spike response on the prior trial, which is consistent with the frequently discussed requirement of an ‘eligibility trace’ for cerebellar long-term depression so that a complex spike can affect the synaptic inputs that were active 100 ms earlier^{23–26}.

An independent confirmation of the trial-over-trial effect of a single complex spike came from the analysis of an unanticipated sharp increase in the probability of complex-spike responses just after the offset of target motion for on-direction learning trials (Fig. 7). At this time, smooth eye velocity was in the on direction of the Purkinje cell (for simple-spike responses) and off direction image motion (the main stimulus for complex-spike responses^{20,21}) occurred because the target was suddenly stationary while the eyes kept moving (Fig. 7a).

The presence of a complex spike after the offset of target motion in one trial caused a depression of the simple-spike response and the eye velocity on the subsequent trial. The depression of simple-spike firing (Fig. 7d) remained outside of the ± 2 s.d. envelope from the random-draw analysis during the analysis window and has a similar time course to that determined previously (Fig. 6b): it started 150 ms before and peaked 100 ms before the offset of target motion. The companion trial-over-trial decrease in eye velocity had two negative peaks, but returned to zero in the middle of the analysis interval (Fig. 7e). The peak trial-over-trial changes in simple-spike firing rate and eye velocity were 9 spikes s^{-1} and 0.6 deg s^{-1} , respectively, for a ratio that again was 15 spikes s^{-1} per deg s^{-1} (Fig. 7d,e).

Finally, we asked whether the complex-spike response at the offset of on-direction target motion were linked to any learning expressed as a

Figure 7 Neural and behavioral learning after single complex spikes were emitted at the offset of target motion in on-direction learning trials. (a–c) Black and red traces indicate average eye velocity (a) and simple-spike firing (c) in the first 10 learning trials and after 100 learning trials. In a, the black dashed trace shows target velocity. The arrow indicates the time when off-direction image motion was present because the eyes were moving in the on direction and the target was stationary. The probability of a complex spike in the learning trials is shown in b. Traces in a–c are averages across 13 Purkinje cells. Legend below a also applies to c. (d,e) Average trial-over-trial change in simple-spike firing (d) and eye velocity (e) for the 0-0 sequences (blue) and the 1-0 sequences (red). The target ceased moving at $t = 0$ and the two vertical dashed lines bound the interval when learning was analyzed. The black traces show the average change obtained by performing the trial-over-trial subtraction on 200 randomly chosen sequences of two consecutive learning trials and the gray envelope indicates ± 2 s.d. (f) Red and black traces show learned changes in simple-spike firing rate and eye velocity. Vertical dashed line indicates the time of the offset of target motion. Horizontal dashed line indicates zero. (g) Quantitative relationship between complex-spike probability and learned changes in simple-spike activity. Blue symbols show data from the offset of target motion in on-direction learning experiments and gray symbols are replotted from Figure 4, showing data from the change in target direction in the off-direction learning experiments.



difference in simple-spike and eye velocity responses before versus after 100 learning trials. It was, even though the offset of target motion was not intended as a learning stimulus. Difference traces computed each millisecond as the average response in late learning trials minus that in early learning trials revealed increases in both simple-spike firing (Fig. 7f) and eye velocity at times that were commensurate with the time of the stimuli that led to complex-spike response: both reached peaks near the time of target motion offset ($t = 700$ ms). The learned eye velocity lagged the learned simple-spike firing by a few milliseconds (as in Fig. 3c). Note that it is difficult to interpret the noise in the traces (Fig. 7f) at times between 400 and 600 ms after the onset of target motion because we were not able to remove the effects of saccades in this interval from the simple-spike firing.

Why do we see a local depression of the simple-spike response on the trial following a complex spike, but see learned increases in simple-spike responses and eye velocity at the offset of on-direction target motion after 100 learning trials? When we plotted the magnitude of the learned change in simple-spike firing as a function of the complex-spike probability 75–175 ms after the offset of on-direction target motion in the learning trials (Fig. 7g), the data for the individual Purkinje cells followed the same relationship found during off-direction learning (Fig. 4). Thus, all our data are consistent with the notion that learning will be associated with an increase in simple-spike firing over the course of a block of learning trials unless complex-spike probability is more than threefold that of baseline levels, in which case the simple-spike responses during the learned behavior will decrease.

DISCUSSION

To understand the neural basis of motor learning, we must examine how learning stimuli, neural plasticity and behavior interact at the systems level. We need to identify all of the loci of learning in a neural system and establish at each site the linkages among three key elements: properly timed instructive signals that indicate errors and activate mechanisms of plasticity, learning-related changes in the firing rate of neurons that provide the output from that site, and the adaptive modification of the behavior. In contrast with prior studies, which restricted themselves to one or two of these links at a time, the major contribution of our study is to demonstrate all three links with a single set of neural and behavioral data obtained by recording cerebellar responses before, during and after learning of smooth pursuit eye movements in awake monkeys.

By evaluating both the trial-over-trial changes that take place during the learning process and the static state of the system before and after learning, we were able to evaluate local complex spike–instructed mechanisms of plasticity in the cerebellar cortex in the larger context of system-level changes in the neural circuits that mediate motor learning in pursuit. Our trial-over-trial analysis revealed that a complex spike error signal on one learning trial was linked to depression of the same Purkinje cell's simple-spike activity on the subsequent trial. The depression was present selectively from 250 to 100 ms before the time in the trial when the complex spike had occurred, which is consistent with the timing of learning in pursuit¹⁵, as well as with a number of prior reports that suggest that a complex spike must cause long-term depression in parallel fiber synapses that were active 100–200 ms before the complex spike^{23–26}. The depression had the same temporal characteristics and amplitude in two different and independent sets of learning trials, providing strong statistical support for our results. We think that the trial-over-trial changes in simple-spike firing are a consequence of plasticity mechanisms in the cerebellar cortex, demonstrated here, to the best of our knowledge, during the learning process in awake, behaving animals for the first time.

Cerebellar learning has often been considered to be synonymous with motor learning, but it is important to remember that a complex circuit of neurons generates pursuit and that a whole concert of plastic changes are probably needed at multiple sites in the circuit to cause stable behavioral learning^{10–12,14}. Even though the trial-by-trial analysis indicated that complex spike–instructed depression of simple-spike activity is one of the local mechanisms contributing to pursuit learning, there must be other mechanisms of plasticity. For example, we found that 100 learning trials caused a learned change in simple-spike responses that was related, in a way that was not predicted by the cerebellar learning theory, to the probability of complex-spike responses during those 100 learning trials. If the complex-spike responses of a Purkinje cell during the learning trials were at baseline levels, or even up to twice baseline levels, then behavioral learning was linked to a learned increase in the simple-spike firing of that particular Purkinje cell. This finding complements, and does not contradict, our trial-over-trial finding of complex spike–linked depression. It suggests that there is a competition where different local mechanisms of potentiation and depression are balanced when the probability of a complex spike is about threefold higher than during baseline conditions. The interplay of multiple plasticity mechanisms is not entirely unexpected. There are examples of cellular plasticity at many sites in the cerebellar cortex^{9,27–29} and previous work has indicated multiple sites of plasticity even in simple forms of cerebellar-dependent motor learning, such as adaptation of the vestibulo-ocular reflex^{30,31} and classical delay conditioning of eyelid responses^{32,33}.

Comparison of eye velocity and simple-spike responses before versus after learning showed, as others have shown before^{34–37}, that learned simple-spike responses could drive the learned eye movement. For this to be a causal link, however, changes in simple-spike firing must cause the changes in behavior and not vice versa. As Purkinje cells normally fire in relationship to smooth eye velocity³⁸ and eye velocity changes during our learning task, the changes in simple-spike firing of group 1 Purkinje cells could result simply from mossy fiber inputs that report the kinematic parameters of the current eye movement³⁹. We think that this scenario is unlikely. First, there is a disynaptic connection from floccular Purkinje cells to extraocular motoneurons⁴⁰ and electrical stimulation in the floccular complex causes smooth eye movement with a latency of 10 ms^{30,41}, making it difficult to argue that a change in simple-spike firing would not drive a change in eye movement. Second eye velocity lagged the simple-spike response of group 1 Purkinje cells by more than 10 ms (Fig. 3c). Therefore, we think that the simple-spike correlates of behavioral learning are primary learned responses that are driving, rather than responding to, the learned behavior.

We also think that the complex spike–linked, trial-over-trial depression in simple-spike firing is much too large to be a consequence of cerebellar inputs that report trial-over-trial changes in eye velocity. Purkinje cells in our sample had an average eye-velocity sensitivity of approximately 2.5 spikes s^{-1} per deg s^{-1} measured during the expression of pursuit learning. Given this measurement, and the fact that the trial-over-trial change in eye velocity had a peak of 0.8 deg s^{-1} , the trial-over-trial depression of simple-spike firing rate would be 0.8×2.5 , or 2 spikes s^{-1} , if Purkinje cells were simply responding to the kinematics of the trial-over-trial change in eye movement. In fact, the trial-over-trial depression of simple-spike firing was 12 spikes s^{-1} . Therefore, the trial-over-trial depression of simple-spike firing cannot be simply reporting the behavioral response (see **Supplementary Appendix** for more information).

The highly stochastic nature of complex-spike activity during learning paints a picture that contrasts with the view that motor learning occurs as a series of gradual steps whose size is proportional to the size

of the error on a particular trial^{42–45} (but see ref. 46). A complex spike occurs, at most, on half of learning trials, but is linked to a large depression of simple-spike firing on the subsequent trial. Perhaps the contribution of individual Purkinje cells to motor learning is a random walk with larger steps in the learning direction than the reverse. It remains to be seen whether this also is true at the population or behavioral levels, a question that may ultimately be answered by applying properly contrived quantitative methods to both behavioral and neural data⁴⁷. It also will be informative to determine whether a dynamic trial-by-trial interplay of potentiation and depression is uncovered when *in vitro* experiments use plasticity protocols that mimic the stochastic nature of the *in vivo* complex-spike responses.

Our study is, to the best of our knowledge, the first to examine trial-over-trial neural changes in awake animals actively engaged in a motor learning task, allowing us to make direct estimates of local cellular plasticity mechanisms engaged during the dynamic learning process. By also evaluating how a block of learning trials changes the state of cerebellar and behavioral outputs, we have begun to assemble an understanding of the interaction between cellular and systems-level mechanisms of motor learning. The approach that we have initiated is built on the foundation provided by earlier studies carried out in anesthetized or paralyzed animals⁴⁸, but goes further because it can assess the relationship between plasticity and behavior simultaneously and because it can be deployed during learning with natural sensory stimuli. We found evidence that strongly supports the predictions of the cerebellar learning theory^{3–5}, while also reminding us that learning in any single behavior probably involves many cellular mechanisms at a variety of loci in the full neural circuit.

METHODS

Animal preparation. We recorded the responses of single Purkinje cells in two awake, behaving rhesus monkeys (*Macaca mulatta*). Monkeys had been instrumented with an implanted head-holder to allow us to immobilize the head in a way that did not cause distress⁴⁹, a coil of wire on one eye to allow the use of the search coil system to monitor eye movement⁴⁹, and a recording cylinder that allowed us to gain access to the floccular complex of the cerebellum with metal microelectrodes¹⁹. These devices were implanted in surgical procedures that used isoflurane as an anesthetic and were conducted with sterile procedure. Monkeys received several days of analgesics following each surgery. All procedures had been approved in advance by the Institutional Animal Care and Use Committee at UCSF and were in accordance with the US National Institutes of Health Guide for the Care and Use of Laboratory Animals.

Pursuit task. Monkeys were trained to fixate and track a target presented on a screen in front of them in exchange for fluid rewards. Targets were created by imaging the beam from a fiber-optic light source and reflecting it off a pair of computer-controlled, moveable mirror galvanometers onto the back of a tangent screen. Experiments were conducted as a series of trials in which each trial started with a brief fixation period (500–1,000 ms), a sequence of target motions was then provided, and the trial was terminated by holding the target stationary for 500 ms. The onset of target motion followed a standard step/ramp trajectory, with the size of the step being adjusted to minimize the occurrence of saccades. Monkeys were required to fixate on the target when it appeared, track it as best they could as it moved during the trial and fixate it again at the end of its motion, keeping eye position in a 2×2 -deg window centered on the moving target. Successful performance on a trial was rewarded with a droplet of fluid. Reward contingencies were suspended for 250 ms whenever the target underwent a change in position or velocity to ensure that the monkey was not punished for the inescapable latencies of the visual system. To generate eye velocity traces, we used an analog circuit to differentiate the eye position records for frequencies below 25 Hz, while filtering out higher

frequencies. Analog signals related to horizontal and vertical target position, eye position, and eye velocity were sampled and digitized at 1 kHz per channel.

Experiments were conducted as three separate blocks. The baseline block comprised approximately 100 trials in which the target moved at constant speed (20 deg s⁻¹) for 750 ms in one of eight directions. It was followed by two learning blocks, each one comprising approximately 300 trials. Most of the trials in a learning block were learning trials that started with 250 ms of target motion at 20 deg s⁻¹ in a direction that was orthogonal to the on direction of the Purkinje cell under study. Then, the target provided an instructional change in direction by acquiring a component of motion at 30 deg s⁻¹ in the on or off direction of the Purkinje cell and continuing to move for an additional 450 ms. The instructional change in target direction was the same in all of the learning trials of a given learning block and was in the on or off direction of the Purkinje cell being studied in the two learning blocks. To assess the extent of learning, we randomly interleaved approximately 20–30 ‘probe’ trials among the learning trials; in probes, the target began to move as in the learning trials, but did not change direction.

Single-unit recording. Platinum-iridium microelectrodes manufactured in our laboratory were introduced daily and driven through the cerebral cortex and the tentorium of the cerebellum until they reached the floccular complex. Entrance in to the cerebellum was heralded by a large increase in the general level of spontaneous activity and the presence of complex-spike responses. Entrance into the floccular complex was signaled by the presence of eye movement-related activity and the sound of Purkinje cells modulating strongly as the monkey tracked sinusoidal target motion along the horizontal or vertical axis. Extracellular activity was passed through a standard head stage, amplified, filtered (bandpass, 100 Hz to 10 kHz), digitized at 25 KHz and stored in the computer for further processing. Because it was challenging to keep the complex-spike responses isolated throughout a full experiment, we were able to include only 17 Purkinje cells in our analysis of complex-spike responses. However, we were able to document the learned simple-spike responses of 60 Purkinje cells that were held through the full learning protocol, with each Purkinje cell being identified by the presence of a clear complex-spike response for part of the recording.

Data analysis. Digitized eye velocity traces from individual trials were displayed on the computer screen and all saccades were excised manually. We then used a software window discriminator to go through each trial in the dataset and marked simple-spikes and complex spikes by hand. The simple-spike response was transformed into a firing rate using the reciprocal interval algorithm⁵⁰. Complex-spike responses were accumulated for many trials in bins with widths of 100 ms and then converted to the probability of a complex spike in a particular bin.

We also used a new trial-by-trial analysis derived from a previous study⁴⁷. We characterized each pair of successive learning trials according to the presence or absence of a complex spike in the interval from 75 to 175 ms after the change in target direction. We retained pairs that lacked a complex spike in both trials of the pair (‘0-0’) or that had a complex spike only in the first trial of the pair (‘1-0’), but only if the learning trial immediately before the pair had lacked a complex spike. The requirement for the absence of a complex spike in a learning trial immediately before the pair reduced the total number of sequences that met all criteria, but was necessary to establish a consistent history of visual stimulus and complex-spike response for every sequence included in our analysis. We then computed the change in simple-spike firing rate and in the eye velocity between the trials of each 1-0 and 0-0 pair. The trial-over-trial changes in simple-spike firing rate and eye velocity were averaged separately for each Purkinje cell as long as there were at least four ‘1-0’ and four ‘0-0’ pairs (Figs. 6 and 7); the averages were then pooled to provide a grand average of trial-by-trial changes across Purkinje cells.

To assess statistical significance, we generated a series of 100 control distributions of trial-over-trial changes for each Purkinje cell separately. Each distribution contained 200 random sequences of two consecutive learning trials drawn from all available sequences (including 0-0, 0-1, 1-0 and 1-1 pairs). The mean of the control distributions represents the average trial-over-trial change that we would expect for each Purkinje cell over two consecutive learning trials,

regardless of whether there was a complex spike in any of the trials or not. The s.d. of the control distributions allowed us to assess statistical significance by asking whether the measured trial-over-trial changes for 1-0 and 0-0 pairs lay outside the 95% confidence intervals.

Note: Supplementary information is available on the Nature Neuroscience website.

ACKNOWLEDGMENTS

This work was supported by the Howard Hughes Medical Institute and by US National Institutes of Health grant P50 MH77970.

AUTHOR CONTRIBUTIONS

J.F.M. conducted recordings from the floccular complex and analyzed data. J.F.M. and S.G.L. wrote the manuscript. S.G.L. provided advice, guidance and suggestions throughout.

Published online at <http://www.nature.com/natureneuroscience/>

Reprints and permissions information is available online at <http://npg.nature.com/reprintsandpermissions/>

- Eccles, J.C., Ito, M. & Szentágothai, J. *The Cerebellum as a Neuronal Machine* (Springer-Verlag, Berlin, New York, 1967).
- Ito, M. *The Cerebellum and Neural Control* (Raven Press, New York, 1984).
- Albus, J.S. A theory of cerebellar function. *Math. Biosci.* **10**, 25–61 (1971).
- Marr, D. A theory of cerebellar cortex. *J. Physiol. (Lond.)* **202**, 437–470 (1969).
- Ito, M. Neural design of the cerebellar motor control system. *Brain Res.* **40**, 81–84 (1972).
- Simpson, J.I., Wylie, D.R. & de Zeeuw, C.I. On climbing fiber signals and their consequence(s). *Behav. Brain Sci.* **19**, 384–398 (1996).
- de Zeeuw, C.I. *et al.* Microcircuitry and function of the inferior olive. *Trends Neurosci.* **21**, 391–400 (1998).
- Ito, M. Cerebellar long-term depression: characterization, signal transduction and functional roles. *Physiol. Rev.* **81**, 1143–1195 (2001).
- Hansel, C., Linden, D.J. & D'Angelo, E. Beyond parallel fiber LTD: the diversity of synaptic and non-synaptic plasticity in the cerebellum. *Nat. Neurosci.* **4**, 467–475 (2001).
- De Zeeuw, C.I. & Yeo, C.H. Time and tide in cerebellar memory formation. *Curr. Opin. Neurobiol.* **15**, 667–674 (2005).
- Boyden, E.S., Katoh, A. & Raymond, J.L. Cerebellum-dependent learning: the role of multiple plasticity mechanisms. *Annu. Rev. Neurosci.* **27**, 581–609 (2004).
- Carey, M. & Lisberger, S. Embarrassed, but not depressed: eye opening lessons for cerebellar learning. *Neuron* **35**, 223–226 (2002).
- Ito, M. Mechanisms of motor learning in the cerebellum. *Brain Res.* **886**, 237–245 (2000).
- Raymond, J.L., Lisberger, S.G. & Mauk, M.D. The cerebellum: a neuronal learning machine? *Science* **272**, 1126–1131 (1996).
- Medina, J.F., Carey, M.R. & Lisberger, S.G. The representation of time for motor learning. *Neuron* **45**, 157–167 (2005).
- Robinson, F.R. & Fuchs, A.F. The role of the cerebellum in voluntary eye movements. *Annu. Rev. Neurosci.* **24**, 981–1004 (2001).
- Lisberger, S.G., Morris, E.J. & Tychsen, L. Visual motion processing and sensory-motor integration for smooth pursuit eye movements. *Annu. Rev. Neurosci.* **10**, 97–129 (1987).
- Stone, L.S. & Lisberger, S.G. Visual responses of Purkinje cells in the cerebellar flocculus during smooth-pursuit eye movements in monkeys. I. Simple spikes. *J. Neurophysiol.* **63**, 1241–1261 (1990).
- Medina, J.F. & Lisberger, S.G. Variation, signal, and noise in cerebellar sensory-motor processing for smooth-pursuit eye movements. *J. Neurosci.* **27**, 6832–6842 (2007).
- Stone, L.S. & Lisberger, S.G. Detection of tracking errors by visual climbing fiber inputs to monkey cerebellar flocculus during pursuit eye movements. *Neurosci. Lett.* **72**, 163–168 (1986).
- Stone, L.S. & Lisberger, S.G. Visual responses of Purkinje cells in the cerebellar flocculus during smooth-pursuit eye movements in monkeys. II. Complex spikes. *J. Neurophysiol.* **63**, 1262–1275 (1990).
- Sato, Y., Miura, A., Fushiki, H. & Kawasaki, T. Short-term modulation of cerebellar Purkinje cell activity after spontaneous climbing fiber input. *J. Neurophysiol.* **68**, 2051–2062 (1992).
- Raymond, J.L. & Lisberger, S.G. Neural learning rules for the vestibulo-ocular reflex. *J. Neurosci.* **18**, 9112–9129 (1998).
- Doi, T., Kuroda, S., Michikawa, T. & Kawato, M. Inositol 1,4,5-trisphosphate-dependent Ca^{2+} threshold dynamics detect spike timing in cerebellar Purkinje cells. *J. Neurosci.* **25**, 950–961 (2005).
- Wang, S.S., Denk, W. & Hausser, M. Coincidence detection in single dendritic spines mediated by calcium release. *Nat. Neurosci.* **3**, 1266–1273 (2000).
- Houk, J.C. & Alford, S. Computational significance of the cellular mechanisms for synaptic plasticity in Purkinje cells. *Behav. Brain Sci.* **19**, 457–461 (1996).
- Jorntell, H. & Hansel, C. Synaptic memories upside down: bidirectional plasticity at cerebellar parallel fiber–Purkinje cell synapses. *Neuron* **52**, 227–238 (2006).
- Mittmann, W. & Hausser, M. Linking synaptic plasticity and spike output at excitatory and inhibitory synapses onto cerebellar Purkinje cells. *J. Neurosci.* **27**, 5559–5570 (2007).
- Kano, M., Rexhausen, U., Dreessen, J. & Konnerth, A. Synaptic excitation produces a long-lasting rebound potentiation of inhibitory synaptic signals in cerebellar Purkinje cells. *Nature* **356**, 601–604 (1992).
- Lisberger, S.G. Neural basis for motor learning in the vestibuloocular reflex of primates. III. Computational and behavioral analysis of the sites of learning. *J. Neurophysiol.* **72**, 974–998 (1994).
- Blazquez, P.M., Hirata, Y. & Highstein, S.M. Chronic changes in inputs to dorsal Y neurons accompany VOR motor learning. *J. Neurophysiol.* **95**, 1812–1825 (2006).
- Medina, J.F., Noes, W.L., Ohya, T. & Mauk, M.D. Mechanisms of cerebellar learning suggested by eyelid conditioning. *Curr. Opin. Neurobiol.* **10**, 717–724 (2000).
- Christian, K.M. & Thompson, R.F. Neural substrates of eye blink conditioning: acquisition and retention. *Learn. Mem.* **10**, 427–455 (2003).
- Krauzlis, R.J. Population coding of movement dynamics by cerebellar Purkinje cells. *Neuroreport* **11**, 1045–1050 (2000).
- Krauzlis, R.J. & Lisberger, S.G. Simple spike responses of gaze velocity Purkinje cells in the floccular lobe of the monkey during the onset and offset of pursuit eye movements. *J. Neurophysiol.* **72**, 2045–2050 (1994).
- Shidara, M., Kawano, K., Gomi, H. & Kawato, M. Inverse-dynamics model eye movement control by Purkinje cells in the cerebellum. *Nature* **365**, 50–52 (1993).
- Kahlon, M. & Lisberger, S.G. Changes in the responses of Purkinje cells in the floccular complex of monkeys after motor learning in smooth pursuit eye movements. *J. Neurophysiol.* **84**, 2945–2960 (2000).
- Lisberger, S.G. & Fuchs, A.F. Role of primate flocculus during rapid behavioral modification of vestibuloocular reflex. II. Mossy fiber firing patterns during horizontal head rotation and eye movement. *J. Neurophysiol.* **41**, 764–777 (1978).
- Miles, F.A., Fuller, J.H., Braitman, D.J. & Dow, B.M. Long-term adaptive changes in primate vestibuloocular reflex. III. Electrophysiological observations in flocculus of normal monkeys. *J. Neurophysiol.* **43**, 1437–1476 (1980).
- Highstein, S.M. Synaptic linkage in the vestibulo-ocular and cerebello-vestibular pathways to the Vth nucleus in the rabbit. *Exp. Brain Res.* **17**, 301–314 (1973).
- Ron, S. & Robinson, D.A. Eye movements evoked by cerebellar stimulation in the alert monkey. *J. Neurophysiol.* **36**, 1004–1022 (1973).
- Wolpert, D. & Flanagan, J. in *Handbook of Brain Theory and Neural Networks* (Arbib, M.A.) 1020–1023 (MIT Press, Cambridge, Massachusetts, 2002).
- Kawato, M., Furukawa, K. & Suzuki, R. A hierarchical neural-network model for control and learning of voluntary movement. *Biol. Cybern.* **57**, 169–185 (1987).
- Pouget, A. & Snyder, L.H. Computational approaches to sensorimotor transformations. *Nat. Neurosci.* **3** Suppl: 1192–1198 (2000).
- Jordan, M. & Rumelhart, D. Forward models: supervised learning with a distal teacher. *Cogn. Sci.* **16**, 307–354 (1992).
- Fine, M.S. & Thoroughman, K.A. Trial-by-trial transformation of error into sensorimotor adaptation changes with environmental dynamics. *J. Neurophysiol.* **98**, 1392–1404 (2007).
- Cheng, S. & Sabes, P.N. Calibration of visually guided reaching is driven by error-corrective learning and internal dynamics. *J. Neurophysiol.* **97**, 3057–3069 (2007).
- Jirenhed, D.A., Bengtsson, F. & Hesselro, G. Acquisition, extinction and reacquisition of a cerebellar cortical memory trace. *J. Neurosci.* **27**, 2493–2502 (2007).
- Ramachandran, R. & Lisberger, S.G. Transformation of vestibular signals into motor commands in the vestibuloocular reflex pathways of monkeys. *J. Neurophysiol.* **96**, 1061–1074 (2006).
- Lisberger, S.G. & Pavelko, T.A. Vestibular signals carried by pathways subserving plasticity of the vestibulo-ocular reflex in monkeys. *J. Neurosci.* **6**, 346–354 (1986).
- Gilbert, P.F. & Thach, T.J. Purkinje cell activity during motor learning. *Brain Res.* **128**, 309–328 (1977).
- Ojakangas, C.L. & Ebner, T.J. Purkinje cell complex and simple spike changes during a voluntary arm movement learning task in the monkey. *J. Neurophysiol.* **68**, 2222–2236 (1992).

Corrigendum: Links from complex spikes to local plasticity and motor learning in the cerebellum of awake-behaving monkeys

Javier F Medina and Stephen G Lisberger

Nat. Neurosci. **11**, 1185–1192 (2008); published online 21 September 2008; corrected after print 15 January 2009

In the version of this article initially published, two citations were inadvertently omitted. To correct this, the following two sentences were added to the second paragraph of the introduction, following the sixth sentence. “One line of work has supported the theory by demonstrating that arm movement errors evoke complex spikes^{51,52} and that subsequent learned changes in motor behavior are associated with suitable changes in simple spike responses⁵¹. This work demonstrates a strong correlation, but stops short of showing cause-and-effect links between individual complex spikes, changes in simple spikes and behavioral learning.” Two references were also added to the reference list as follows:

51. Gilbert, P.F. & Thach W.T. Purkinje cell activity during motor learning. *Brain Res.* **128**, 309–328 (1977).

52. Ojakangas C.L & Ebner T.J. Purkinje cell complex and simple spike changes during a voluntary arm movement learning task in the monkey. *J. Neurophysiol.* **68**, 2222–2236 (1992).

The error has been corrected in the HTML and PDF versions of the article.

Corrigendum: Links from complex spikes to local plasticity and motor learning in the cerebellum of awake-behaving monkeys

Javier F Medina & Stephen G Lisberger

Nat. Neurosci. 11, 1185–1192 (2008); published online 21 September 2008; corrected after print 15 January and 30 April 2009

The second sentence of the abstract should read “Many elements of this hypothesis have been supported by prior experiments, and correlations have been found between complex spikes, simple-spike plasticity and behavior during the learning process.” This error has been corrected in the HTML and PDF versions of the article.