

NEWTON POLYGONS ARISING FROM SPECIAL FAMILIES OF CYCLIC COVERS OF THE PROJECTIVE LINE

WANLIN LI, ELENA MANTOVAN, RACHEL PRIES, AND YUNQING TANG

ABSTRACT. By a result of Moonen, there are exactly 20 positive-dimensional families of cyclic covers of the projective line for which the Torelli image is open and dense in the associated Shimura variety. For each of these, we compute the Newton polygons, and the μ -ordinary Ekedahl–Oort type, occurring in the characteristic p reduction of the Shimura variety. We prove that all but a few of the Newton polygons appear on the open Torelli locus. As an application, we produce multiple new examples of Newton polygons and Ekedahl–Oort types of Jacobians of smooth curves in characteristic p . Under certain congruence conditions on p , these include: the supersingular Newton polygon for genus 5, 6, 7; fourteen new non-supersingular Newton polygons for genus 5 – 7; eleven new Ekedahl–Oort types for genus 4 – 7 and, for all $g \geq 6$, the Newton polygon with p -rank $g - 6$ with slopes $1/6$ and $5/6$.

MSC10: primary 11G18, 11G20, 11M38, 14G10, 14G35; secondary 11G10, 14H10, 14H30, 14H40, 14K22

Keywords: curve, cyclic cover, Jacobian, abelian variety, Shimura variety, PEL-type, moduli space, reduction, p -rank, supersingular, Newton polygon, p -divisible group, Kottwitz method, Dieudonné module, Ekedahl–Oort type.

1. INTRODUCTION

A fundamental problem in arithmetic geometry is to determine which abelian varieties arise as Jacobians of (smooth) curves. This question is equivalent to studying (the interior of) the Torelli locus in certain Siegel varieties. For $g = 1, 2, 3$, the Torelli locus is open and dense in the moduli space \mathcal{A}_g of principally polarized abelian varieties of dimension g . For any family T of cyclic covers of the projective line, Deligne and Mostow construct the smallest PEL-type Shimura variety containing the image of T under the Torelli morphism [6]. In [25], Moonen shows that there are precisely twenty positive-dimensional families of cyclic covers of \mathbb{P}^1 for which the image of T under the Torelli morphism is open and dense in the associated Shimura variety; these families are called special.

In positive characteristic p , the p -rank, Newton polygon, and Ekedahl–Oort type are discrete invariants of an abelian variety. It is a natural question to ask which of these invariants can be realized by the Jacobian of a smooth curve. As each discrete invariant yields a stratification of the reduction modulo p of the Siegel variety, this question is equivalent to understanding which strata intersect the interior of the Torelli locus.

Ultimately, the goal is to understand the geometry of the induced stratifications of the Torelli locus (e.g., the connected components of each stratum and their closures), in the same way that the geometry of these stratifications is understood for Siegel varieties. For example, for any prime p , genus g and integer f such that $0 \leq f \leq g$, Faber and van der Geer prove in [9] that the p -rank f stratum is non-empty and has the expected codimension $g - f$ in the moduli space \mathcal{M}_g of curves of genus g . See also [1], [2], and [12].

Beyond genus 3, much less is known about the Newton polygon, more precisely the Newton polygon of the characteristic polynomial of Frobenius. In [26, Expectation 8.5.4], for $g \geq 9$, using a dimension count, Oort observed that the interior of the Torelli locus is unlikely to intersect the supersingular locus or other Newton polygon stratum of high codimension in \mathcal{A}_g . This suggests that it is unlikely for all Newton polygons to occur for Jacobians of smooth curves for each prime p .

In our earlier work [22], we studied Newton polygons of cyclic covers of the projective line branched at 3 points. In this case, the Jacobian is an abelian variety with complex multiplication and its Newton polygon can be computed using the Shimura–Taniyama theorem. We used this to find several new Newton polygons having p -rank 0 which occur for Jacobians of smooth curves.

The Newton polygon stratification of PEL-type Shimura varieties is well understood by the work of Viehmann and Wedhorn [35], based on earlier work of Kottwitz, Rapoport, and Richartz [16], [17], [19], [29]. In this paper, we demonstrate this theory by determining all the Newton polygons of Jacobians that occur for the 20 special families of cyclic covers from Moonen’s paper [25]. Furthermore, using [24], we compute the μ -ordinary mod p Dieudonné modules (e.g., Ekedahl–Oort types) for the 20 special families.

We then investigate which Newton polygons and Dieudonné modules on these lists arise for a smooth curve in the family. We conclude affirmatively in three distinct cases:

- (i) for the μ -ordinary Newton polygon and Dieudonné module, Proposition 5.1;
- (ii) for any PEL-indecomposable Newton polygon, Proposition 5.6; and
- (iii) for the Newton polygon of the closed stratum (called the basic Newton polygon) if the family has dimension 1 and if p is sufficiently large, Proposition 5.8.

For (iii), we refer to Theorem 8.1 in the Appendix, where we prove, under some restrictions on the prime p , that the number of geometrically irreducible components of the basic locus at p of certain unitary Shimura variety grows to infinity with p . More precisely, Theorem 8.1 holds for unitary Shimura varieties of signature $(1, n - 1)$ at one real place and definite signature at all other real places. In particular, our results hold for the simple Shimura varieties considered by Harris–Taylor in [14]. Earlier instances of such a statement are due to Vollaard [36], and Vollaard–Wedhorn [37] who prove a similar result under the further assumption that the implicit CM field defining the unitary group is quadratic-imaginary, and to Liu–Tian–Xiao–Zhang–Zhu [23] at primes which are totally split in the CM field. Here, we only exclude the case of primes which are inert in the quadratic imaginary extension for n even. We regard Theorem 8.1 to be of independent interest.

We calculate that these three criteria together with the de Jong–Oort purity result imply that all but a few of the Newton polygons for Moonen’s special families occur for Jacobians of smooth curves. More precisely, we prove the following result.

Theorem 1.1. *(See Theorem 5.11) Let (m, N, a) denote the monodromy datum for one of Moonen’s special families from [25, Table 1]. Assume $p \nmid m$. Let $v \in v(B(\mu_m, \mathfrak{f}))$ be a Newton polygon occurring on the closure in \mathcal{A}_g of the image of the family under the Torelli morphism. Then v occurs as the Newton polygon of the Jacobian of a smooth curve in the family unless either:*

- (1) *the dimension of the family is one, v is supersingular, and p is not sufficiently large; or*
- (2) *the dimension of the family is at least two and v is supersingular.*

In cases (1) and (2), we also expect that there exists a smooth curve in the family which is supersingular; (we prove this for large p in [21, Theorem 6.1]).

We deduce the following applications.

First, we verify the existence of supersingular curves of genus $5 - 7$ for primes satisfying certain congruence conditions. See [22, Theorem 1.1] for a complementary result.

Theorem 1.2 (See Theorem 7.1). *There exists a smooth supersingular curve of genus g defined over $\overline{\mathbb{F}}_p$ for all sufficiently large primes satisfying the given congruence condition: $g = 5$ when $p \equiv 7 \pmod{8}$; $g = 6$ when $p \equiv 2, 3, 4 \pmod{5}$; and $g = 7$ when $p \equiv 2 \pmod{3}$.*

The second application is in Theorems 7.3 and 7.4: under certain congruence conditions on p , we verify that 14 new non-supersingular Newton polygons and 11 new Dieudonné modules occur for Jacobians of smooth curves of genus $4 - 7$.

Every abelian variety is isogenous to a factor of a Jacobian. Thus every $\lambda \in [0, 1] \cap \mathbb{Q}$ is known to occur as a slope for the Newton polygon of the Jacobian of a smooth curve. Usually, however, there is no control over the other slopes. In the third application, for $p \equiv 2, 4 \pmod{7}$ and $g \geq 6$, we show that the slopes $1/6$ and $5/6$ occur for a smooth curve of genus g defined over $\overline{\mathbb{F}}_p$ with the other slopes of the Newton polygon all 0 and 1, Theorem 7.5.

As a final application, we consider one non-special family of curves in Section 7.2. Consider a prime $m \equiv 3 \pmod{4}$ (with $m \neq 3$) and a prime p which is a quadratic non-residue modulo m . In Theorem 7.8, we find smooth curves of genus $g_m = (m - 5)(m - 1)/4$ defined over $\overline{\mathbb{F}}_p$ whose Newton polygon only has slopes 0, $1/2$, and 1 and such that the multiplicity of the slope $1/2$ is at least $2\sqrt{g_m}$.

In future work, we solve questions about Newton polygons for curves of arbitrarily large genera using a new induction argument for Newton polygons of cyclic covers of \mathbb{P}^1 . We use the Newton polygons from this paper as base cases in that induction process.

Organization of the paper.

Section 2 recalls basic definitions and facts about group algebras, families of cyclic covers of \mathbb{P}^1 , Newton polygons, and mod p Dieudonné modules.

Section 3 contains information about Shimura varieties and positive-dimensional special families of cyclic covers of the projective line.

In Section 4, we study the Newton polygon stratification of PEL-type Shimura varieties. We demonstrate the general theory by computing the Newton polygons for several prototypical examples from [25] in Section 4.4. In Section 4.5, we determine the μ -ordinary Dieudonné modules (Ekedahl–Oort types) for the special families.

Section 5 discusses which Newton polygon strata intersect the smooth Torelli locus.

Section 6 contains tables of data, including the Newton polygons and the μ -ordinary Dieudonné modules for each of the twenty special families of cyclic covers from [25].

Section 7 contains the proofs of the main theorems.

In the appendix, we compute a lower bound for the number of irreducible components of the basic locus of simple Shimura varieties.

Acknowledgements.

This project began at the *Women in Numbers 4* workshop at the Banff International Research Station. Pries was partially supported by NSF grant DMS-15-02227. We thank

Liang Xiao, Xinwen Zhu, and Rong Zhou for discussions about the appendix and thank Liang Xiao for the detailed suggestions on the writing of the appendix. We would like to thank the referee for many helpful comments.

2. NOTATION AND BACKGROUND

2.1. The group algebra $\mathbb{Q}[\mu_m]$.

For an integer $m \geq 2$, let $\mu_m := \mu_m(\mathbb{C})$ denote the group of m -th roots of unity in \mathbb{C} . For each positive integer d , we fix a primitive d -th root of unity $\zeta_d = e^{\frac{2\pi i}{d}} \in \mathbb{C}$. Let $K_d = \mathbb{Q}(\zeta_d)$ be the d -th cyclotomic field over \mathbb{Q} of degree $\phi(d)$.

Let $\mathbb{Q}[\mu_m]$ denote the group algebra of μ_m over \mathbb{Q} . Then $\mathbb{Q}[\mu_m] = \prod_{0 < d | m} K_d$. The involution $*$ on $\mathbb{Q}[\mu_m]$ induced by the inverse map on μ_m preserves each cyclotomic factor K_d , and for each $d | m$, the restriction of $*$ to K_d agrees with complex conjugation.

Set $\mathcal{T} := \text{Hom}_{\mathbb{Q}}(\mathbb{Q}[\mu_m], \mathbb{C})$. Each $(\mathbb{Q}[\mu_m] \otimes_{\mathbb{Q}} \mathbb{C})$ -module W has an eigenspace decomposition $W = \bigoplus_{\tau \in \mathcal{T}} W_{\tau}$, where W_{τ} is the subspace of W on which $a \otimes 1 \in \mathbb{Q}[\mu_m] \otimes_{\mathbb{Q}} \mathbb{C}$ acts as $\tau(a)$. We fix an identification $\mathcal{T} = \mathbb{Z}/m\mathbb{Z}$ by defining, for all $n \in \mathbb{Z}/m\mathbb{Z}$,

$$\tau_n(\zeta) := \zeta^n, \text{ for all } \zeta \in \mu_m.$$

For all $n \in \mathbb{Z}/m\mathbb{Z}$, and $a \in \mathbb{Q}[\mu_m]$, $\tau_{-n}(a) = \tau_n(a^*)$. We write $\tau_n^* := \tau_{-n}$.

For $\tau = \tau_n \in \mathcal{T}$, the *order of τ* is the order of n in $\mathbb{Z}/m\mathbb{Z}$. The homomorphism $\tau : \mathbb{Q}[\mu_m] \rightarrow \mathbb{C}$ factors via K_d if and only if τ has order d .

For each rational prime p , we fix an algebraic closure $\overline{\mathbb{Q}}_p^{\text{alg}}$ of \mathbb{Q}_p , and an identification $\iota : \mathbb{C} \simeq \mathbb{C}_p$, where \mathbb{C}_p denotes the p -adic completion of $\overline{\mathbb{Q}}_p^{\text{alg}}$. Let $\overline{\mathbb{Q}}_p^{\text{un}}$ denote the maximal unramified extension of \mathbb{Q}_p in $\overline{\mathbb{Q}}_p^{\text{alg}}$, and by σ the Frobenius of $\overline{\mathbb{Q}}_p^{\text{un}}$.

If $p \nmid m$, then $\mathbb{Q}[\mu_m]$ is unramified at p . Via ι , we identify $\mathcal{T} = \text{Hom}_{\mathbb{Q}}(\mathbb{Q}[\mu_m], \overline{\mathbb{Q}}_p^{\text{un}})$. Hence, there is a natural action of σ on \mathcal{T} , defined by $\tau \mapsto \tau^{\sigma} := \sigma \circ \tau$. Note that $\tau_n^{\sigma} = \tau_{pn}$ for all $n \in \mathbb{Z}/m\mathbb{Z}$. For each σ -orbit $\mathfrak{o} \subseteq \mathcal{T}$, the order of τ is the same for all $\tau \in \mathfrak{o}$. We write $d_{\mathfrak{o}}$ for the order of $\tau \in \mathfrak{o}$.

Let \mathfrak{D} be the set of σ -orbits \mathfrak{o} in \mathcal{T} . For each $\tau \in \mathcal{T}$, let \mathfrak{o}_{τ} denote its σ -orbit. Each $\tau : \mathbb{Q}[\mu_m] \rightarrow \overline{\mathbb{Q}}_p^{\text{un}}$ determines a prime \mathfrak{p} of $\mathbb{Q}[\mu_m]$ above p , depending only on the orbit \mathfrak{o}_{τ} . Hence, the set \mathfrak{D} is in bijection with the set of primes \mathfrak{p} of $\mathbb{Q}[\mu_m]$ above p . For an orbit $\mathfrak{o} \in \mathcal{T}$, we write $\mathfrak{p}_{\mathfrak{o}}$ for the associated prime above p and $K_{d_{\mathfrak{o}}, \mathfrak{p}_{\mathfrak{o}}}$ for the completion of $K_{d_{\mathfrak{o}}}$ along the prime $\mathfrak{p}_{\mathfrak{o}}$.

2.2. Families of cyclic covers of the projective line.

We follow [25, §§2,3]. Fix integers $m \geq 2, N \geq 3$ and an N -tuple of positive integers $a = (a(1), \dots, a(N))$. Then a is an *inertia type* for m and (m, N, a) is a *monodromy datum* if

- (1) $a(i) \not\equiv 0 \pmod{m}$, for all $i = 1, \dots, N$,
- (2) $\gcd(m, a(1), \dots, a(N)) = 1$, and
- (3) $\sum_i a(i) \equiv 0 \pmod{m}$.

Fix a monodromy datum (m, N, a) . Working over $\mathbb{Z}[1/m]$, let $U \subset (\mathbb{A}^1)^N$ be the complement of the weak diagonal, which consists of points where any two of the coordinates

are the same. Consider the smooth projective (relative) curve C over U whose fiber at each point $t = (t(1), \dots, t(N)) \in U$ has affine model

$$(2.1) \quad y^m = \prod_{i=1}^N (x - t(i))^{a(i)}.$$

The function x on C yields a map $C \rightarrow \mathbb{P}_U^1$ and there is a μ_m -action on C over U given by $\zeta \cdot (x, y) = (x, \zeta \cdot y)$ for all $\zeta \in \mu_m$. Thus $C \rightarrow \mathbb{P}_U^1$ is a μ_m -Galois cover.

For a point $t \in U$, the cover $C_t \rightarrow \mathbb{P}^1$ is a μ_m -Galois cover, branched at N points $t(1), \dots, t(N)$ in \mathbb{P}^1 , and with local monodromy $a(i)$ at $t(i)$. By the hypotheses on the monodromy datum, the fibers of $C \rightarrow U$ are geometrically irreducible curves of genus g , where

$$(2.2) \quad g = g(m, N, a) = 1 + \frac{(N-2)m - \sum_{i=1}^N \gcd(a(i), m)}{2}.$$

Let \mathcal{M}_g denote the moduli space of smooth projective curves of genus g and let \mathcal{A}_g denote the Siegel moduli space of principally polarized abelian varieties of dimension g over \mathbb{Z} .¹ The composition of the Torelli map with the morphism $U \rightarrow \mathcal{M}_g$ defined by the curve $C \rightarrow U$ yields a morphism over $\mathbb{Z}[1/m]$ denoted by

$$j = j(m, N, a) : U \rightarrow \mathcal{M}_g \rightarrow \mathcal{A}_g.$$

Definition 2.1. If (m, N, a) is a monodromy datum, let $Z^0(m, N, a)$ be the image of $j(m, N, a)$ in \mathcal{A}_g and let $Z(m, N, a)$ be its closure in \mathcal{A}_g .

Remark 2.2. By definition, $Z(m, N, a)$ is a closed, reduced substack of \mathcal{A}_g . It is also irreducible [10, Corollary 7.5], [38, Corollary 4.2.3].

The substack $Z(m, N, a)$ depends uniquely on the equivalence class of the monodromy datum (m, N, a) , where (m, N, a) and (m', N', a') are equivalent if $m = m'$, $N = N'$, and the images of a, a' in $(\mathbb{Z}/m\mathbb{Z})^N$ are in the same orbit under $(\mathbb{Z}/m\mathbb{Z})^* \times \text{Sym}_N$.

Fix a point $t \in U(\mathbb{C})$ and let V be the first Betti cohomology group $H^1(C_t(\mathbb{C}), \mathbb{Q})$. Then V has a Hodge structure of type $(1, 0) + (0, 1)$ with the $(1, 0)$ piece given by $H^0(C_t(\mathbb{C}), \Omega_{C_t}^1)$ via the Betti–de Rham comparison. Let V^+ (resp. V^-) denote the $(1, 0)$ (resp. $(0, 1)$) piece.

Then V, V^+ , and V^- are $\mathbb{Q}[\mu_m]$ -modules and there are decompositions

$$V \otimes_{\mathbb{Q}} \mathbb{C} = \bigoplus_{\tau \in \mathcal{T}} V_{\tau}, \quad V^+ = \bigoplus_{\tau \in \mathcal{T}} V_{\tau}^+, \quad V^- = \bigoplus_{\tau \in \mathcal{T}} V_{\tau}^-.$$

Set:

$$(2.3) \quad g(\tau) := \dim_{\mathbb{C}} V_{\tau}, \quad f(\tau) := \dim_{\mathbb{C}} V_{\tau}^+.$$

For any $x \in \mathbb{Q}$, let $\langle x \rangle$ denote the fractional part of x . By [25, Lemma 2.7, §3.2] (or [6]),

$$(2.4) \quad f(\tau_n) = \begin{cases} -1 + \sum_{i=1}^N \langle \frac{-na(i)}{m} \rangle & \text{if } n \not\equiv 0 \pmod{m} \\ 0 & \text{if } n \equiv 0 \pmod{m}. \end{cases}$$

The dimension $f(\tau_n)$ is independent of the choice of $t \in U$. The *signature type* of the monodromy datum (m, N, a) is

$$\mathfrak{f} = (f(\tau_1), \dots, f(\tau_{m-1})).$$

¹When we talk about universal families on the stacks, it means that we pass to a suitable level structure for such universal objects to exist.

Remark 2.3. For all $\tau \in \mathcal{T}$, $\dim_{\mathbb{C}} V_{\tau^*}^+ = \dim_{\mathbb{C}} V_{\tau}^-$ and thus $g(\tau) = f(\tau) + f(\tau^*)$. The value $g(\tau)$ is constant on the σ -orbit of τ , so we sometimes write $g(\mathfrak{o}) = g(\tau)$ for $\tau \in \mathfrak{o}$.

2.3. Newton polygons.

Let X denote a g -dimensional abelian scheme over an algebraically closed field \mathbb{F} of positive characteristic p (e.g., $X = \mathcal{X}_x$, where \mathcal{X} is the universal abelian scheme on \mathcal{A}_g , and $x \in \mathcal{A}_g(\mathbb{F})$).

If \mathbb{F} is an algebraic closure of \mathbb{F}_p , the finite field of p elements, then there exists a finite subfield $\mathbb{F}_0 \subset \mathbb{F}$ such that X is isomorphic to the base change to \mathbb{F} of an abelian scheme X_0 over \mathbb{F}_0 . Let $W(\mathbb{F}_0)$ denote the Witt vector ring of \mathbb{F}_0 . Consider the action of Frobenius φ on the crystalline cohomology group $H_{\text{cris}}^1(X_0/W(\mathbb{F}_0))$. There exists an integer n such that φ^n , the composite of n Frobenius actions, is a linear map on $H_{\text{cris}}^1(X_0/W(\mathbb{F}_0))$. The *Newton polygon* $\nu(X)$ of X is defined as the multi-set of rational numbers λ , called the *slopes*, such that $n\lambda$ are the valuations at p of the eigenvalues of Frobenius for this action. Note that the Newton polygon is independent of the choice of X_0 , \mathbb{F}_0 , and n . For a more general \mathbb{F} , one can use the Dieudonné–Manin classification to define the Newton polygon.

The p -rank of X is the multiplicity of the slope 0 in $\nu(X)$; it equals $\dim_{\mathbb{F}_p} \text{Hom}(\mu_p, X)$.

If \mathcal{A} is an abelian variety or p -divisible group defined over a local field of mixed characteristic $(0, p)$, we write $\nu(\mathcal{A})$ for the Newton polygon of its special fiber.

For $\lambda \in \mathbb{Q} \cap [0, 1]$, the multiplicity m_λ is the multiplicity of λ in the multi-set; if $c, d \in \mathbb{N}$ are relatively prime integers such that $\lambda = c/(c + d)$, then $(c + d)$ divides m_λ . The Newton polygon is *symmetric* if $m_\lambda = m_{1-\lambda}$ for every $\lambda \in \mathbb{Q} \cap [0, 1]$. The Newton polygon is typically drawn as a lower convex polygon, with slopes equal to the values of λ occurring with multiplicity m_λ . The Newton polygon of a g -dimensional abelian variety X is symmetric and, when drawn as a polygon, it has endpoints $(0, 0)$ and $(2g, g)$ and integral break points.

Let *ord* denote the Newton polygon $\{0, 1\}$ and *ss* denote the Newton polygon $\{1/2, 1/2\}$. Thus an ordinary (resp. supersingular) abelian variety of dimension g has Newton polygon *ord* ^{g} (resp. *ss* ^{g}). For $s, t \in \mathbb{N}$, with $s \leq t/2$ and $\gcd(s, t) = 1$, let $(s/t, (t - s)/t)$ denote the Newton polygon with slopes s/t and $(t - s)/t$, each with multiplicity t .

We denote the union of multi-sets by \oplus , and for any multi-set ν , and $n \in \mathbb{N}$, we write ν^n for $\nu \oplus \cdots \oplus \nu$, n -times. For convex polygons, we write $\nu_1 \geq \nu_2$ if ν_1, ν_2 share the same endpoints and ν_1 lies below ν_2 . Under this convention, the ordinary Newton polygon is maximal and the supersingular Newton polygon is minimal.

2.4. Dieudonné modules modulo p and Ekedahl–Oort types.

The p -torsion $X[p]$ of X is a symmetric BT_1 -group scheme (of rank $2g$) annihilated by p . The a -number of X is $\dim_{\mathbb{F}}(\alpha_p, X[p])$, where α_p is the kernel of Frobenius on \mathbb{G}_a .

Let \mathbb{E} denote the non-commutative ring over $\overline{\mathbb{F}}_p$, generated by a σ -linear F and σ^{-1} -linear V with $FV = VF = 0$. The mod p Dieudonné module of X is an \mathbb{E} -module of finite dimension $(2g)$. Over \mathbb{F} , there is an equivalence of categories between finite commutative group schemes of rank $2g$ annihilated by p and \mathbb{E} -modules of dimension $2g$. These can equivalently be classified by Ekedahl–Oort types.

Let $\mathbb{E}(w)$ denote the left ideal of \mathbb{E} generated by w . The mod p Dieudonné module for an ordinary elliptic curve is isomorphic to $L := \mathbb{E}/\mathbb{E}(F, V-1) \oplus \mathbb{E}/\mathbb{E}(V, F-1)$; for a supersingular elliptic curve, it is isomorphic to $N_{1,1} := \mathbb{E}/\mathbb{E}(F-V)$.

3. SHIMURA VARIETIES

Let (m, N, a) be a monodromy datum with $N \geq 4$, and \mathfrak{f} the associated signature type given by (2.4). In [6] Deligne and Mostow construct the smallest PEL-type Shimura variety containing $Z(m, N, a)$, which we will denote by $\text{Sh}(\mu_m, \mathfrak{f})$. In this section, we recall the basic setting for PEL-type Shimura varieties, and the construction of [6]. We follow [25].

3.1. Shimura datum for the moduli space of abelian varieties.

Let $V = \mathbb{Q}^{2g}$, and let $\Psi : V \times V \rightarrow \mathbb{Q}$ denote the standard symplectic form.² Let $G := \text{GSp}(V, \Psi)$ denote the group of symplectic similitudes over \mathbb{Q} . Let \mathfrak{h} denote the space of homomorphisms $h : \mathbb{S} = \text{Res}_{\mathbb{C}/\mathbb{R}} \mathbb{G}_m \rightarrow \mathbb{G}_{\mathbb{R}}$ which define a Hodge structure of type $(-1, 0) + (0, -1)$ on $V_{\mathbb{Z}}$ such that $\pm(2\pi i)\Psi$ is a polarization on V . The pair (G, \mathfrak{h}) is the Shimura datum for \mathcal{A}_g .

Let $H \subset G$ be an algebraic subgroup over \mathbb{Q} such that the subspace

$$\mathfrak{h}_H := \{h \in \mathfrak{h} \mid h \text{ factors through } H_{\mathbb{R}}\}$$

is non-empty. Then $H(\mathbb{R})$ acts on \mathfrak{h}_H by conjugation, and for each $H(\mathbb{R})$ -orbit $Y_H \subset \mathfrak{h}_H$, the Shimura datum (H, Y_H) defines an algebraic substack $\text{Sh}(H, Y_H)$ of \mathcal{A}_g . In the following, for $h \in Y_H$, we sometimes write (H, h) for the Shimura datum (H, Y_H) . For convenience, we also write $\text{Sh}(H, \mathfrak{h}_H)$ for the finite union of the Shimura stacks $\text{Sh}(H, Y_H)$, as Y_H varies among the $H(\mathbb{R})$ -orbits in \mathfrak{h}_H .

3.2. Shimura data of PEL-type.

Now we focus on Shimura data of PEL-type. Let B be a semisimple \mathbb{Q} -algebra, together with an involution $*$. Suppose there is an action of B on V such that $\Psi(bv, w) = \Psi(v, b^*w)$, for all $b \in B$ and all $v, w \in V$. Let

$$H_B := \text{GL}_B(V) \cap \text{GSp}(V, \Psi).$$

We assume that $\mathfrak{h}_{H_B} \neq \emptyset$.

For each $H_B(\mathbb{R})$ -orbit $Y_B := Y_{H_B} \subset \mathfrak{h}_{H_B}$, the associated Shimura stack $\text{Sh}(H_B, Y_B)$ arise as moduli spaces of polarized abelian varieties endowed with a B -action, and are called of PEL-type. In the following, we also write $\text{Sh}(B) := \text{Sh}(H_B, \mathfrak{h}_{H_B})$.

Each homomorphism $h \in Y_B$ defines a decomposition of $B_{\mathbb{C}}$ -modules

$$V_{\mathbb{C}} = V^+ \oplus V^-$$

where V^+ (respectively, V^-) is the subspace of $V_{\mathbb{C}}$ on which $h(z)$ acts by z (respectively, by \bar{z}). The isomorphism class of the $B_{\mathbb{C}}$ -module V^+ depends only on Y_B . Moreover, Y_B is

²In Section 2.2, we use V to denote $H^1(C_x, \mathbb{Q})$ for $x \in U(\mathbb{C})$. Here, by the convention of Deligne, V is identified with $H_1(\mathcal{X}_x, \mathbb{Q})$ for any $x \in \mathcal{A}_g(\mathbb{C})$. The symplectic form will be identified with the one induced by the polarization on \mathcal{X}_x . This symplectic form, along with Poincaré duality, induces a natural isomorphism $H^1(C_x, \mathbb{Q})(1) \cong H_1(C_x, \mathbb{Q})$. Hence the only difference between them is given by the Tate twist and we use the same notation V for both.

determined by the isomorphism class of V^+ as a $B_{\mathbb{C}}$ -submodule of $V_{\mathbb{C}}$. In the following, we prescribe Y_B in terms of the $B_{\mathbb{C}}$ -module V^+ . By construction, $\dim_{\mathbb{C}} V^+ = g$.

3.3. Shimura subvariety attached to a monodromy datum.

We consider cyclic covers of the projective line branched at more than three points; fix a monodromy datum (m, N, a) with $N \geq 4$. Take $B = \mathbb{Q}[\mu_m]$ with involution $*$ as in Section 2.1.

As in Section 2.2, let $C \rightarrow U$ denote the universal family of μ_m -covers of \mathbb{P}^1 branched at N points with inertia type a ; let $j = j(m, N, a) : U \rightarrow \mathcal{A}_g$ be the composition of the Torelli map with the morphism $U \rightarrow \mathcal{M}_g$. From Definition 2.1, recall that $Z = Z(m, N, a)$ is the closure in \mathcal{A}_g of the image of $j(m, N, a)$.

The pullback of the universal abelian scheme \mathcal{X} on \mathcal{A}_g via j is the relative Jacobian \mathcal{J} of $C \rightarrow U$. Since μ_m acts on C , there is a natural action of the group algebra $\mathbb{Z}[\mu_m]$ on \mathcal{J} . We also use \mathcal{J} to denote the pullback of \mathcal{X} to Z . The action of $\mathbb{Z}[\mu_m]$ extends naturally to \mathcal{J} over Z . Hence the substack $Z = Z(m, N, a)$ is contained in $\text{Sh}(\mathbb{Q}[\mu_m])$ for an appropriate choice of a structure of $\mathbb{Q}[\mu_m]$ -module on V . More precisely, fix $x \in Z(\mathbb{C})$, and let (\mathcal{J}_x, θ) denote the corresponding Jacobian with its principal polarization θ . Choose a symplectic similitude, meaning an isomorphism

$$\alpha : (H_1(\mathcal{J}_x, \mathbb{Q}), \psi_{\theta}) \rightarrow (V, \Psi),$$

such that the pull back of the symplectic form Ψ to $H_1(\mathcal{J}_x, \mathbb{Q})$ is a scalar multiple of ψ_{θ} , where ψ_{θ} denotes the Riemannian form on $H_1(\mathcal{J}_x, \mathbb{Q})$ corresponding to the polarization θ . Via α , the $\mathbb{Q}[\mu_m]$ -action on \mathcal{J}_x induces an action on V . This action satisfies

$$\mathfrak{h}_{\mathbb{Q}[\mu_m]} \neq \emptyset, \text{ and } \Psi(bv, w) = \Psi(v, b^*w),$$

for all $b \in \mathbb{Q}[\mu_m]$, all $v, w \in V$, and $Z \subset \text{Sh}(\mathbb{Q}[\mu_m])$.

The isomorphism class of V^+ as a $\mathbb{Q}[\mu_m] \otimes_{\mathbb{Q}} \mathbb{C}$ -module is determined by and determines the signature type $\{\mathfrak{f}(\tau) = \dim V_{\tau}^+\}_{\tau \in \mathcal{T}}$. By [6, 2.21, 2.23] (see also [25, §§3.2, 3.3, 4.5]), the $H_{\mathbb{Q}[\mu_m]}(\mathbb{R})$ -orbit $Y_{\mathbb{Q}[\mu_m]}$ in $\mathfrak{h}_{H_{\mathbb{Q}[\mu_m]}}$ such that

$$Z \subset \text{Sh}(H_{\mathbb{Q}[\mu_m]}, Y_{\mathbb{Q}[\mu_m]})$$

corresponds to the isomorphism class of V^+ with \mathfrak{f} given by (2.4). From now on, since $\text{Sh}(H_{\mathbb{Q}[\mu_m]}, Y_{\mathbb{Q}[\mu_m]})$ depends only on μ_m and \mathfrak{f} , we denote it by $\text{Sh}(\mu_m, \mathfrak{f})$.

The irreducible component of $\text{Sh}(\mu_m, \mathfrak{f})$ containing Z is the largest closed, reduced and irreducible substack S of \mathcal{A}_g containing Z such that the action of $\mathbb{Z}[\mu_m]$ on \mathcal{J} extends to the universal abelian scheme over S . To emphasis the dependence on the monodromy datum, we denote this irreducible substack by $S(m, N, a)$.

4. NEWTON POLYGONS FOR SPECIAL SHIMURA VARIETIES

In this section, we determine the Newton polygons and the μ -ordinary Dieudonné modules that occur for the special families of [25, Table 1]. We refer to [22, Section 3] for a review of the Shimura–Taniyama method for computing the Newton polygon of a cyclic cover of the projective line branched at exactly three points.

4.1. Newton polygon stratification for PEL-type Shimura varieties.

We recall some of the key results about the Newton polygon stratification for PEL-type Shimura varieties at unramified primes of good reduction from [35]. Similar results are now known for abelian-type Shimura varieties by [31] and we refer to [15] for a survey of previous work.

Let (H, h) denote a Shimura datum of PEL-type, with H connected. Assume p is an unramified prime of good reduction for (H, h) , then the Shimura variety $\mathrm{Sh}(H, h)$ has a smooth canonical integral model at p , and we denote by $\mathrm{Sh}(H, h)_{\overline{\mathbb{F}}_p}$ its special fiber (base changed to $\overline{\mathbb{F}}_p$).³ For the Shimura data introduced in Section 3.3, H is always connected and p is an unramified prime of good reduction if $p \nmid m$.

There exists a p -adic cocharacter $\mu = \mu_h$ which factors through $H(\overline{\mathbb{Q}}_p)$ and is conjugate by an element in H to the Hodge cocharacter induced by h . Given the local datum $(H_{\mathbb{Q}_p}, \mu)$, in [17, §5] and [19, §6], Kottwitz introduced a partially ordered set $B(H_{\mathbb{Q}_p}, \mu)$, and showed that $B(H_{\mathbb{Q}_p}, \mu)$ has a canonical maximal element (called the μ -ordinary element) and a minimal one (called *basic*). For the Shimura datum $(G = \mathrm{GSp}(V, \Psi), h)$ as in Section 3.1, the set $B(G_{\mathbb{Q}_p}, \mu)$ is canonically isomorphic (as a partially ordered set) to the set of all symmetric convex polygons, with endpoints $(0, 0)$ and $(2g, g)$, integral break-points, and slopes in $\mathbb{Q} \cap [0, 1]$ (see [16, §4]). Under this identification, the μ -ordinary and basic elements are respectively the ordinary and supersingular Newton polygons.

Furthermore, for a Shimura datum $(H, h) \subseteq (G, h)$, Kottwitz constructed a canonical map (the Newton map)

$$\nu : B(H_{\mathbb{Q}_p}, \mu) \rightarrow B(G_{\mathbb{Q}_p}, \mu).$$

For any $b \in B(H_{\mathbb{Q}_p}, \mu)$, consider the Newton polygon stratum

$$\mathrm{Sh}(H, h)_{\overline{\mathbb{F}}_p}(b) := \{x \in |\mathrm{Sh}(H, h)_{\overline{\mathbb{F}}_p}| \mid \nu(\mathcal{X}_x) = \nu(b)\},$$

where, as in Section 2.3, \mathcal{X} denotes the universal abelian scheme over \mathcal{A}_g . Viehmann and Wedhorn prove:

Theorem 4.1. [35, Theorem 1.6] *For any $b \in B(H_{\mathbb{Q}_p}, \mu)$ and any irreducible component S of $\mathrm{Sh}(H, h)$, there exists $x \in |S_{\overline{\mathbb{F}}_p}|$ such that $\nu(\mathcal{X}_x) = \nu(b)$.*

Definition 4.2. Let (m, N, a) be a monodromy datum with signature \mathfrak{f} . For the associated Shimura datum $(H_{\mathbb{Q}[\mu_m]}, Y_{\mathbb{Q}[\mu_m]})$ as defined in Section 3.3, we write $B(\mu_m, \mathfrak{f})$ in place of $B((H_{\mathbb{Q}[\mu_m]})_{\mathbb{Q}_p}, \mu(\mathfrak{f}))$ and denote by $\nu(B(\mu_m, \mathfrak{f}))$ its image in $B(G_{\mathbb{Q}_p}, \mu)$.

We note that the sets $B(\mu_m, \mathfrak{f})$ and $\nu(B(\mu_m, \mathfrak{f}))$ depend on p , specifically on the congruence of p modulo m .

By Theorem 4.1, $\nu(B(\mu_m, \mathfrak{f}))$ is the set of Newton polygons appearing on the irreducible component $S = S(m, N, a)$. We refer to the images under ν of the μ -ordinary and basic elements of $B(\mu_m, \mathfrak{f})$ respectively as the μ -ordinary and basic Newton polygons.

³More precisely, for p an unramified prime for the datum (H, h) , the group $H_{\mathbb{Q}_p}$ is the generic fiber of some reductive group \mathcal{H} over \mathbb{Z}_p with connected geometric fibers, and we assume that the level at p of the Shimura variety $\mathrm{Sh}(H, h)$ is the hyperspecial subgroup $\mathcal{H}(\mathbb{Z}_p) \subset H(\mathbb{Q}_p)$. The condition for p to be an unramified prime of good reduction for (H, h) and the definition of the canonical integral model of $\mathrm{Sh}(H, h)$ at p can be found in [35, §1].

4.2. The μ -ordinary Newton polygon.

For a PEL-type Shimura variety, Moonen [24] explicitly computes the slopes of the μ -ordinary Newton polygon at an unramified prime of good reduction in terms of the signature type. Here, for convenience, we recall the formula in the case of the Shimura variety $\text{Sh}(\mu_m, \mathfrak{f})$, following [7, §2.8].

Recall from Sections 2.1-2.2 that σ is the Frobenius action on $\mathcal{T} = \text{Hom}(\mathbb{Q}[\mu_m], \mathbb{C})$ and \mathfrak{D} is the set of σ -orbits in \mathcal{T} . The signature type \mathfrak{f} is given by (2.4). For $\tau \in \mathcal{T}$, recall that $\mathfrak{f}(\tau)$ is the dimension of the eigenspace of V^+ for τ under the μ_m -action.

For a σ -orbit $\mathfrak{o} \in \mathfrak{D}$, let $s(\mathfrak{o})$ be the number of distinct values of $\{\mathfrak{f}(\tau) \mid \tau \in \mathfrak{o}\}$ in $[1, g(\mathfrak{o}) - 1]$. Let $E(1), \dots, E(s(\mathfrak{o}))$ denote these distinct values, ordered such that

$$g(\mathfrak{o}) > E(1) > E(2) > \dots > E(s(\mathfrak{o})) > 0.$$

For convenience, we also write $E(0) := g(\mathfrak{o})$ and $E(s(\mathfrak{o}) + 1) := 0$.

Proposition 4.3. *The μ -ordinary Newton polygon for $\text{Sh}(H, h)$ is given by $\nu_{\mathfrak{o}} = \bigoplus_{\mathfrak{o} \in \mathfrak{D}} \mu(\mathfrak{o})$ where the polygon $\mu(\mathfrak{o})$ has exactly $s(\mathfrak{o}) + 1$ distinct slopes,*

$$0 \leq \lambda(0) < \lambda(1) < \dots < \lambda(s(\mathfrak{o})) \leq 1,$$

and, for each integer t such that $0 \leq t \leq s(\mathfrak{o})$, the slope

$$(4.1) \quad \lambda(t) = \frac{1}{\#\mathfrak{o}} \sum_{i=0}^t \#\{\tau \in \mathfrak{o} \mid \mathfrak{f}(\tau) = E(i)\}$$

occurs in $\mu(\mathfrak{o})$ with multiplicity

$$(4.2) \quad m(\lambda(t)) = (\#\mathfrak{o}) \cdot (E(t) - E(t+1)).$$

Example 4.4. If $\#\mathfrak{o} = 1$ with signature (f_1) , then $\mu(\mathfrak{o})$ has two slopes: 0 with multiplicity $g(\mathfrak{o}) - f_1$ and 1 with multiplicity f_1 .

Example 4.5. If $\mathfrak{o} = \{\tau, \tau^*\}$ with signature (f_1, f_2) and $f_1 \leq f_2$, then $\mu(\mathfrak{o}) = \text{ord}^{2f_1} \oplus \text{ss}^{f_2 - f_1}$.

Proof. If $f_1 \neq 0$ and $f_1 \neq f_2$, then $s(\mathfrak{o}) = 2$; also $E(0) = f_1 + f_2$, $E(1) = f_2$, and $E(2) = f_1$. Then $\lambda(0) = 0$ with $m(0) = 2f_1$; $\lambda(1) = 1/2$ with $m(1/2) = 2(f_2 - f_1)$; and $\lambda(2) = 1$ with $m(2) = 2f_1$. If $f_1 = 0$ or $f_1 = f_2$, then a similar calculation applies. In all cases, $\nu_{\mathfrak{o}} = \text{ord}^{2f_1} \oplus \text{ss}^{f_2 - f_1}$. \square

A similar calculation applies for orbits with $\#\mathfrak{o} = 2$ that are not self-dual, again showing that $\mu(\mathfrak{o})$ has slopes in $\{0, 1/2, 1\}$. Hence, we focus on the orbits of length > 2 when computing the μ -ordinary Newton polygon.

4.3. The Kottwitz method.

Recall that $S = S(m, N, a)$ is an irreducible component of the smallest Shimura subvariety of \mathcal{A}_g containing $Z(m, N, a)$. In [25, Table 1], Moonen lists the 20 monodromy data (m, N, a) (up to equivalence) for which $Z(m, N, a) = S(m, N, a)$. He obtains the list by comparing the dimensions of Z and S : on one hand, $\dim(Z) = N - 3$ since 3 out of the N branch points can be fixed via fractional linear transformations; on the other hand, the signature type \mathfrak{f} determines $\dim(S)$ by [25, Equation (3.3.1)].

As above, p is a rational prime such that $p \nmid m$. We now give an explicit description of the set $\nu(B(\mu_m, f))$ of Newton polygons from Definition 4.2. Let M be the (contravariant) Dieudonné module of an abelian variety $X = \mathcal{X}_x$ for any $x \in \text{Sh}(H_{\mathbb{Q}[\mu_m]}, \mu)(\overline{\mathbb{F}}_p)$. The $\mathbb{Q}[\mu_m]$ -action on X induces a $\mathbb{Q}[\mu_m] \otimes_{\mathbb{Q}} \mathbb{Q}_p$ -action on $X[p^\infty]$ and thus a canonical decomposition

$$X[p^\infty] = \bigoplus_{\mathfrak{o} \in \mathfrak{D}} X[\mathfrak{p}_\mathfrak{o}^\infty].$$

There is an analogous decomposition of Dieudonné modules

$$M = \bigoplus_{\mathfrak{o} \in \mathfrak{D}} M_\mathfrak{o}.$$

Let $\nu(\mathfrak{o})$ denote the Newton polygon of $X[\mathfrak{p}_\mathfrak{o}^\infty]$. By (2.4), for $\mathfrak{o} = \{\tau_0\}$, we have $X[\mathfrak{p}_\mathfrak{o}^\infty] = 0$ and hence $M_\mathfrak{o} = 0$. For convenience, we set $\nu(\{\tau_0\}) = 0$.

Theorem 4.6 (special case of [35, Theorem 1.6]). *The Newton polygons that occur on $S(m, N, a)_{\overline{\mathbb{F}}_p}$ are the Newton polygons $\bigoplus_{\mathfrak{o} \in \mathfrak{D}} \nu(\mathfrak{o})$ satisfying conditions (D), (WA), and (M) below.*

The polarization on X induces a prime-to- p isogeny $X \rightarrow X^\vee$ and the Rosati involution acts on $\mathbb{Q}[\mu_m]$ via the involution $*$. Thus there is an isomorphism $M \cong M^\vee(1)$ compatible with the decomposition and this isomorphism induces certain restrictions on $\nu(\mathfrak{o})$ in the following sense:

condition (D):

- (1) if $\mathfrak{o}^* \neq \mathfrak{o}$, then $M_\mathfrak{o} \cong M_{\mathfrak{o}^*}^\vee(1)$, i.e., $\nu(\mathfrak{o})$ determines $\nu(\mathfrak{o}^*)$. More precisely, if $\nu(\mathfrak{o})$ has slopes $\lambda_1, \dots, \lambda_s$ with multiplicities m_1, \dots, m_s , then $\nu(\mathfrak{o}^*)$ has slopes $1 - \lambda_1, \dots, 1 - \lambda_s$ with multiplicities m_1, \dots, m_s .
- (2) if $\mathfrak{o}^* = \mathfrak{o}$, then $M_\mathfrak{o} \cong M_\mathfrak{o}^\vee(1)$, i.e., $\nu(\mathfrak{o})$ is symmetric. In other words, if λ is a slope of $\nu(\mathfrak{o})$ of multiplicity m_λ , then $1 - \lambda$ is a slope of $\nu(\mathfrak{o})$ of multiplicity m_λ .

By [29, Theorem 4.2], for any $\mathfrak{o} \in \mathfrak{D}$, the Newton polygon $\nu(\mathfrak{o})$ satisfies the weakly admissible condition:

$$\text{condition (WA): } \nu(\mathfrak{o}) \geq \mu(\mathfrak{o}).$$

Recall that $\mathfrak{p}_\mathfrak{o}$ is a prime of $K_{d_\mathfrak{o}}$ with inertia degree $\#\mathfrak{o}$. Note that $K_{d_\mathfrak{o}, \mathfrak{p}_\mathfrak{o}}$ acts on $M_\mathfrak{o}$, which yields:

$$\text{condition (M): the multiplicities of slopes in } \nu(\mathfrak{o}) \text{ are divisible by } \#\mathfrak{o}.$$

4.4. Computation.

We compute all possible Newton polygons for the 20 families from [25, Table 1]; the data can be found in the tables in §6. Here, the label $M[r]$ denotes the r th label in [25, Table 1]. Since the computations are similar for all 20 cases, we give complete details for two examples: first, family $M[17]$ where $m = 7$ is prime; and second, family $M[19]$ where $m = 9$ is not a prime.

Set $\mathfrak{D}' = \mathfrak{D} - \{\{\tau_0\}\}$; from now on, $\mathfrak{o} \in \mathfrak{D}'$. The Newton polygon for X is given by $\bigoplus_{\mathfrak{o} \in \mathfrak{D}'} \nu(\mathfrak{o})$. In the following two examples, since non-symmetric Newton polygons arise, we use different notation from §2.3: when $\gcd(a, b) = 1$, with small abuse of notation, we use $G_{a,b}$ instead of $\nu(G_{a,b})$ to denote the Newton polygon of pure slope $a/(a+b)$ with multiplicity $a+b$.

Example 4.7. We consider family $M[17]$ where $m = 7$, $N = 4$, and $a = (2, 4, 4, 4)$. The signature type is $(1, 2, 0, 2, 0, 1)$. Since $N = 4$, the Shimura variety is 1-dimensional. For each congruence class of $p \neq 7$, there are exactly two possible Newton polygons.

- (1) If $p \equiv 2, 4 \pmod{7}$, then $\#\mathfrak{o} = 3$ and $\#\mathfrak{D} = 2$. In this case, $\mathfrak{D} = \{\mathfrak{o}_1, \mathfrak{o}_2\}$, where $\mathfrak{o}_1 = \{\tau_1, \tau_2, \tau_4\}$ and $\mathfrak{o}_2 = \{\tau_3, \tau_5, \tau_6\}$. Since $\mathfrak{o}_1^* = \mathfrak{o}_2$, by (D), $M_{\mathfrak{o}_1} \cong M_{\mathfrak{o}_2}^\vee(1)$. By Section 4.2, $\mu(\mathfrak{o}_2) = G_{0,1}^3 \oplus G_{1,2}$. By (WA) and (M), if $\nu(\mathfrak{o}_2) \neq \mu(\mathfrak{o}_2)$, then $\nu(\mathfrak{o}_2) = G_{1,5}$. Therefore, by (D),

$$\nu_{\mathfrak{o}} = G_{0,1}^3 \oplus G_{1,2} \oplus G_{2,1} \oplus G_{1,0}^3, \text{ and } \nu_b = G_{1,5} \oplus G_{5,1}.$$

- (2) If $p \equiv 3, 5 \pmod{7}$, then $\#\mathfrak{o} = 6$ and $\#\mathfrak{D} = 1$ and there is only one orbit $\mathfrak{o} = \mathcal{T}$.

By Section 4.2,

$$\nu_{\mathfrak{o}} = \mu(\mathfrak{o}) = G_{1,2}^2 \oplus G_{2,1}^2.$$

By (M), every eigenspace of the Frobenius action on the Dieudonné module M has dimension divisible by 6. Since the interval $(1/3, 1/2)$ contains no rational number with denominator 6, the basic Newton Polygon ν_b is supersingular.

Example 4.8. We compute all possible Newton polygons for family $M[19]$ where $m = 9$, $N = 4$ and $a = (3, 5, 5, 5)$. The signature is $(1, 2, 0, 2, 0, 1, 0, 1)$. Then τ_3, τ_6 have order 3 and τ_i have order 6 for $i = 1, 2, 4, 5, 7, 8$. By (2.4) and Remark 2.3, $g(\tau_3) = g(\tau_6) = 0 + 1 = 1$ and $g(\tau_i) = 2$ for $i = 1, 2, 4, 5, 7, 8$.

To illustrate the idea, we show the computation for $p \equiv 4, 7 \pmod{9}$. In this case, $\#\mathfrak{D}' = 4$, with orbits $\mathfrak{o}_1 = \{\tau_1, \tau_4, \tau_7\}$, $\mathfrak{o}_2 = \{\tau_2, \tau_5, \tau_8\}$, $\mathfrak{o}_3 = \{\tau_3\}$, $\mathfrak{o}_4 = \{\tau_6\}$.

By Section 4.2, $\mu(\mathfrak{o}_3) = G_{0,1}$ and by (WA), this is the only possibility for the Newton polygon of $M_{\mathfrak{o}_3}$. By (D), the only possible Newton polygon on $M_{\mathfrak{o}_3} \oplus M_{\mathfrak{o}_4}$ is $G_{0,1} \oplus G_{1,0}$. Again by Section 4.2, $\mu(\mathfrak{o}_1) = G_{1,2} \oplus G_{2,1}$ on $M_{\mathfrak{o}_1}$. By (M), every slope in the Newton polygon of $M_{\mathfrak{o}_3}$ has multiplicity 3 or 6 and hence the denominator divides 6. By (WA), the first slope lies in the interval $[1/3, 1/2]$. So, there are two possible Newton polygons

$$\nu_{\mathfrak{o}}(\mathfrak{o}_1) = G_{1,2} \oplus G_{2,1}, \text{ and } \nu_b(\mathfrak{o}_1) = G_{1,1}^3.$$

Then, by (D), the only possible Newton polygons on $M_{\mathfrak{o}_1} \oplus M_{\mathfrak{o}_2}$ are

$$\nu_{\mathfrak{o}}(\mathfrak{o}_1, \mathfrak{o}_2) = G_{1,2}^2 \oplus G_{2,1}^2, \text{ and } \nu_b(\mathfrak{o}_1, \mathfrak{o}_2) = G_{1,1}^6.$$

In conclusion, the two possible Newton polygons of the Dieudonné module M are:

$$\nu_{\mathfrak{o}} = G_{0,1} \oplus G_{1,2}^2 \oplus G_{2,1}^2 \oplus G_{1,0}, \text{ and } \nu_b = G_{0,1} \oplus G_{1,1}^6 \oplus G_{1,0}.$$

4.5. The μ -ordinary Ekedahl–Oort type.

The Ekedahl–Oort type is a combinatorial invariant classifying the structure of the reduction modulo p of the Dieudonné module. In [24, Theorem 1.3.7], Moonen shows that the largest Ekedahl–Oort stratum coincides with the μ -ordinary Newton stratum. We recall the \mathbb{E} -module structure of the mod p Dieudonné module for the μ -ordinary locus, given in [24, §1.2.3].

Fix $\mathfrak{o} \in \mathfrak{D}$ and $t \in \{0, \dots, s(\mathfrak{o})\}$. We define the \mathbb{E} -module:

$$N_t(\mathfrak{o}) := \bigoplus_{\tau \in \mathfrak{o}} \overline{\mathbb{F}}_p e_\tau,$$

by

$$\begin{aligned} F(e_\tau) &= e_{\tau^\sigma} \text{ if } f(\tau) \leq E(t+1), 0 \text{ otherwise,} \\ V(e_{\tau^\sigma}) &= e_\tau \text{ if } f(\tau) > E(t+1), 0 \text{ otherwise.} \end{aligned}$$

Then $N_t(\mathfrak{o})$ arises as the mod p -reduction of an isoclinic Dieudonné module of slope $\lambda(t)$.

The reduction modulo p of the μ -ordinary Dieudonné module of $\text{Sh}(H, h)_{\overline{\mathbb{F}}_p}$ is

$$(4.3) \quad N_{\mathfrak{o}} = \bigoplus_{\mathfrak{o} \in \mathfrak{D}} N(\mathfrak{o}), \text{ where } N(\mathfrak{o}) = \bigoplus_{t=0}^{s(\mathfrak{o})} N_t(\mathfrak{o})^{E(t)-E(t+1)}.$$

This agrees with the formula in [24, §1.2.3]: note that Moonen defines an \mathbb{E} -module for each $j \in \{1, \dots, g(\mathfrak{o})\}$; if $g(\mathfrak{o}) - E(t+1) < j \leq g(\mathfrak{o}) - E(t)$, then the \mathbb{E} -modules have the same structure, which determines the multiplicity of each factor $N_t(\mathfrak{o})$ in (4.3).

Recall from Section 2.4 that L denotes the \mathbb{E} -module of an ordinary elliptic curve, and $N_{1,1}$ that of a supersingular elliptic curve.

Example 4.9. If $\mathfrak{o} = \{\tau, \tau^* = \tau^\sigma\}$ with signature (f_1, f_2) with $f_1 \leq f_2$, then $N(\mathfrak{o}) \simeq L^{2f_1} \oplus N_{1,1}^{f_2-f_1}$.

Note that Example 4.9 is not automatic from Example 4.5 since several Dieudonné modules can occur with Newton polygon $ss^{f_2-f_1}$; the one in Example 4.9 has a -number $f_2 - f_1$; it decomposes into dimension 2 subspaces stable under F and V .

Proof. If $f_1 \neq 0$ and $f_1 \neq f_2$, then by the computation in Example 4.5 and (4.3) $N_0(\mathfrak{o})$ is generated by e_τ, e_{τ^σ} with $F = \text{id}$, the identity map, and $V = 0$; $N_2(\mathfrak{o})$ is generated by e_τ, e_{τ^σ} with $V = \text{id}$ and $F = 0$; and $N_1(\mathfrak{o})$ is generated by e_τ, e_{τ^σ} with $F(e_\tau) = V(e_\tau) = e_{\tau^\sigma}$ and $F(e_{\tau^\sigma}) = V(e_{\tau^\sigma}) = 0$. A similar calculation works when $f_1 = f_2$ or $f_1 = 0$. In all cases, $N(\mathfrak{o}) \simeq L^{2f_1} \oplus N_{1,1}^{f_2-f_1}$. \square

A similar calculation applies for all orbits with $\#\mathfrak{o} = 2$. Hence, we focus on orbits of length > 2 when computing the μ -ordinary mod p Dieudonné module.

Example 4.10. We compute the μ -ordinary mod p Dieudonné module $N_{\mathfrak{o}}$ for family $M[17]$ where $m = 7$. The signature is $(1, 2, 0, 2, 0, 1)$ and $g(\tau) = 2$.

(1) If $p \equiv 2, 4 \pmod{7}$, then the p -rank is 3, and the local-local part of $N_{\mathfrak{o}}$ has 6 generators with action of F and V given by:

$$\begin{array}{c|ccc} t = 0, E(t+1) = 1 & e_1 & e_2 & e_4 \\ \hline F & e_2 & 0 & 0 \\ \hline V & e_4 & 0 & e_2 \end{array} \text{ and } \begin{array}{c|ccc} t = 1, E(t+1) = 0 & e'_3 & e'_5 & e'_6 \\ \hline F & e'_5 & e'_6 & 0 \\ \hline V & e'_6 & 0 & 0 \end{array}$$

Thus $N_{\mathfrak{o}} \simeq L^3 \oplus (\mathbb{E}/\mathbb{E}(F - V^2))e_1 \oplus (\mathbb{E}/\mathbb{E}(F^2 - V))e'_3$. One can compute that $N_{\mathfrak{o}}$ has Ekedahl-Oort type $[1, 2, 3, 3, 4, 4]$.

(2) If $p \equiv 3, 5 \pmod{7}$, then $N_{\mathfrak{o}}$ has 12 generators with the actions of F and V given by:

$$\begin{array}{c|ccccc} t = 0, E(t+1) = 1 & e_1 & e_3 & e_2 & e_6 & e_4 & e_5 \\ \hline F & e_3 & e_2 & 0 & e_4 & 0 & e_1 \\ \hline V & 0 & 0 & 0 & e_2 & 0 & e_4 \end{array} \text{ and } \begin{array}{c|ccccc} t = 1, E(t+1) = 0 & e'_1 & e'_3 & e'_2 & e'_6 & e'_4 & e'_5 \\ \hline F & 0 & e'_2 & 0 & 0 & 0 & e'_1 \\ \hline V & 0 & e'_1 & 0 & e'_2 & e'_6 & e'_4 \end{array}$$

Thus $N_o = \mathbb{E}\langle e_5, e_6 \rangle / \mathbb{E}(F^3 e_5 - V e_6, V e_5 - F e_6) \oplus \mathbb{E}\langle e'_3, e'_5 \rangle / \mathbb{E}(F e'_3 - V^3 e'_5, V e'_3 - F e'_5)$. In total, N_o has 4 generators, thus a -number 4. One can compute that N_o has Ekedahl-Oort type $[0, 1, 2, 2, 2, 2]$.

Example 4.11. We compute the μ -ordinary mod p Dieudonné module N_o for family M [19] where $m = 9$. The signature is $(1, 2, 0, 2, 0, 1, 0, 1)$ and $g(\tau_1) = 2$ and $g(\tau_3) = 1$.

- (1) If $p \equiv 2, 5 \pmod{9}$, there are two orbits \mathfrak{o}_3 and \mathfrak{o}_1 . By Example 4.9, $N(\mathfrak{o}_3) \simeq (\mathbb{E}/\mathbb{E}(F - V))e_3$. For the orbit \mathfrak{o}_1 of τ_1 , we see that $N(\mathfrak{o}_1)$ has 12 generators with the actions of F and V given by:

$t = 0, E(t+1) = 1$	e_1	e_2	e_4	e_8	e_7	e_5
F	e_2	0	0	e_7	e_5	e_1
V	0	0	e_2	e_4	0	0

and

$t = 1, E(t+1) = 0$	e'_1	e'_2	e'_4	e'_8	e'_7	e'_5
F	0	0	0	0	e'_5	e'_1
V	0	e'_1	e'_2	e'_4	e'_8	0

Then $N(\mathfrak{o}_1) = (\mathbb{E}/\mathbb{E}(F^4 - V^2))e_8 \oplus (\mathbb{E}/\mathbb{E}(V^4 - F^2))e'_7$. The Ekedahl-Oort type of $N(\mathfrak{o}_1)$ is $[0, 1, 2, 2, 3, 4]$. In total, N_o has 3 generators, thus a -number 3.

- (2) If $p \equiv 4, 7 \pmod{9}$, there are four orbits. We check that:

$$\begin{aligned} N(\mathfrak{o}_3) \oplus N(\mathfrak{o}_6) &\simeq L \\ N(\mathfrak{o}_1) &\simeq (\mathbb{E}/\mathbb{E}(F^2 - V))e_7 \oplus (\mathbb{E}/\mathbb{E}(F - V^2))e'_7 \\ N(\mathfrak{o}_2) &\simeq (\mathbb{E}/\mathbb{E}(F^2 - V))e_8 \oplus (\mathbb{E}/\mathbb{E}(F - V^2))e'_5. \end{aligned}$$

One interesting feature is that the Newton polygon $(1/3, 2/3)^2$ matches with two different Dieudonné modules in Examples 4.10(2) and 4.11(2).

5. NEWTON POLYGONS OF SMOOTH CURVES

Let (m, N, a) be a monodromy datum. If it is the monodromy datum of one of the twenty special families from [25, Table 1], we call it *special*. Fix a prime p with $p \nmid m$.

For a Newton polygon $\nu \in \nu(B(\mu_m, \mathfrak{f}))$, by definition ν occurs as the Newton polygon of an abelian variety represented by a point of $Z(m, N, a) \subset \mathcal{A}_g$. This abelian variety is the Jacobian of a curve of *compact type*. One can ask whether ν occurs as the Newton polygon of the Jacobian of a curve that is *smooth*. By Definition 2.1, it is equivalent to ask whether ν occurs as the Newton polygon for a point of $Z^0(m, N, a)$. In general, this is a subtle and difficult question. In this section, we provide an answer in three settings: when the Newton polygon ν is μ -ordinary in Section 5.1, PEL-indecomposable in Section 5.2, or basic in Section 5.3. In Section 5.4, we use these three criteria and the purity theorem to prove that all of the Newton polygons for the special families in [25, Table 1] occur for Jacobians of smooth curves except for a few of the supersingular ones.

When there is no risk of confusion, we write Z and Z^0 for $Z(m, N, a)$ and $Z^0(m, N, a)$ respectively.

5.1. The μ -ordinary case.

Proposition 5.1. *For any special monodromy datum (m, N, a) , and any prime p with $p \nmid m$, the μ -ordinary Newton polygon $\nu_o \in \nu(B(\mu_m, \mathfrak{f}))$ and the μ -ordinary mod p Dieudonné module N_o occur for the Jacobian of a smooth curve in $Z(m, N, a)$.*

Proof. Both Z^0 and the μ -ordinary Newton polygon (or Ekedahl–Oort) stratum $Z(\nu_0)$ are open and dense in Z , thus their intersection is non-empty. \square

5.2. PEL-indecomposable Newton polygons.

Suppose C is a curve of genus g of compact type. If the Newton polygon of $\text{Jac}(C)$ is indecomposable as a symmetric Newton polygon of height $2g$, (e.g., $G_{1,g-1} \oplus G_{g-1,1}$), then C is necessarily smooth. In this section, we refine that observation for curves of compact type which are μ_m -Galois covers of the projective line.

Remark 5.2. The following material is based on information about the boundary of the Hurwitz space of cyclic covers of the projective line, see e.g., [38, Chapter 4] and [8]. A point ξ of $Z(m, N, a) - Z^0(m, N, a)$ represents the Jacobian of a singular curve Y . This curve Y is of compact type since $\text{Jac}(Y)$ is an abelian variety. Furthermore, there is a μ_m -cover $\psi : Y \rightarrow X$ where X is a singular curve of genus 0. The cover ψ is the join of two μ_m -covers $\psi_1 : Y_1 \rightarrow X_1$ and $\psi_2 : Y_2 \rightarrow X_2$, where the curves are clutched together in ordinary double points. Since ξ is in the closure of $Z^0(m, N, a)$, the cover ψ is *admissible*, meaning that at each clutching point the canonical generators of inertia for ψ_1 and ψ_2 are inverses. Since Y is connected, either Y_1 or Y_2 is connected; without loss of generality, say Y_1 is connected and let r be the number of components of Y_2 . Then ψ_2 is induced from a $\mu_{m/r}$ -cover $\psi_2^{\text{co}} : Y_2^{\text{co}} \rightarrow X_2$, where Y_2^{co} is isomorphic to a connected component of Y_2 . Finally, the fact that Y is of compact type implies that each component of Y_2 is clutched together with Y_1 at exactly one point.

In the situation of Remark 5.2, the monodromy data for the covers satisfy certain numerical conditions as follows.

Notation 5.3. Let (m, N, a) be the monodromy datum for ψ . Let (m, N_1, α_1) be the monodromy datum for ψ_1 . Let $(m/r, N_2, \alpha_2)$ be the monodromy datum for ψ_2^{co} . Write

$$\alpha_1 = (\alpha_1(1), \dots, \alpha_1(N_1)), \alpha_2 = (\alpha_2(1), \dots, \alpha_2(N_2)).$$

Then $N_1 + N_2 = N + 2$. Let $\tilde{\alpha}_2 = \text{Ind}_{m/r}^m \alpha_2 := (r\alpha_2(1), \dots, r\alpha_2(N_2))$; we call it the monodromy datum for ψ_2 , even if $r \neq 1$. The admissible condition is that $\alpha_1(N_1) \equiv -r\alpha_2(1) \pmod{m}$. Then (possibly after rearranging)

$$a = (\alpha_1(1), \dots, \alpha_1(N_1 - 1), r\alpha_2(2), \dots, r\alpha_2(N_2)).$$

Finally, $r = \gcd(m, \alpha_1(N_1))$.

Definition 5.4. We say that the pair $\alpha_1, \tilde{\alpha}_2$ is a degeneration of compact type of the inertia type a if the numerical conditions in Notation 5.3 are satisfied.

Suppose that $\alpha_1, \tilde{\alpha}_2$ is a degeneration of compact type of a . For $i = 1, 2$, write $Z_1 := Z(m, N_1, \alpha_1) \subset \mathcal{A}_{g_1}$ and $Z_2 := Z(m/r, N_2, \alpha_2) \subset \mathcal{A}_{g_2}$. The numerical conditions imply that $g_1 + rg_2 = g$. We define $\oplus : \mathcal{A}_{g_1} \times \mathcal{A}_{g_2} \rightarrow \mathcal{A}_g$, by $(A_1, A_2) \mapsto A_1 \oplus A_2^r$. Let f (resp. f_1, f_2) be the signature for a (resp. α_1, α_2). Consider $\text{Sh}(\mu_m, f_1)$ (resp. $\text{Sh}(\mu_{m/r}, f_2)$), the smallest PEL-type Shimura stack containing Z_1 (resp. Z_2) as introduced in Section 3.3. It follows from the definitions that

$$\oplus (\text{Sh}(\mu_m, f_1) \times \text{Sh}(\mu_{m/r}, f_2)) \subseteq \text{Sh}(\mu_m, f).$$

Definition 5.5. Let (m, N, a) be a monodromy datum and let f be the corresponding signature. A Newton polygon $v \in v(B(\mu_m, f))$ is *PEL-decomposable* if $v = v_1 \oplus v_2'$ for some $v_1 \in v(B(\mu_m, f_1))$ and $v_2 \in v(B(\mu_{m/r}, f_2))$, for some pair f_1, f_2 of signatures arising from a degeneration of compact type of a .

We note that the condition of being PEL-decomposable depends on p modulo m .

Proposition 5.6. For a special monodromy datum (m, N, a) , and a prime p with $p \nmid m$, if $v \in v(B(\mu_m, f))$ is not PEL-decomposable, then v occurs as the Newton polygon of the Jacobian of a smooth curve in $Z(m, N, a)$.

Proof. For a special monodromy datum, every Newton polygon $v \in v(B(\mu_m, f))$ occurs for a point of $Z(m, N, a)$. By Remark 5.2, if this point is the Jacobian of a singular curve, then v is PEL-decomposable by Notation 5.3 and Definitions 5.4 and 5.5. \square

We compute all PEL-decomposable Newton polygons, for families $M[1 - 20]$ from [25, Table 1]; the data for families $M[3 - 20]$ can be found in Lemma 6.4. The computations are similar in all cases, so we give complete details only for $M[15]$.

Example 5.7. For family $M[15]$ where $m = 8$, $N = 4$, and $a = (2, 4, 5, 5)$; the signature is $f = (1, 1, 0, 0, 2, 0, 1)$. There are two degenerations of compact type of a :

$$\alpha_1 = (2, 5, 1), \alpha_2 = (7, 4, 5) \text{ or } \alpha_1' = (5, 5, 6), \tilde{\alpha}_2' = \text{Ind}_4^8(1, 2, 1).$$

The corresponding partitions of the signature f are respectively:

$$\begin{aligned} f_1 &= (1, 1, 0, 0, 1, 0, 0), f_2 = (0, 0, 0, 0, 1, 0, 1); \\ f_1' &= (0, 1, 0, 0, 1, 0, 1), \tilde{f}_2' = (1, 0, 0, 0, 1, 0, 0). \end{aligned}$$

Thus, for each non-trivial congruence class of $p \pmod 8$, there are at most two PEL-decomposable Newton polygons η, η' . We compute η, η' using the Kottwitz method.

- (1) If $p \equiv 1 \pmod 8$: $\eta = \eta' = v_o = G_{0,1}^5 \oplus G_{1,0}^5$.
- (2) If $p \equiv 3 \pmod 8$: $\eta = \eta' = v_o = G_{0,1}^2 \oplus G_{1,1}^3 \oplus G_{1,0}^2$.
- (3) If $p \equiv 5 \pmod 8$: $\eta = \eta' = v_o = G_{0,1}^3 \oplus G_{1,1}^2 \oplus G_{1,0}^3$.
- (4) If $p \equiv 7 \pmod 8$: $\eta = \eta' = v_b = G_{1,1}^5$.

We deduce that the following Newton polygons are not PEL-decomposable and thus occur as the Newton polygon of the Jacobian of a smooth curve in $Z^0(8, 4, a)$:

- (1) for $p \equiv 1 \pmod 8$, $v_b = G_{0,1}^3 \oplus G_{1,1}^2 \oplus G_{1,0}^3$;
- (2) for $p \equiv 3 \pmod 8$, $v_b = G_{1,3} \oplus G_{3,1} \oplus G_{1,1}$;
- (3) for $p \equiv 5 \pmod 8$, $v_b = G_{0,1} \oplus G_{1,3} \oplus G_{3,1} \oplus G_{1,0}$.
- (4) for $p \equiv 7 \pmod 8$, $v_o = G_{0,1}^2 \oplus G_{1,1}^3 \oplus G_{1,0}^2$.

Thus, combining Propositions 5.1 and 5.6, in this example we conclude that all Newton polygons $v \in B(\mu_m, f)$ occur on $Z^0(8, 4, a)$, thus occur as Newton polygons of the Jacobian of a smooth curve, except for $p \equiv 7 \pmod 8$ and $v = G_{1,1}^5$. This last case is addressed by Proposition 5.8.

5.3. Basic Newton polygons.

Consider a special monodromy datum (m, N, a) , and a prime p with $p \nmid m$, such that the set $\nu(B(\mu_m, \mathfrak{f}))$ contains exactly two Newton polygons, namely the μ -ordinary polygon ν_o and the basic polygon ν_b . In other words, assume that the closed (basic) Newton stratum has codimension 1. Note that this condition depends on p , as seen in the tables. For the special families of [25, Table 1], this condition implies that the dimension of $Z = Z(m, N, a)$ is either 1 or 2, or equivalently $N = 4, 5$.

Under this condition, we prove that the basic polygon ν_b occurs as the Newton polygon of a smooth curve in Z .

Proposition 5.8. *For any special monodromy datum (m, N, a) with $N = 4$, and any sufficiently large prime p with $p \nmid m$, the basic polygon $\nu_b \in \nu(B(\mu_m, \mathfrak{f}))$ occurs as the Newton polygon of the Jacobian of a smooth curve in $Z(m, N, a)$.*

The key input of the proof is that the number of irreducible components of the basic locus in the mod p reduction of a simple unitary Shimura variety is unbounded as p goes to ∞ . We prove this statement in Theorem 8.1 in the appendix.

Proof. For $N = 4$ (dimension 1), the set $\nu(B(\mu_m, \mathfrak{f}))$ contains exactly two Newton polygons for any prime $p \nmid m$. We argue by contradiction; let us assume that the basic Newton stratum $Z_p(\nu_b)$ is contained in the mod p fiber ∂Z_p of the boundary $\partial Z := Z - Z^0$. Since the closed substack $Z_p(\nu_b)$ has codimension 1 in Z_p , if $Z_p(\nu_b) \subset \partial Z_p$, then it is equal to a union of irreducible components of ∂Z_p , which represent the singular curves in the family. On one hand, the number of irreducible components of the mod p fiber ∂Z_p has an natural upper bound which is independent of p .

On the other hand, the number of irreducible components of the basic Newton stratum of the mod p reduction of any irreducible component of the unitary Shimura variety attached to the special monodromy datum $(m, 4, a)$ grows to infinity with p (see appendix Theorem 8.1, Corollary 8.3). Hence the contradiction, for p sufficiently large. \square

Example 5.9. For family $M[17]$, the Shimura variety $S(7, 4, (2, 4, 4, 4))$ has dimension 1. We deduce that for p sufficiently large, and $p \equiv 3, 5, 6 \pmod{7}$, the basic Newton polygon $\nu_b = G_{1,1}^6$ occurs as the Newton polygon of the Jacobian of a smooth curve in Z .

Remark 5.10. When $N = 5$ (dimension 2), to apply Theorem 8.1 we have to further assume p is split in the quadratic imaginary extension. This condition excludes primes $p \equiv -1 \pmod{m}$ for the special families $M[6]$, $M[8]$, and $M[14]$, and primes $p \not\equiv 1 \pmod{5}$ for $M[16]$.

5.4. Conclusion.

We apply the three criteria above to the special families in [25, Table 1] to prove that all but a few of the Newton polygons occur for the Jacobian of a smooth curve in the family.

Let $M[r]$ denote one of the special families from [25, Table 1] and (m, N, a) its monodromy datum. Section 6 contains the list of all the Newton polygons $\nu \in \nu(B(\mu_m, \mathfrak{f}))$. Note that $\dim(Z(m, N, a))$ equals 3 for the family $M[r]$ when $r = 2, 10$, equals 2 for $r = 6, 8, 14, 16$, and equals 1 otherwise.

Theorem 5.11. *Let (m, N, a) denote the monodromy datum for one of the special families from [25, Table 1]. Assume $p \nmid m$. Let $v \in \nu(B(\mu_m, \mathfrak{f}))$ be a Newton polygon occurring on $Z(m, N, a)$. Then v occurs on $Z^0(m, N, a)$, meaning that v occurs as the Newton polygon of the Jacobian of a smooth curve in the family unless either:*

- (1) $\dim(Z(m, N, a)) = 1$, v is supersingular, and p is not sufficiently large; or
- (2) $\dim(Z(m, N, a)) \geq 2$, and v is supersingular.

In cases (1) and (2), we prove in [21, Theorem 6.1] that there exists a smooth curve in the family which is supersingular for sufficiently large p .

Proof. For the family $M[1]$ (resp. $M[2]$), where $m = 2$, the Shimura variety $Z(m, N, a)$ is simply \mathcal{A}_1 (resp. \mathcal{A}_2). Thus all Newton polygons for $M[1]$ and $M[2]$ are already known to occur for Jacobians of smooth curves.

Consider one of the families $M[3 - 20]$, where $m \geq 3$. For the μ -ordinary Newton polygon v_o , the result follows from Proposition 5.1. Suppose that $v \in \nu(B(\mu_m, \mathfrak{f}))$ is not μ -ordinary. We distinguish two cases: $\dim(Z(m, N, a)) = 1$ and $\dim(Z(m, N, a)) \geq 2$.

Suppose $\dim(Z(m, N, a)) = 1$. Then the set $\nu(B(\mu_m, \mathfrak{f}))$ contains exactly two polygons; since v is not μ -ordinary, then v is the basic Newton polygon v_b . Direct computations (Lemma 6.4) show that if v_b is not supersingular, then it is PEL-indecomposable and the result follows from Proposition 5.6. If v_b is supersingular, by Proposition 5.8, v_b occurs for a smooth curve if p is sufficiently large.

Suppose $\dim(Z(m, N, a)) \geq 2$ and the basic Newton polygon $v_b \in \nu(B(\mu_m, \mathfrak{f}))$ is not supersingular. From the tables in Section 6, this occurs exactly when $p \equiv 1 \pmod{m}$. Again, direct computations (Lemma 6.4) show that v_b is PEL-indecomposable. By Proposition 5.6, v_b occurs for the Jacobian of a smooth curve in the family, i.e., $Z^0[v_b] \neq \emptyset$, where $Z^0 = Z^0(m, N, a)$. Furthermore, note that $\nu(B(\mu_m, \mathfrak{f}))$ is totally ordered and

$$\text{codim}(Z^0[v_b], Z^0) = \#\nu(B(\mu_m, \mathfrak{f})) - 1.$$

By the de Jong–Oort purity result for the stratification by Newton polygons [5, Theorem 4.1], we deduce that $Z^0[v]$ is non empty, for all $v \in \nu(B(\mu_m, \mathfrak{f}))$.

Suppose $\dim(Z(m, N, a)) \geq 2$ and the basic Newton polygon $v_b \in \nu(B(\mu_m, \mathfrak{f}))$ is supersingular. The proof is complete unless $\nu(B(\mu_m, \mathfrak{f}))$ contains more than 2 Newton polygons. From the tables in Section 6, this happens only for the family $M[10]$ when $p \equiv 2 \pmod{3}$, in which case $\nu(B(\mu_m, \mathfrak{f}))$ contains exactly three polygons: the μ -ordinary v_o ; the basic Newton polygon v_b ; and $\eta = (1/4, 3/4)$, which is PEL-indecomposable. By Proposition 5.6, η occurs for the Jacobian of a smooth curve in the family $M[10]$. \square

6. TABLES

The following tables contain all the Newton polygons and the μ -ordinary Dieudonné modules which occur for the cyclic covers of the projective line arising from the special families of Moonen. The label $M[r]$ denotes the r th label in [25, Table 1]. We organize the tables for $M[1] - M[20]$ by the value of m starting with the ones with $\phi(m) = 1, 2$.

Notation 6.1. The degree of the cover is m , the inertia type is $a = (a_1, \dots, a_N)$, the signature is $\mathfrak{f} = (f(1), \dots, f(m-1))$. The prime orbits are denoted by \mathfrak{D} .

Notation 6.2. Let $ord = \{0, 1\}$ (resp. $ss = \{1/2, 1/2\}$) be the ordinary (resp. supersingular) Newton polygon. For $s, t \in \mathbb{N}$, with $s < t$ and $\gcd(s, t) = 1$, let $(s/t, (t-s)/t)$ denote the Newton polygon with slopes s/t and $(t-s)/t$, each occurring with multiplicity t .

Notation 6.3. As in Section 2.4, let \mathbb{E} denote the non-commutative ring generated by F and V with $FV = VF = 0$. Let EO-type stand for Ekedahl-Oort type. Define:

- $L = \mathbb{E}/\mathbb{E}(F, V - 1) \oplus \mathbb{E}/\mathbb{E}(V, F - 1)$; it has rank 2, p -rank 1, and EO-type [1];
- $N_{r,1} = \mathbb{E}/\mathbb{E}(F^r - V^r)$; rank $2r$, p -rank 0, a -number 1, and EO-type $[0, 1, \dots, r-1]$;
- $N_{r,2} = \mathbb{E}/\mathbb{E}(F^{r-1} - V) \oplus \mathbb{E}/\mathbb{E}(V^{r-1} - F)$; it has rank $2r$, p -rank 0, a -number 2, and EO-type $[0, 1, \dots, r-3, r-2, r-2]$.

6.1. Tables for m with $m = 2, 3, 4, 6$.

Here are the tables for Moonen's special families when $\phi(m) \leq 2$. Because of previous work, e.g., [3, 28, 22], we do not produce any new examples of Newton polygons and Dieudonné modules for smooth curves in this subsection.

When $\phi(m) \leq 2$, each orbit of the eigenspaces under Frobenius has length 1 or 2. By Examples 4.4 and 4.5, this implies that the μ -ordinary Dieudonné module N_o is determined by the μ -ordinary Newton polygon v_o ; if $v_o = ord^a \oplus ss^b$, then $N_o \simeq L^a \oplus N_{1,1}^b$.

For $m = 2$ with p odd: Family $M[1]$ has $a = (1, 1, 1, 1)$, $f = (1)$, and ord and ss occur. Family $M[2]$ has $a = (1, 1, 1, 1, 1, 1)$, $f = (2)$, and ord^2 , $ord \oplus ss$, and ss^2 occur.

$m = 3$	a	f	$p \equiv 1 \pmod{3}$	$p \equiv 2 \pmod{3}$
$M[3]$	$(1, 1, 2, 2)$	$(1, 1)$	ord^2 ss^2	ord^2 ss^2
$M[6]$	$(1, 1, 1, 1, 2)$	$(2, 1)$	ord^3 $ord \oplus ss^2$ $(1/3, 2/3)$	$ord^2 \oplus ss$ ss^3
$M[10]$	$(1, 1, 1, 1, 1, 1)$	$(3, 1)$	ord^4 $ord^2 \oplus ss^2$ $ord \oplus (1/3, 2/3)$ $(1/4, 3/4)$	$ord^2 \oplus ss^2$ $(1/4, 3/4)$ ss^4

$m = 4$	a	f	$p \equiv 1 \pmod{4}$	$p \equiv 3 \pmod{4}$
$M[4]$	$(1, 2, 2, 3)$	$(1, 0, 1)$	ord^2 ss^2	ord^2 ss^2
$M[7]$	$(1, 1, 1, 1)$	$(2, 1, 0)$	ord^3 $ord^2 \oplus ss$	$ord \oplus ss^2$ ss^3
$M[8]$	$(1, 1, 2, 2, 2)$	$(2, 0, 1)$	ord^3 $ord \oplus ss^2$ $(1/3, 2/3)$	$ord^2 \oplus ss$ ss^3

$m = 6$	p	1 mod 6	5 mod 6
a	$\begin{array}{c} \mathfrak{D} \\ \mathfrak{f} \end{array}$	split	(1, 5), (2, 4), (3)
(2,3,3,4) $M[5]$	(1,0,0,0,1)	ord^2 ss^2	ord^2 ss^2
(1,3,4,4) $M[9]$	(1,1,0,0,1)	ord^3 $ord \oplus ss^2$	$ord^2 \oplus ss$ ss^3
(1,1,1,3) $M[12]$	(2,1,1,0,0)	ord^4 $ord^3 \oplus ss$	$ord \oplus ss^3$ ss^4
(1,1,2,2) $M[13]$	(2,1,0,1,0)	ord^4 $ord^2 \oplus ss^2$	$ord^2 \oplus ss^2$ ss^4
(2,2,2,3,3) $M[14]$	(2,0,0,1,1)	ord^4 $ord^2 \oplus ss^2$ $ord \oplus (1/3, 2/3)$	$ord^2 \oplus ss^2$ ss^4

6.2. Tables for m with $m = 5, 7, 8, 9, 10, 12$.

Here are the tables of Newton polygons for Moonen's special families when $\phi(m) > 2$. We include the μ -ordinary Dieudonné module N_o when the prime orbits have $\#o > 2$; when $\#o \leq 2$, then N_o can be computed from Examples 4.4 and 4.9.

$m = 5$	p	1 mod 5	2, 3 mod 5	4 mod 5
a	$\begin{array}{c} \mathfrak{D} \\ \mathfrak{f} \end{array}$	split	(1, 2, 3, 4)	(1, 4), (2, 3)
(1, 3, 3, 3) $M[11]$	(1, 2, 0, 1)	ord^4 $ord^2 \oplus ss^2$	(1/4, 3/4) ss^4	$ord^2 \oplus ss^2$ ss^4
(2, 2, 2, 2, 2) $M[16]$	(2, 0, 3, 1)	ord^6 $ord^4 \oplus ss^2$ $ord^3 \oplus (1/3, 2/3)$	(1/4, 3/4) \oplus ss^2 ss^6	$ord^2 \oplus ss^4$ ss^6

When $p \equiv 2, 3 \pmod{5}$: for $M[11]$, $N_o \simeq N_{4,2}$; for $M[16]$, $N_o \simeq N_{4,2} \oplus N_{2,1}$.

$m = 7$	p	1 mod 7	2, 4 mod 7	3, 5 mod 7	6 mod 7
a	$\begin{array}{c} \mathfrak{D} \\ \mathfrak{f} \end{array}$	split	(1, 2, 4), (3, 5, 6)	(1, 2, 3, 4, 5, 6)	(1, 6), (2, 5), (3, 4)
(2,4,4,4) $M[17]$	(1,2,0,2,0,1)	ord^6 $ord^4 \oplus ss^2$	$ord^3 \oplus (1/3, 2/3)$ (1/6, 5/6)	(1/3, 2/3) ² ss^6	$ord^2 \oplus ss^4$ ss^6

By Example 4.10, for $M[17]$: when $p \equiv 2, 4 \pmod{7}$, then $N_o \simeq L^3 \oplus N_{3,2}$; when $p \equiv 3, 5 \pmod{7}$, then $N_o \simeq \mathbb{E}\langle e_5, e_6 \rangle / \mathbb{E}(F^3 e_5 - Ve_6, Ve_5 - Fe_6) \oplus \mathbb{E}\langle e'_3, e'_5 \rangle / \mathbb{E}(Fe'_3 - V^3 e'_5, Ve'_3 - Fe'_5)$.

$m = 8$	p	1 mod 8	3 mod 8	5 mod 8	7 mod 8
a	$\begin{array}{c} \mathfrak{D} \\ \mathfrak{f} \end{array}$	split	(1, 3), (2, 6) (5, 7), (4)	(1, 5), (3, 7) (2), (4), (6)	(1, 7), (2, 6) (3, 5), (4)
(2,4,5,5) $M[15]$	(1,1,0,0,2,0,1)	ord^5 $ord^3 \oplus ss^2$	$ord^2 \oplus ss^3$ (1/4, 3/4) \oplus ss	$ord^3 \oplus ss^2$ $ord \oplus (1/4, 3/4)$	$ord^2 \oplus ss^3$ ss^5

$m = 9$	p	1 mod 9	2, 5 mod 9	4, 7 mod 9	8 mod 9
a	$\begin{array}{c} \mathfrak{D} \\ \diagdown \\ \mathfrak{f} \end{array}$	split	(1, 2, 4, 8, 7, 5) (3, 6)	(1, 4, 7), (2, 8, 5) (3), (6)	(1, 8), (2, 7) (4, 5), (3, 6)
(3, 5, 5, 5) $M[19]$	(1, 2, 0, 2, 0, 1, 0, 1)	ord^7 $ord^5 \oplus ss^2$	$(1/3, 2/3)^2 \oplus ss$ ss^7	$ord \oplus (1/3, 2/3)^2$ $ord \oplus ss^6$	$ord^2 \oplus ss^5$ ss^7

By Example 4.11, for $M[19]$: when $p \equiv 4, 7 \pmod{9}$, then $N_o \simeq L \oplus N_{3,2}^2$; when $p \equiv 2, 5 \pmod{9}$, then $N_o \simeq \mathbb{E}/\mathbb{E}(F^4 - V^2) \oplus \mathbb{E}/\mathbb{E}(V^4 - F^2) \oplus \mathbb{E}/\mathbb{E}(F - V)$.

$m = 10$	p	1 mod 10	3, 7 mod 10	9 mod 10
a	$\begin{array}{c} \mathfrak{D} \\ \diagdown \\ \mathfrak{f} \end{array}$	split	(1, 3, 9, 7) (2, 6, 8, 4), (5)	(1, 9), (2, 8) (3, 7), (4, 6), (5)
(3, 5, 6, 6) $M[18]$	(1, 1, 0, 1, 0, 0, 2, 0, 1)	ord^6 $ord^4 \oplus ss^2$	$(1/4, 3/4) \oplus ss^2$ ss^6	$ord^2 \oplus ss^4$ ss^6

For $M[18]$: when $p \equiv 3, 7 \pmod{10}$, then $N_o \simeq N_{4,2} \oplus N_{2,1}$.

$m = 12$	p	1 mod 12	5 mod 12	7 mod 12	11 mod 12
a	$\begin{array}{c} \mathfrak{D} \\ \diagdown \\ \mathfrak{f} \end{array}$	split	(1, 5), (2, 10), (3) (4, 8), (6), (7, 11), (9)	(1, 7), (3, 9) (2), (4), (5, 11) (6), (8), (10)	(1, 11), (4, 8) (3, 9), (2, 10) (5, 7), (6)
(4, 6, 7, 7) $M[20]$	(1, 1, 0, 1, 0, 0, 2, 0, 1, 0, 1)	ord^7 $ord^5 \oplus ss^2$	$ord^3 \oplus ss^4$ $ord \oplus (1/4, 3/4) \oplus ss^2$	$ord^4 \oplus ss^3$ $ord^2 \oplus ss^5$	$ord^2 \oplus ss^5$ ss^7

6.3. PEL-decomposable Newton polygons. The next lemma contains the data of all the PEL-decomposable Newton polygons for families $M[3 - 20]$ from [25, Table 1]. We refer to Remark 5.2 and Notation 5.3 for explanation and to Example 5.7 for a sample calculation.

Lemma 6.4. *In the following tables, for the special families from [25, Table 1], the fourth column lists the degenerations of compact type of the inertia type and the last column lists the PEL-decomposable Newton polygons for the family under the given congruence condition on p modulo m (excluding the μ -ordinary Newton polygon when $p \equiv 1 \pmod{m}$ ⁴).*

⁴The μ -ordinary Newton polygon for $p \equiv 1 \pmod{m}$ is ordinary, and always PEL-decomposable.

<i>label</i>	<i>m</i>	<i>inertia type</i>	<i>degeneration</i>	<i>PEL-decomp. NP (congruence class)</i>
M[3]	3	(1,1,2,2)	(1,1,1)+(2,2,2)	$ss^2, p \equiv 2$
M[6]	3	(1,1,1,1,2)	(1,1,1)+(1,1,2,2)	$ord \oplus ss^2, p \equiv 1$ $ord^2 \oplus ss, p \equiv 2$ $ss^3, p \equiv 2$
M[10]	3	(1,1,1,1,1,1)	(1,1,1) + (2,1,1,1,1)	$ord^2 \oplus ss^2, p \equiv 1, 2$ $ord \oplus (1/3, 2/3), p \equiv 1$ $ss^4, p \equiv 2$
M[4]	4	(1,2,2,3)	(1,1,2)+(2,3,3)	$ss^2, p \equiv 3$
M[7]	4	(1,1,1,1)	None	None
M[8]	4	(1,1,2,2,2)	(1,2,2,3)+(1,1,2) (1,1,2) + $\text{Ind}_2^4(1,1,1,1)$	$ord \oplus ss^2, p \equiv 1$ $ord^2 \oplus ss, p \equiv 3$ $ss^3, p \equiv 3$
M[5]	6	(2,3,3,4)	(1,2,3)+(3,4,5)	$ss^2, p \equiv 5$
M[9]	6	(1,3,4,4)	(1,1,4)+(3,4,5) (1,2,3) + $\text{Ind}_3^6(2,2,2)$	$ss^3, p \equiv 5$
M[12]	6	(1,1,1,3)	(1,1,4) + (2,1,3)	$ss^4, p \equiv 5$
M[13]	6	(1,1,2,2)	(1,1,4) + $\text{Ind}_3^6(1,1,1)$	$ss^4, p \equiv 5$
M[14]	6	(2,2,2,3,3)	(2,3,1) + (5,3,2,2) (2,3,3,4) + $\text{Ind}_3^6(1,1,1)$	$ord^2 \oplus ss^2, p \equiv 1, 5$ $ss^4, p \equiv 5$

<i>label</i>	<i>m</i>	<i>a</i>	<i>degenerations</i>	<i>PEL-dec. NP, congruence on p</i>
M[11]	5	(1,3,3,3)	(1,3,1) + (4,3,3)	$ss^4, p \not\equiv 1$
M[16]	5	(2,2,2,2,2)	(2,2,2,4) + (1,2,2)	$(1/4, 3/4) \oplus ss^2, p \equiv 2, 3$ $ord^2 \oplus ss^4, p \equiv 4$ $ord^4 \oplus ss^2, p \equiv 1$ $ss^6, p \not\equiv 1$
M[17]	7	(2,4,4,4)	(4,4,6) + (1,4,2)	$ord^3 \oplus (1/3, 2/3), p \equiv 2, 4$ $ss^6, p \equiv 3, 5, 6$
M[15]	8	(2,4,5,5)	(5,5,6) + $\text{Ind}_4^8(1,2,1)$ (5,2,1)+(7,4,5)	$ord^2 \oplus ss^3, p \equiv 3$ $ord^3 \oplus ss^2, p \equiv 5$ $ss^5, p \equiv 7$
M[19]	9	(3,5,5,5)	(3,5,1) + (8,5,5)	$ord \oplus (1/3, 2/3)^2, p \equiv 4, 7$ $ss^7, p \equiv 2, 5, 8$
M[18]	10	(3,5,6,6)	(3,5,2) + $\text{Ind}_5^{10}(4,3,3)$ (6,3,1) + (9,5,6)	$ss^6, p \not\equiv 1$
M[20]	12	(4,6,7,7)	(7,7,10) + $\text{Ind}_6^{12}(1,2,3)$ (7,4,1) + (11,6,7)	$ord^3 \oplus ss^4, p \equiv 5$ $ord^4 \oplus ss^3, p \equiv 7$ $ss^7, p \equiv 11$

7. APPLICATIONS

The method of the previous sections produces numerous Newton polygons and mod p Dieudonné modules that occur for the Jacobian of a smooth curve. In Section 7.1, we collect a list of these, focusing on the ones which are new, and prove a result about curves of arbitrary large genus whose Newton polygon has slopes $1/6, 5/6$. In Section 7.2, for an infinite sequence of $g \in \mathbb{N}$ and a set of primes of density $1/2$, we produce an explicit family of smooth curves of genus g such that the multiplicity of the slope $1/2$ in the Newton polygon is at least $4\sqrt{g}$.

7.1. Newton polygons and Dieudonné modules arising from special families. The label $M[r]$ denotes the r th label in [25, Table 1].

Theorem 7.1 (Theorem 1.2). *There exists a smooth supersingular curve of genus g defined over $\overline{\mathbb{F}}_p$ for p sufficiently large in the congruence class in the following cases:*

- when $g = 5$, $p \equiv 7 \pmod{8}$, from $M[15]$;
- when $g = 6$, $p \equiv 2, 3, 4 \pmod{5}$, from $M[18]$;
- when $g = 7$, $p \equiv 2 \pmod{3}$, from $M[19, 20]$.

Proof. The results appear in Section 6 in the basic loci of families. By Proposition 5.8, for p sufficiently large in each congruence class, each of the supersingular Newton polygons above appears as the Newton polygon of the Jacobian of a smooth curve. \square

Remark 7.2. (1) In Theorem 7.1, we could also include: genus 4 when $p \equiv 2 \pmod{3}$ from $M[12 - 13]$ or $p \equiv 2, 3, 4 \pmod{5}$ from $M[11]$; and genus 6 when $p \equiv 3, 5, 6 \pmod{7}$ from $M[17]$. We do not include these cases since they already appear in [22, Theorem 1.1] for all p satisfying the congruence condition.

(2) We would like to thank Voight and Long for pointing out that the genus 6 case of Theorem 7.1 can likely also be proven (without the hypothesis that p is sufficiently large) using truncated hypergeometric functions.

We collect new examples of Newton polygons and Dieudonné modules from the μ -ordinary locus. The two Newton polygons with $*$ appear earlier in [3, 28].

Theorem 7.3. *There exists a smooth curve of genus g defined over $\overline{\mathbb{F}}_p$ with the given Newton polygon NP and mod p Dieudonné module DM in the following cases:*

genus	NP	DM	congruence on p	where
4	$*(1/4, 3/4)$	$N_{4,2}$	$2, 3 \pmod{5}$	$M[11]$
5	$ord^2 \oplus ss^3$	$L^2 \oplus N_{1,1}^3$	$3, 7 \pmod{8}$	$M[15]$
6	$(1/4, 3/4) \oplus ss^2$	$N_{4,2} \oplus N_{2,1}$	$2, 3 \pmod{5}$	$M[16, 18]$
6	$ord^2 \oplus ss^4$	$L^2 \oplus N_{1,1}^4$	$4 \pmod{5}, 6 \pmod{7},$ $or 9 \pmod{10}$	$M[16, 17]$ $M[18]$
6	$*ord^3 \oplus (1/3, 2/3)$	$L^3 \oplus N_{3,2}$	$2, 4 \pmod{7}$	$M[17]$
6	$(1/3, 2/3)^2$	Example 4.10(2)	$3, 5 \pmod{7}$	$M[17]$
7	$(1/3, 2/3)^2 \oplus ss$	Example 4.11(1)	$2, 5 \pmod{9}$	$M[19]$
7	$ord \oplus (1/3, 2/3)^2$	$L \oplus N_{3,2}^2$	$4, 7 \pmod{9}$	$M[19]$
7	$ord^4 \oplus ss^3$	$L^4 \oplus N_{1,1}^3$	$7 \pmod{12}$	$M[20]$
7	$ord^3 \oplus ss^4$	$L^3 \oplus N_{1,1}^4$	$5 \pmod{12}$	$M[20]$
7	$ord^2 \oplus ss^5$	$L^2 \oplus N_{1,1}^5$	$8 \pmod{9} or 11 \pmod{12}$	$M[19, 20]$

Proof. The result follows from Proposition 5.1 since these Newton polygons and Dieudonné modules occur in μ -ordinary cases in Section 6 under these congruence conditions. \square

Theorem 7.4. *There exists a smooth curve of genus g defined over $\overline{\mathbb{F}}_p$ with the given Newton polygon NP in the following cases:*

genus	NP	congruence on p	where
5	$(1/4, 3/4) \oplus ss$	$3 \pmod{8}$	$M[15]$
6	$(1/6, 5/6)$	$2, 4 \pmod{7}$	$M[17]$
7	$ord \oplus ss^6$	$4, 7 \pmod{9}$	$M[19]$
7	$ord^2 \oplus ss^5$	$7 \pmod{12}$	$M[20]$
7	$ord \oplus (1/4, 3/4) \oplus ss^2$	$5 \pmod{12}$	$M[20]$

Proof. These Newton polygons appear in basic loci cases from Section 6 under the given congruence conditions. By Lemma 6.4, these cases are PEL-indecomposable. Thus, by Proposition 5.6, they occur for Jacobians of smooth curves. \square

The p -divisible group $G_{1,d-1} \oplus G_{d-1,1}$ has slopes $1/d, (d-1)/d$. The next result was proven for all p , when $d = 2$ [9, Theorem 2.6]; $d = 3$ [28, Theorem 4.3]; $d = 4$ [3, Corollary 5.6]; and (under congruence conditions on p) when $d = 5$ or $d = 11$ in [22, Theorem 5.6].

Theorem 7.5. *When $d = 6$ and $p \equiv 2, 4 \pmod{7}$, for all $g \geq d$, there exists a smooth curve of genus g defined over $\overline{\mathbb{F}}_p$ whose Jacobian has p -divisible group isogenous to $(G_{1,d-1} \oplus G_{d-1,1}) \oplus (G_{0,1} \oplus G_{1,0})^{g-d}$.*

Proof. From the basic locus for $M[17]$ when $m = 7$ in Section 6, there exists a curve of genus $d = 6$ defined over $\overline{\mathbb{F}}_p$ whose p -divisible group is isogenous to $G_{1,d-1} \oplus G_{d-1,1}$. By Proposition 5.6, this curve is smooth. The Newton polygon for $G_{1,d-1} \oplus G_{d-1,1}$ is the lowest Newton polygon in dimension d with p -rank 0. Thus there is at least one component of the p -rank 0 stratum of \mathcal{M}_d whose generic geometric point has p -divisible group isogenous to $G_{1,d-1} \oplus G_{d-1,1}$. The result is then immediate from [27, Corollary 6.4]. \square

7.2. A non-special family.

Let QR (resp. QNR) be an abbreviation for quadratic residue (resp. non-residue).

Notation 7.6. Let $m > 7$ be a prime such that $m \equiv 3 \pmod{4}$. Let $p \geq m(m-7)/2$ be a prime which is a QNR modulo m . Let $N = (m-1)/2$. Let $\alpha_m = (a_1, \dots, a_N)$ be an ordering of the QRs modulo m .

The triple (m, N, α_m) is a monodromy datum because $\sum_{i=1}^N a_i \equiv 0 \pmod{m}$.

Consider the family $C = C(m, N, \alpha_m) \rightarrow U$ of curves defined as in (2.1). For $t \in U$, the genus of the curve C_t is $g_m = (m-5)(m-1)/4$ by (2.2). Recall that $Z(m, N, \alpha_m)$ is the closure in \mathcal{A}_{g_m} of the image of C under the Torelli map, and $S(m, N, \alpha_m)$ is the smallest Shimura subvariety in \mathcal{A}_{g_m} containing $Z(m, N, \alpha_m)$.

Proposition 7.7. *Consider the monodromy datum (m, N, α_m) with m, p, N, α_m as in Notation 7.6. There are explicitly computable constants $E(1), E(2) \in \mathbb{N}$, with $E(1) > E(2)$ and $E(1) + E(2) = (m-5)/2$, such that the μ -ordinary Newton polygon v_o and μ -ordinary mod p Dieudonné module N_o of the reduction modulo p of $S(m, N, \alpha_m)$ are*

$$(7.1) \quad v_o = (ord^{2E(2)} \oplus ss^{E(1)-E(2)})^{(m-1)/2},$$

$$(7.2) \quad N_o = (L^{2E(2)} \oplus N_{1,1}^{E(1)-E(2)})^{(m-1)/2}.$$

Proof. Let $n \in (\mathbb{Z}/m\mathbb{Z})^*$. Since the inertia type α_m consists of all the QRs, by (2.4), there exist $c_1, c_2 \in \mathbb{Z}^{>0}$ such that $f(\tau_n) = c_1$ when n is a QR and $f(\tau_n) = c_2$ when n is a QNR. Since $m \equiv 3 \pmod{4}$, exactly one of n and $-n$ is a QR. So $g(\tau_n) = f(\tau_n) + f(\tau_{-n}) = c_1 + c_2 = (m-5)/2$. Also $c_1 \neq c_2$ since $(m-5)/2$ is odd.

Since p is a QNR in $(\mathbb{Z}/m\mathbb{Z})^*$, each orbit \mathfrak{o} has the same number of QRs and QNRs. By §4.2, $s(\mathfrak{o}) = 2$, $E(0) = c_1 + c_2$, $E(1) = \max\{c_1, c_2\}$ and $E(2) = \min\{c_1, c_2\}$. By (4.1)-(4.2):

$$\begin{aligned}\lambda(0) &= 0, \quad m(\lambda(0)) = (\#\mathfrak{o}) \cdot E(2); \\ \lambda(1) &= 1/2, \quad m(\lambda(1)) = (\#\mathfrak{o}) \cdot (E(1) - E(2)); \text{ and} \\ \lambda(2) &= 1, \quad m(\lambda(2)) = (\#\mathfrak{o}) \cdot E(2).\end{aligned}$$

Hence, $\mu(\mathfrak{o}) = (\text{ord}^{2E(2)} \oplus \text{ss}^{E(1)-E(2)})^{\#\mathfrak{o}/2}$ and $\nu_{\mathfrak{o}} = \sum \mu(\mathfrak{o}) = (\text{ord}^{2E(2)} \oplus \text{ss}^{E(1)-E(2)})^{(m-1)/2}$. Because $E(1) - E(2) \neq 0$, the multiplicity of the slope $1/2$ in $\nu_{\mathfrak{o}}$ is at least $m-1$.

Since $s(\mathfrak{o}) = 2$ for any orbit \mathfrak{o} and $E(1) = \max\{c_1, c_2\}$ and $E(2) = \min\{c_1, c_2\}$ are independent of \mathfrak{o} , then $N_{\mathfrak{o}} = (L^{2E(2)} \oplus N_{1,1}^{E(1)-E(2)})^{(m-1)/2}$ by Example 4.9. \square

For $m > 7$, the image of the Torelli morphism is not open and dense in $S(m, N, \alpha_m)$. This makes it extremely difficult to determine the generic Newton polygon for the curve $C \rightarrow U$; In this case, a result of Bouw allows us to prove that the μ -ordinary Newton polygon $\nu_{\mathfrak{o}}$ occurs for the curves in the family.

Theorem 7.8. *Consider the monodromy datum (m, N, α_m) with m, p, N, α_m as in Notation 7.6. Consider the family $C(m, N, \alpha_m) \rightarrow U$ of curves of genus g_m defined as in (2.1). For t in an open dense subset of U , the curve C_t is smooth and has Newton polygon $\mu_{\mathfrak{o}}$ and mod p Dieudonné module $N_{\mathfrak{o}}$ as in (7.1). In particular, the Newton polygon only has slopes $0, 1/2$, and 1 and the multiplicity of the slope $1/2$ is at least $m-1 \geq 2\sqrt{g_m}$.*

Proof. Let S_p denote the reduction modulo p of $S(m, N, \alpha_m)$; it has μ -ordinary Newton polygon $\mu_{\mathfrak{o}}$ by Proposition 7.7. The Newton polygon $\mu_{\mathfrak{o}}$ is the lowest among all Newton polygons of the same p -rank. Thus in S_p , the maximal p -rank stratum agrees with the μ -ordinary Newton polygon (and Dieudonné module) stratum. In particular, the Jacobian of C_t has Newton polygon $\nu_{\mathfrak{o}}$ if and only if it has maximal possible p -rank. By [4, Theorem 6.1], for $p \geq m(N-3) = m(m-7)/2$, the maximal possible p -rank is achieved on an open dense subset of U . \square

In the first few examples, the multiplicity of the slope $1/2$ in the Newton polygon matches the lower bound $(m-1)/2$.

Example 7.9. When $m = 7$ and $a = (1, 2, 4)$, then $c_1 = 1$ and $c_2 = 0$. Thus, for $p \equiv 3, 5, 6 \pmod{7}$, the Newton polygon is ss^3 .

Example 7.10. When $m = 11$ and $a = (1, 3, 4, 5, 9)$, then $c_1 = 2$ and $c_2 = 1$. Thus, for $p \equiv 2, 6, 7, 8, 10 \pmod{11}$, the Newton polygon is $\text{ord}^{10} \oplus \text{ss}^5$.

Example 7.11. When $m = 19$ and $a = (1, 4, 5, 6, 7, 9, 11, 16, 17)$, then $c_1 = 4$ and $c_2 = 3$. Thus, for $p \equiv 2, 3, 8, 10, 12, 13, 14, 15, 18 \pmod{19}$, the Newton polygon is $\text{ord}^{54} \oplus \text{ss}^9$.

8. APPENDIX: BOUNDING THE NUMBER OF IRREDUCIBLE COMPONENTS OF THE BASIC LOCUS OF SIMPLE SHIMURA VARIETIES

In this section, we study the basic locus of a certain type of unitary Shimura variety. Under some natural restrictions on the prime p , we prove that the number of irreducible components of the basic locus of its reduction modulo p is unbounded as p goes to ∞ .

8.1. Notation. Let E be a CM field, Galois over \mathbb{Q} , and F its maximal totally real subfield. Recall from Section 3.1, that $V = \mathbb{Q}^{2g}$ has a standard symplectic form Ψ . There is an action of E on V which is compatible with Ψ . In other words, if we view V as an E -vector space, the symplectic form on V naturally induces an E/F -Hermitian form ψ on V .

Using the notation in Section 3.2, we consider the PEL-Shimura datum (H_B, h) for $B = E$. In the following, we write $G = H_E$. Note that we have an exact sequence

$$1 \rightarrow \text{Res}_{\mathbb{Q}}^E U(V, \psi) \rightarrow G \rightarrow \mathbb{G}_m \rightarrow 1$$

where $U(V, \psi)$ is the unitary group over F with respect to ψ . For example, in the notation of Section 3.2, when $E = K_m$ is a cyclotomic field, for each embedding $\tau : K_m \rightarrow \mathbb{C}$, then the signature of ψ at the real place of F induced by τ is $(f(\tau), f(\tau^*))$. Note that the reflex field of the Shimura datum (G, h) is contained in E .

For any rational prime p , we fix a prime \mathfrak{p} of E above p . In the following, with some abuse of notation, we still denote by \mathfrak{p} the corresponding primes of F and of the reflex field. We write \mathbb{A}_f for the finite adeles of \mathbb{Q} . Let $K \subset G(\mathbb{A}_f)$ be an open compact subgroup, and denote by $\text{Sh}(G, h)$ the associated Shimura variety of level K . Assume p is unramified in E , and $K = K_{\mathfrak{p}} K^p \subset G(\mathbb{Q}_p) \times G(\mathbb{A}_f^p)$ with $K_{\mathfrak{p}}$ hyperspecial.⁵ Note that this assumption holds for any prime p sufficiently large. Then, we denote by \mathcal{S} the canonical integral model of $\text{Sh}(G, h)$ at \mathfrak{p} , and write $\mathcal{S}_{\mathfrak{p}}$ for the mod \mathfrak{p} reduction of \mathcal{S} , and $\mathcal{S}_{\mathfrak{p}}(\nu_b)$ for its basic locus.

8.2. Main theorem.

Theorem 8.1. *Assume that the signature of the unitary group $U(V, \psi)$ is $(1, n - 1)$ or $(n - 1, 1)$ at one real place of F and is $(0, n)$ or $(n, 0)$ at any other real place. If $\mathfrak{p} \mid p$ is inert in E/F , we further assume that n is even. Then for any such prime \mathfrak{p} , the number of (geometrically) irreducible components of $\mathcal{S}_{\mathfrak{p}}(\nu_b)$ grows to infinity with p .*

Remark 8.2. When n is odd or when the signature has another form, the statement of Theorem 8.1 does not hold in general. For example, when the center Z_G of G is connected, Xiao and Zhu show that if the dimension of the basic locus is half the dimension of a Hodge type Shimura variety, then the number of its irreducible components is the same for all unramified primes [39, Lemma 1.1.3, Theorem 1.1.4 (1), Remark 1.1.5 (2), Proposition 7.4.2]. This dimension requirement is satisfied for any unitary Shimura variety with the signature as in Theorem 8.1 and n odd at inert prime \mathfrak{p} . For more examples, see [39, Remark 4.2.11].

Related material is in [40].

Corollary 8.3. *The statement of Theorem 8.1 holds true for any connected component of a product of finitely many unitary Shimura varieties satisfying the assumptions of Theorem 8.1.*

⁵If p is unramified in E , then $G_{\mathbb{Q}_p}$ admits a smooth reductive model \mathcal{G} over \mathbb{Z}_p , and $K_{\mathfrak{p}} = \mathcal{G}(\mathbb{Z}_p)$ is hyperspecial.

Definition 8.4. The group of self-isogenies and certain open compact subgroups. Fix a point x in $\mathcal{S}_p(\nu_b)$, and let A_x denote the associated abelian variety, endowed with the additional PEL-structure. We write $I = I_x$ for the group of quasi-isogenies of A_x which are compatible with the E -action and the \mathbb{Q} -polarization. Then I is an inner form of the algebraic reductive group G . Furthermore, $I(\mathbb{Q}_\ell) = G(\mathbb{Q}_\ell)$ at any finite prime $\ell \neq p$, and $I(\mathbb{R})/\mathbb{R}^*$ is compact (due to the positivity of the Rosati involution).

In the following, we construct an open compact subgroup C_p of $I(\mathbb{Q}_p)$ such that the mass of $C_p K^p$ gives a lower bound of the number of irreducible components of $\mathcal{S}_p(\nu_b)$.

- (1) Assume \mathfrak{p} is split in E/F . Then, $I(\mathbb{Q}_p)$ decomposes as $I(\mathbb{Q}_p) = \mathbb{Q}_p^\times \prod_{v|p} I_v$,⁶ where v runs through all places of F above p . Furthermore, $I_v \cong \mathrm{GL}_n(F_v)$ for $v \neq \mathfrak{p}$, and $I_{\mathfrak{p}}$ is isomorphic to $D_{F_{\mathfrak{p}},n}^\times$ the division algebra over $F_{\mathfrak{p}}$ with invariant $1/n$. We define C_p as the maximal compact subgroup of $I(\mathbb{Q}_p)$ given by

$$C_p = \mathbb{Z}_p^\times \mathcal{O}_{D_{F_{\mathfrak{p}},n}}^\times \prod_{v|p, v \neq \mathfrak{p}} \mathrm{GL}_n(\mathcal{O}_{F_v}),$$

where $\mathcal{O}_{D_{F_{\mathfrak{p}},n}}^\times \subset D_{F_{\mathfrak{p}},n}^\times$ is the subgroup of elements with norm in $\mathcal{O}_{F_{\mathfrak{p}}}^\times$. (For more details, see for example [14, Chapter II.1]).

- (2) Assume \mathfrak{p} is inert in E/F . We generalize the discussion in [36, §2], where Vollaard treats the case $F = \mathbb{Q}$. Here, we follow [23] and recall that n is even in this case.

Let V^\bullet be the n -dimensional E/F -Hermitian space such that

- $V^\bullet \otimes_F \mathbb{A}_{F,f}^p \cong V \otimes_F \mathbb{A}_{F,f}^p$ as Hermitian spaces; (we fix such an isomorphism);
- V^\bullet has signature $(n, 0)$ or $(0, n)$ at all archimedean places (more precisely, we only change the signature by 1 at the one indefinite place of V),
- $V^\bullet \otimes_F F_{\mathfrak{p}}$ is a ramified Hermitian space over $E_{\mathfrak{p}}/F_{\mathfrak{p}}$.

Fix an $\mathcal{O}_E \otimes \mathbb{Z}_p$ -lattice $\Lambda_p^\bullet \subset V^\bullet \otimes_{\mathbb{Q}} \mathbb{Q}_p$, such that the dual $\Lambda_p^{\bullet, \vee}$ of Λ_p^\bullet with respect to the Hermitian form satisfies $\Lambda_p^\bullet \subset \Lambda_p^{\bullet, \vee}$ and $\Lambda_p^{\bullet, \vee} / \Lambda_p^\bullet \simeq \mathcal{O}_E / \mathfrak{p}$ (such lattice exists due to the above assumption on V^\bullet). Then I is the unitary similitude group of V^\bullet ⁷ and we define $C_p \subset I(\mathbb{Q}_p)$ to be the stabilizer group of Λ_p^\bullet .

Proposition 8.5. *The number of irreducible components of $\mathcal{S}(\nu_b)_{\mathfrak{p}}$ is bounded below by the mass of $C_p K^p$,*

$$m(C_p K^p) := \#I(\mathbb{Q}) \backslash I(\mathbb{A}_f) / C_p K^p.$$

Proof. By Rapoport and Zink's p -adic uniformization theorem [30, Theorem 6.30], to prove the statement it suffices to show that the irreducible components (of a subset) of the Rapoport–Zink space RZ are indexed by $I(\mathbb{Q}_p)/C_p$.

Assume \mathfrak{p} is split in E/F . By Theorem 4.6, there are n possible Newton polygons on \mathcal{S}_p and hence by [13, Theorem 1.1], the basic locus is 0-dimensional. Note that the group C_p is the stabilizer of the Dieudonné lattice of A_x . Hence $I(\mathbb{Q}_p)$ acts on RZ with stabilizer C_p and the number of irreducible components is bounded below by $I(\mathbb{Q}_p)/C_p$.

⁶As a convention in the appendix, if we write $G = H_1 H_2$, it means that H_1, H_2 are subgroups of G and every element g in G can be written as $h_1 h_2$ for $h_i \in H_i, i = 1, 2$ with *no* assumption that such decomposition is unique. For instance, we do not assume that $H_1 \cap H_2 = \{1\}$.

⁷Step (2) of the inert case of the proof of Proposition 8.5 contains a proof for this well-known fact.

Assume \mathfrak{p} is inert in E/F . For $F = \mathbb{Q}$, the statement is [37, Theorem 5.2 (2), Prop. 6.3]. For F totally split at p , the statement is proved in [23]. Here, we sketch a proof that generalizes [37] and [23]. The proof is in three steps.

- (1) Construct a 0-dimensional Shimura variety $\text{Sh}(GU(V^\bullet))$ of level $C_p K^p$, parametrizing abelian varieties A^\bullet of dimension $[F : \mathbb{Q}]n$ with \mathcal{O}_E -action and polarization λ^\bullet satisfying Kottwitz's determinant condition. Here we assume that the polarization λ^\bullet admits the same polarization type as in $\text{Sh}(G, h)$ outside \mathfrak{p} ; at \mathfrak{p} we assume that $(\text{Ker } \lambda^\bullet)[p^\infty] = (\text{Ker } \lambda^\bullet)[\mathfrak{p}]$ and that the latter is a finite flat group scheme of order $\#\mathcal{O}_E/\mathfrak{p}$. To describe the Kottwitz's condition, let $\Phi \subset \text{Hom}(E, \mathbb{C})$ be the subset of τ such that the signature $(f(\tau), f(\tau^*))$ of V^\bullet is $(n, 0)$. Note that $\Phi \sqcup \Phi^* = \text{Hom}(E, \mathbb{C})$. Kottwitz's determinant condition says that the characteristic polynomial of the action $b \in \mathcal{O}_E$ on $\text{Lie}(A^\bullet)$ is given by $\prod_{\tau \in \Phi} (x - \tau(b))^n$.
- (2) We will construct a Deligne–Lusztig variety DL and, for a fixed A^\bullet as in (1), a family of abelian varieties A in $\mathcal{S}_{\mathfrak{p}}(\nu_b)$ together with a universal isogeny from A^\bullet parametrized by DL^{perf} , the perfection of DL .

Let τ_0 denote the unique indefinite real place for $U(V)$ and let f denote the inertia degree of \mathfrak{p} in F . Consider the σ -orbit $\mathfrak{o}_{\tau_0} = \{\tau_0, \sigma\tau_0, \dots, \sigma^f\tau_0 = \tau_0^*, \sigma\tau_0^*, \dots, \sigma^{f-1}\tau_0^*\}$. For $\tau \in \mathfrak{o}_{\tau_0}$, define $\tilde{F}^f = F_f \circ \dots \circ F_1$, where $F_i : H_1^{\text{dR}}(A^\bullet)_{\sigma^{i-1}\tau_0} \rightarrow H_1^{\text{dR}}(A^\bullet)_{\sigma^i\tau_0}$ is equal to F if $\sigma^{i-1}\tau \in \Phi$, and to V^{-1} otherwise.⁸ Then, there exists a submodule $M \subset H_1^{\text{dR}}(A^\bullet)_{\tau_0} \otimes \bar{\mathbb{F}}_p$ of rank $n/2 + 1$ satisfying the condition

$$\tilde{F}^f((M^{(p^f)}))^\perp \subset M,$$

where $^\perp$ is taken with respect to the pairing

$$\langle -, - \rangle^\bullet : H_1^{\text{dR}}(A^\bullet)_{\tau_0} \times H_1^{\text{dR}}(A^\bullet)_{\tau_0^*} \rightarrow \bar{\mathbb{F}}_p.^9$$

The Deligne–Lusztig variety DL is the moduli of all submodules M satisfying the above conditions.

To construct A over DL^{perf} , we use covariant Dieudonné theory and by a theorem of Gabber (see for instance [20, Theorem D]), we only need realize the Dieudonné module of A as a sub-Dieudonné module of $D(A^\bullet)$.

We define $D(A)_\tau \subset D(A^\bullet)_\tau$ as follows.

- (a) For $\tau \notin \mathfrak{o}_{\tau_0}$: we define $D(A)_\tau = D(A^\bullet)_\tau$ if $\tau \notin \Phi$, and $D(A)_\tau = pD(A^\bullet)_\tau$ if $\tau \in \Phi$.
- (b) For $\tau \in \mathfrak{o}_{\tau_0}$: let \tilde{M} be the preimage of M in $D(A^\bullet)_{\tau_0}$ under the mod p map, and \tilde{M}^* the preimage of M^\perp in $D(A^\bullet)_{\tau_0^*}$. We define $D(A)_{\tau_0^*} = F(\tilde{M})$ and $D(A)_{\tau_0} = V^{-1}(\tilde{M}^*)$. For $\tau = \sigma^i\tau_0$, where $1 \leq i \leq f$ (resp. $f+1 \leq i \leq 2f$), we define $D(A)_\tau = \tilde{V}D(A)_{\sigma\tau}$ inductively from $\tau_0^c = \sigma^f\tau_0$ (resp. $\tau_0 = \tau_0^{2f}$), where $\tilde{V} = V$ if $\tau \notin \Phi$ and $\tilde{V} = p^{-1}V$ otherwise.

⁸We have that $V : H_1^{\text{dR}}(A^\bullet)_{\sigma\tau} \rightarrow H_1^{\text{dR}}(A^\bullet)_\tau$ is invertible if and only if $\tau \notin \Phi$.

⁹For the totally split case, see [23]. To prove that such a submodule M exists, one can argue by induction on n as in [23].

Note that by definition the submodule $D(A)$ of $D(A^\bullet)$ is invariant under F and V . Hence, it is a sub-Dieudonné module.¹⁰ Furthermore, the abelian variety A inherits a polarization and additional PEL-structures from those of A^\bullet , and satisfies Kottwitz's determinant condition.

- (3) Show that step (2) constructs an irreducible component of $\mathcal{S}_p(\nu_b)$, or similarly of RZ . Indeed, this can be proven by showing that the image of DL has the correct dimension. On one hand, by the same argument as in the proof of [36, Proposition 2.13] (replacing the p -Frobenius by the q -Frobenius), the dimension of the Deligne–Lusztig variety DL in step (2) is $n/2 - 1$. On the other hand, by Theorem 4.6, the μ -ordinary Newton polygon $\nu_o(\sigma_{\tau_0})$ has break points

$$(2f, f), (2f(n-1), (n-1)f-1), \text{ and } (2fn, fn).$$

Hence all possible non-supersingular Newton polygons are in one-to-one correspondence with integer points with abscissa $2ft$, for some $t \in \mathbb{Z} \cap [1, n/2]$. In particular, by [13, Theorem 1.1] (combined with Theorem 4.6) the basic locus (i.e., the supersingular locus) has codim $n/2$, and thus dimension $n/2 - 1$.

To conclude, we observe that C_p is the stabilizer of this irreducible component under the action of $I(\mathbb{Q}_p)$ on RZ arising from its natural action on the associated Dieudonné modules. \square

Proof of Theorem 8.1. By Proposition 8.5, the theorem follows once we provide an asymptotic lower bound for $m(C_p K^p)$ which grows to infinity with p . By [11, Proposition 2.13],¹¹

$$m(C_p K^p) = c \cdot \lambda_S$$

where $\lambda_S = \prod_{p \in S} \lambda_p$, for λ_p an explicit local factor at p and S the set of finite places v of \mathbb{Q} where I_v is not isomorphic to G_v , and $c = 2^{-(n[F:\mathbb{Q}]+1)} c' \cdot L(M_I) \cdot \tau(I)$, where

- M_I is a motive of Artin–Tate type attached to I by Gross,
- $L(M_I)$ is the value of the L -function of M_I at 0,
- $\tau(I)$ is the Tamagawa number of I , and
- c' depends only on the non-hyperspecial piece of the level K .

Note that in our case $S = \{p\}$ (see Section 8.4). We claim that c is independent of p . Indeed, the constant $L(M_I)$ only depends on the quasi-split inner form of I over \mathbb{Q} , which is independent of p ; by [18], $\tau(I)$ is independent of p because the center of the neutral connected component of the Langlands dual group of I is independent of p ; since K is hyperspecial at p , then c' is independent of p .

Hence, to conclude, it suffices to prove that the local factor λ_p is unbounded as p grows to infinity. In [11, Formula (2.12)], the local factor λ_p is explicitly computed as

$$\lambda_p = \frac{p^{-N(\overline{G}_p)} \cdot \#\overline{G}_p(\mathbb{F}_p)}{p^{-N(\overline{I}_p)} \cdot \#\overline{I}_p(\mathbb{F}_p)},$$

¹⁰One may use the non-emptiness of basic locus due to Wedhorn and Viehmann to prove the existence of such M . The argument goes the reverse way: let A be the abelian variety with $\nu(A) = \nu_b$. As in [36], one can construct an F -invariant lattice from $D(A)$ and this lattice recovers Λ_p^\bullet .

¹¹A mass formula for the stabilizer of a maximal lattice in a quadratic or Hermitian space over a totally real field was first proved by Shimura in [32, 33, 34].

where, for $H = G, I$, the group \overline{H}_p denotes the maximal reductive quotient of the special fiber of H_p , and $N(\overline{H}_p)$ denotes the number of positive roots of \overline{H}_p over $\overline{\mathbb{F}}_p$. Note that the integral structure of I_p is given by C_p .

Assume p is split in E/F . Then, G and I only differ at \mathfrak{p} . More precisely, $G_{\mathbb{Z}_p} = \mathbb{G}_m \prod_{v|p} G_v$ and $G_v \cong I_v$ for $v \neq \mathfrak{p}$. At \mathfrak{p} , the group $C_p = \mathcal{O}_{D_{F_p, n}}^\times \subset I_p$ is Iwahori and \overline{I}_p modulo \mathbb{G}_m is a totally non-split torus of rank $n - 1$. More precisely, \overline{I}_p is the multiplicative group of the degree n extension of $\mathcal{O}_F/\mathfrak{p}$. Hence, for $q = \#\mathcal{O}_F/\mathfrak{p}$, we have

$$\lambda_p = \frac{q^{-N(\mathrm{GL}_n)} \#\mathrm{GL}_n(\mathbb{F}_q)}{\#\overline{I}_p(\mathbb{F}_q)} = \frac{q^{(1-n)n/2} \prod_{i=1}^n (q^n - q^{i-1})}{q^n - 1} = \prod_{i=2}^n (q^{n-i+1} - 1).$$

Assume \mathfrak{p} is inert in E/F . In Lemma 8.6 below, we verify that the $\mathcal{O}_E \otimes \mathbb{Z}_p$ -lattice Λ_p^\bullet is maximal. Hence, the computation in [11, §3, Table 2] applies for n even. For $q = \#\mathcal{O}_F/\mathfrak{p}$, we have

$$\lambda_p = (q^n - 1)/(q + 1).$$

□

Lemma 8.6. *Maintaining the same assumptions as in Theorem 8.1, suppose \mathfrak{p} is inert in E/F and n is even. Then the $\mathcal{O}_E \otimes \mathbb{Z}_p$ -lattice Λ_p^\bullet is maximal. More precisely, let \mathbb{H} be the split rank 2 Hermitian space over E_v with a standard basis $\{e, f\}$ and let Δ be the maximal lattice $\mathcal{O}_{E,v}e \oplus \mathcal{O}_{E,v}f$.*

- (1) *if $v \neq \mathfrak{p}$, then $\Lambda_v^\bullet \cong \Delta^n$ as Hermitian lattices.*
- (2) *if $v = \mathfrak{p}$, then, as Hermitian lattices, $\Lambda_v^\bullet \cong \Delta^{n-1} \oplus \mathcal{O}_D$, where \mathcal{O}_D is the maximal order of the unique quaternion algebra over F_v .¹²*

Proof. Let Λ be a maximal lattice containing Λ_p^\bullet . We use integral Witt decomposition for the maximal lattice Λ to check that Λ itself satisfies the duality condition $\Lambda^\vee/\Lambda \simeq \mathcal{O}_E/\mathfrak{p}$. Hence, $\Lambda = \Lambda_p^\bullet$ and it is maximal. (For definition and results on the integral Witt decomposition, we refer to [32, Chapter 1]. Also, note that since p is unramified in E/Q , the inverse of the different ideal \mathfrak{d}^{-1} is relatively prime to p .) □

REFERENCES

1. Jeffrey D. Achter and Rachel Pries, *Monodromy of the p -rank strata of the moduli space of curves*, Int. Math. Res. Not. IMRN (2008), no. 15, Art. ID rnn053, 25. MR 2438069
2. ———, *The p -rank strata of the moduli space of hyperelliptic curves*, Adv. Math. **227** (2011), no. 5, 1846–1872. MR 2803789
3. ———, *Generic Newton polygons for curves of given p -rank*, Algebraic curves and finite fields, Radon Ser. Comput. Appl. Math., vol. 16, De Gruyter, Berlin, 2014, pp. 1–21. MR 3287680
4. Irene I. Bouw, *The p -rank of ramified covers of curves*, Compositio Math. **126** (2001), no. 3, 295–322. MR 1834740
5. A. J. de Jong and F. Oort, *Purity of the stratification by Newton polygons*, J. Amer. Math. Soc. **13** (2000), no. 1, 209–241. MR 1703336
6. P. Deligne and G. D. Mostow, *Monodromy of hypergeometric functions and nonlattice integral monodromy*, Inst. Hautes Études Sci. Publ. Math. (1986), no. 63, 5–89. MR 849651

¹²More precisely, we equip D with the following Hermitian form. As an E_v vector space, write $D = E_v + zE_v$ such that $z^2 = \alpha := \mathrm{disc}(V^\bullet \otimes_E E_v)$ and $xz = zx^c$ for $x \in E_v$. Then the Hermitian form is given by $\langle x_1 + zx_2, y_1 + zy_2 \rangle = x_1y_1^c - \alpha x_2y_2^c$

7. E. Eischen and E. Mantovan, *p-adic families of automorphic forms in the μ -ordinary setting*, preprint, available on arXiv:1710.01864.
8. Torsten Ekedahl, *Boundary behaviour of Hurwitz schemes*, The moduli space of curves (Texel Island, 1994), Progr. Math., vol. 129, Birkhäuser Boston, Boston, MA, 1995, pp. 173–198. MR 1363057
9. Carel Faber and Gerard van der Geer, *Complete subvarieties of moduli spaces and the Prym map*, J. Reine Angew. Math. **573** (2004), 117–137. MR 2084584
10. William Fulton, *Hurwitz schemes and irreducibility of moduli of algebraic curves*, Ann. of Math. (2) **90** (1969), 542–575. MR 0260752
11. Wee Teck Gan, Jonathan P. Hanke, and Jiu-Kang Yu, *On an exact mass formula of Shimura*, Duke Math. J. **107** (2001), no. 1, 103–133. MR 1815252
12. Darren Glass and Rachel Pries, *Hyperelliptic curves with prescribed p -torsion*, Manuscripta Math. **117** (2005), no. 3, 299–317. MR 2154252
13. Paul Hamacher, *The almost product structure of Newton strata in the deformation space of a Barsotti-Tate group with crystalline Tate tensors*, Math. Z. **287** (2017), no. 3-4, 1255–1277. MR 3719535
14. Michael Harris and Richard Taylor, *The geometry and cohomology of some simple Shimura varieties*, Annals of Mathematics Studies, vol. 151, Princeton University Press, Princeton, NJ, 2001, With an appendix by Vladimir G. Berkovich. MR 1876802
15. X. He and M. Rapoport, *Stratifications in the reduction of Shimura varieties*, Manuscripta Math. **152** (2017), no. 3-4, 317–343. MR 3608295
16. R. Kottwitz and M. Rapoport, *On the existence of F -crystals*, Comment. Math. Helv. **78** (2003), no. 1, 153–184. MR 1966756
17. Robert E. Kottwitz, *Isocrystals with additional structure*, Compositio Math. **56** (1985), no. 2, 201–220. MR 809866
18. ———, *Tamagawa numbers*, Ann. of Math. (2) **127** (1988), no. 3, 629–646. MR 942522
19. ———, *Isocrystals with additional structure. II*, Compositio Math. **109** (1997), no. 3, 255–339. MR 1485921
20. Eike Lau, *Smoothness of the truncated display functor*, J. Amer. Math. Soc. **26** (2013), no. 1, 129–165. MR 2983008
21. Wanlin Li, Elena Mantovan, Rachel Pries, and Yunqing Tang, *Newton polygon stratification of the Torelli locus in PEL-type Shimura varieties*, Available on arXiv:1811.00604.
22. ———, *Newton polygons of cyclic covers of the projective line branched at three points*, Research Directions in Number Theory: Women in Numbers IV. To appear, available on arXiv:1805.04598.
23. Yifeng Liu, Yichao Tian, Liang Xiao, Wei Zhang, and Xinwen Zhu, *On the Beilinson–Bloch–Kato conjecture for Rankin–Selberg*, in preparation.
24. Ben Moonen, *Serre-Tate theory for moduli spaces of PEL type*, Ann. Sci. École Norm. Sup. (4) **37** (2004), no. 2, 223–269. MR 2061781
25. ———, *Special subvarieties arising from families of cyclic covers of the projective line*, Doc. Math. **15** (2010), 793–819. MR 2735989
26. Frans Oort, *Abelian varieties isogenous to a Jacobian in problems from the Workshop on Automorphisms of Curves*, Rend. Sem. Mat. Univ. Padova **113** (2005), 129–177. MR 2168985 (2006d:14027)
27. Rachel Pries, *Current results on Newton polygons of curves*, to appear in Questions in Arithmetic Algebraic Geometry, Advanced Lectures in Mathematics, Chapter 6.
28. ———, *The p -torsion of curves with large p -rank*, Int. J. Number Theory **5** (2009), no. 6, 1103–1116. MR 2569747
29. M. Rapoport and M. Richartz, *On the classification and specialization of F -isocrystals with additional structure*, Compositio Math. **103** (1996), no. 2, 153–181. MR 1411570
30. M. Rapoport and Th. Zink, *Period spaces for p -divisible groups*, Annals of Mathematics Studies, vol. 141, Princeton University Press, Princeton, NJ, 1996. MR 1393439
31. Xu Shen and Chao Zhang, *Stratification in good reductions of Shimura varieties of abelian type*, preprint, available on arXiv:1707.00439.
32. Goro Shimura, *Euler products and Eisenstein series*, CBMS Regional Conference Series in Mathematics, vol. 93, Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1997. MR 1450866
33. ———, *An exact mass formula for orthogonal groups*, Duke Math. J. **97** (1999), no. 1, 1–66. MR 1681092

34. ———, *Some exact formulas on quaternion unitary groups*, J. Reine Angew. Math. **509** (1999), 67–102. MR 1679167
35. Eva Viehmann and Torsten Wedhorn, *Ekedahl-Oort and Newton strata for Shimura varieties of PEL type*, Math. Ann. **356** (2013), no. 4, 1493–1550. MR 3072810
36. Inken Vollaard, *The supersingular locus of the Shimura variety for $GU(1, s)$* , Canad. J. Math. **62** (2010), no. 3, 668–720. MR 2666394
37. Inken Vollaard and Torsten Wedhorn, *The supersingular locus of the Shimura variety of $GU(1, n - 1)$ II*, Invent. Math. **184** (2011), no. 3, 591–627. MR 2800696
38. Stefan Wewers, *Construction of Hurwitz spaces*, Dissertation, 1998.
39. Liang Xiao and Xinwen Zhu, *Cycles on Shimura varieties via geometric Satake*, preprint, available on arXiv:1707.05700, 2017.
40. Chia-Fu Yu, *Simple mass formulas on Shimura varieties of PEL-type*, Forum Math. **22** (2010), no. 3, 565–582. MR 2652712

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WI 53706, USA

E-mail address: wanlin@math.wisc.edu

DEPARTMENT OF MATHEMATICS, CALIFORNIA INSTITUTE OF TECHNOLOGY PASADENA, CA 91125, USA

E-mail address: mantovan@caltech.edu

DEPARTMENT OF MATHEMATICS, COLORADO STATE UNIVERSITY, FORT COLLINS, CO 80523, USA

E-mail address: pries@math.colostate.edu

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NJ 08540, USA

E-mail address: yunqingt@math.princeton.edu