

NEWTON POLYGON STRATIFICATION OF THE TORELLI LOCUS IN PEL-TYPE SHIMURA VARIETIES

WANLIN LI, ELENA MANTOVAN, RACHEL PRIES, AND YUNQING TANG

ABSTRACT. We study the intersection of the Torelli locus with the Newton polygon stratification of the modulo p reduction of certain PEL-type Shimura varieties. We develop a clutching method to show that the intersection of the open Torelli locus with some Newton polygon strata is non-empty. This allows us to give a positive answer, under some compatibility conditions, to a question of Oort about smooth curves in characteristic p whose Newton polygons are an amalgamate sum.

As an application, we produce infinitely many new examples of Newton polygons that occur for smooth curves that are cyclic covers of the projective line. Most of these arise in inductive systems which demonstrate unlikely intersections of the open Torelli locus with the Newton polygon stratification in Siegel modular varieties. In addition, for the twenty special PEL-type Shimura varieties found in Moonen's work, we prove that all Newton polygon strata intersect the open Torelli locus (if $p \gg 0$ in the supersingular cases).

Keywords: curve, cyclic cover, Jacobian, abelian variety, moduli space, Shimura variety, PEL-type, reduction, Frobenius, supersingular, Newton polygon, p -rank, Dieudonné module, p -divisible group.

MSC10 classifications: primary 11G18, 11G20, 11M38, 14G10, 14G35; secondary 11G10, 14H10, 14H30, 14H40, 14K10

1. INTRODUCTION

1.1. Overview. Consider the moduli space \mathcal{A}_g of principally polarized abelian varieties of dimension g in characteristic $p > 0$. It can be stratified by Newton polygon. In general, it is unknown how these strata intersect the open Torelli locus \mathcal{T}_g° , which is the image of the moduli space \mathcal{M}_g of smooth genus g curves under the Torelli morphism. For a symmetric Newton polygon ν of height $2g$, in most cases it is not known whether the stratum $\mathcal{A}_g[\nu]$ intersects \mathcal{T}_g° or, equivalently, whether there exists a smooth curve of genus g in characteristic p whose Jacobian has Newton polygon ν . This question is answered only when ν is close to ordinary, meaning that the codimension of $\mathcal{A}_g[\nu]$ in \mathcal{A}_g is small.

In this paper, we develop a framework to study Newton polygons of Jacobians of μ_m -covers of the projective line \mathbb{P}^1 for an integer $m \geq 2$. We study the Newton polygon stratification on the Hurwitz spaces which represent such covers. Using clutching morphisms, we produce singular curves with prescribed Newton polygons. Under an *admissible* condition Definition 3.3, these singular curves can be deformed to smooth curves which are μ_m -covers of \mathbb{P}^1 ; we prove this can be done *without changing the Newton polygon* under a *balanced* condition Definition 4.2, or a further *compatible* condition Definition 6.3.

We then find systems of Hurwitz spaces of μ_m -covers of \mathbb{P}^1 for which the admissible, balanced, and compatible conditions, together with an expected codimension condition on the Newton polygon strata, can be verified inductively. The base cases we use involve cyclic covers of \mathbb{P}^1 branched at 3 points or the 20 special families found by Moonen [25].

As an application, we find numerous infinite sequences of unusual Newton polygons which occur for Jacobians of smooth curves. Most of these arise in an *unlikely intersection* of the open Torelli locus \mathcal{T}_g° with the Newton polygon strata of \mathcal{A}_g Definition 8.2.

In essence, our strategy is to generalize earlier techniques for studying the intersection of the Torelli locus with the Newton polygon stratification from the context of the system of moduli spaces \mathcal{A}_g to the context of appropriate inductive systems of PEL-type Shimura varieties. Hurwitz spaces of μ_m -covers of \mathbb{P}^1 determine unitary Shimura varieties associated with the group algebra of μ_m , as constructed by Deligne–Mostow [8]. By work of Kottwitz [16, 18], Viehmann–Wedhorn [30], and Hamacher [14], the Newton polygon stratification of the modulo p reduction of these Shimura varieties is well understood in terms of their signature types and the congruence class of p modulo m . It is difficult to understand the intersection of the Newton polygon stratum $Sh[v]$ with the open Torelli locus because the codimension of the latter in Sh grows with g ; our main results verify that this intersection is non-trivial when v is close to μ -ordinary, for infinitely many Shimura varieties Sh of PEL-type.

1.2. Comparison with earlier results. In 2005, Oort proposed the following conjecture.

Conjecture 1.1. ([27, Conjecture 8.5.7]) *For $i = 1, 2$, let $g_i \in \mathbb{Z}_{\geq 1}$ and let v_i be a symmetric Newton polygon appearing on $\mathcal{T}_{g_i}^\circ$. Write $g = g_1 + g_2$. Let v be the amalgamate sum of v_1 and v_2 . Then v appears on \mathcal{T}_g° .*

It is not clear whether Oort’s conjecture is true in complete generality. Theorems 4.5 and 6.11 show that Oort’s conjecture has an affirmative answer in many cases.

The results in Section 4 can be viewed as a generalization of Bouw’s work [5] about the intersection of \mathcal{T}_g° with the stratum of maximal p -rank in a PEL-type Shimura variety. For most families of μ_m -covers and most congruence classes of p modulo m , the maximal p -rank does not determine the Newton polygon.

Clutching morphisms were also used to study the intersection of \mathcal{T}_g° with the p -rank stratification of \mathcal{A}_g in Faber and Van der Geer’s work [11, Theorem 2.3]; also [2], [13], [3].

The results in Section 6 generalize Pries’ work [28, Theorem 6.4], which states that if a Newton polygon v occurs on \mathcal{M}_g with the expected codimension, then the Newton polygon $v \oplus (0, 1)^n$ occurs on \mathcal{M}_{g+n} with the expected codimension for $n \in \mathbb{Z}_{\geq 1}$. However, the expected codimension condition is difficult to verify for most Newton polygons v .

1.3. Outline of main results. In Section 2, we review key background about Hurwitz spaces, PEL-type Shimura varieties, and Newton polygon stratifications. In Section 3, we analyze the image of a clutching morphism κ on a pair of μ_m -covers of \mathbb{P}^1 .

In Section 4, we study whether the open Torelli locus \mathcal{T}_g° intersects the μ -ordinary Newton polygon stratum, see Definition 2.5, inside the Shimura variety S . The first main result Theorem 4.5 provides a method to leverage information about this question from lower to higher genus. Under a balanced condition on the signatures Definition 4.2, we can determine the μ -ordinary Newton polygon as the Shimura variety varies in the clutching system (Proposition 4.4, which we prove in Section 5).

The most ground-breaking results in the paper are in Section 6, where we study the intersection of the open Torelli locus \mathcal{T}_g° with the non μ -ordinary Newton polygon strata inside the Shimura variety S . Theorem 6.11 also provides a method to leverage information from lower to higher genus. Under an additional compatibility condition on the

signatures Definition 6.3, we can control the codimension of the Newton polygon strata as the Shimura variety varies in the clutching system (Proposition 6.8).

In Sections 4.3 and 6.4, we find situations where Theorems 4.5 and 6.11 can be implemented recursively, infinitely many times, which yields smooth curves with arbitrarily large genera and prescribed (unusual) Newton polygons. We do this by constructing suitable *infinite clutching systems* of PEL-type Shimura varieties which satisfy the admissible, balanced, (and compatible) conditions at every level. For example, we prove:

Theorem 1.2 (Special case of Corollary 4.9). *Let $\gamma = (m, N, a)$ be a monodromy datum as in Definition 2.1. Let p be a prime such that $p \nmid m$. Let u be the μ -ordinary Newton polygon associated to γ as in Definition 2.5. Suppose there exists a cyclic cover of \mathbb{P}^1 defined over $\overline{\mathbb{F}}_p$ with monodromy datum γ and Newton polygon u .¹ Then, for any $n \in \mathbb{Z}_{\geq 1}$, there exists a smooth curve over $\overline{\mathbb{F}}_p$ with Newton polygon $u^n \oplus (0, 1)^{(m-1)(n-1)}$.²*

For a symmetric Newton polygon ν of height $2g$, we say that the open Torelli locus has an *unlikely intersection* with the Newton polygon stratum $\mathcal{A}_g[\nu]$ in \mathcal{A}_g if there exists a smooth curve of genus g with Newton polygon ν and if $\dim(\mathcal{M}_g) < \text{codim}(\mathcal{A}_g[\nu], \mathcal{A}_g)$, Definition 8.2. In Section 8, we study the asymptotic of $\text{codim}(\mathcal{A}_g[\nu], \mathcal{A}_g)$ for the Newton polygons ν appearing in Sections 4.3 and 6.4, and verify that most of our inductive systems produce unlikely intersections once g is sufficiently large, for most congruence classes of p modulo m .

1.4. Applications. In Corollary 7.2 in Section 7, we prove that all the Newton polygons for the Shimura varieties associated to the families in [25, Table 1] occur for smooth curves in the family.

In Section 9, we construct explicit infinite sequences of Newton polygons that occur at odd primes for smooth curves which demonstrate unlikely intersections. For example, by Theorem 1.2 applied to $\gamma = (m, 3, (1, 1, m - 2))$, we prove:

Application 1.3. (Corollary 9.2) Let $m \in \mathbb{Z}_{>1}$ be odd and $h = (m - 1)/2$. Let p be a prime, $p \nmid 2m$, such that the order f of p in $(\mathbb{Z}/m\mathbb{Z})^*$ is even and $p^{f/2} \equiv -1 \pmod{m}$. For $n \in \mathbb{Z}_{\geq 1}$, there exists a μ_m -cover $C \rightarrow \mathbb{P}^1$ defined over $\overline{\mathbb{F}}_p$ where C is a smooth curve of genus $g = h(3n - 2)$ with Newton polygon $\nu = (1/2, 1/2)^{hn} \oplus (0, 1)^{2h(n-1)}$.³ If $n \geq 34/h$, then $\text{Jac}(C)$ lies in the unlikely intersection $\mathcal{T}_g^\circ \cap \mathcal{A}_g[\nu]$.

In Corollary 9.4, we apply Application 1.3 when $m = 3$ to verify, for $p \equiv 2 \pmod{3}$ and $g \in \mathbb{Z}_{\geq 1}$, there exists a smooth curve of genus g defined over $\overline{\mathbb{F}}_p$ whose Newton polygon only has slopes $\{0, 1/2, 1\}$ and the multiplicity of slope $1/2$ is at least $2\lfloor g/3 \rfloor$. To our knowledge, this is the first time for any odd prime p that a sequence of smooth curves has been produced for every $g \in \mathbb{Z}_{\geq 1}$ such that the multiplicity of the slope $1/2$ in the Newton polygon grows linearly in g .

¹See Proposition 4.6 for when this condition is satisfied.

²The slopes of this Newton polygon are the slopes of u (with multiplicity scaled by n) and 0 and 1 each with multiplicity $(m - 1)(n - 1)$.

³Its slopes are $1/2$ with multiplicity $2hn$ and 0 and 1 each with multiplicity $2h(n - 1)$.

Acknowledgements. Pries was partially supported by NSF grant DMS-15-02227. Tang is partially supported by NSF grant DMS-1801237. We thank the Banff International Research Station for hosting *Women in Numbers 4*, the American Institute of Mathematics for supporting our square proposal, and an anonymous referee for valuable suggestions.

2. NOTATIONS AND PRELIMINARIES

More details on this section can be found in [21, Sections 2,3] and [25, Sections 2,3].

2.1. The group algebra of m -th roots of unity. Let $m, d \in \mathbb{Z}_{\geq 1}$. Let $\mu_m := \mu_m(\mathbb{C})$ denote the group of m -th roots of unity in \mathbb{C} . Let K_d be the d -th cyclotomic field over \mathbb{Q} . Let $\mathbb{Q}[\mu_m]$ denote the group algebra of μ_m over \mathbb{Q} . Then $\mathbb{Q}[\mu_m] = \prod_{d|m} K_d$. We endow $\mathbb{Q}[\mu_m]$ with the involution $*$ induced by the inverse map on μ_m , i.e., $\zeta^* := \zeta^{-1}$ for all $\zeta \in \mu_m$.

Set $\mathcal{T} := \text{hom}_{\mathbb{Q}}(\mathbb{Q}[\mu_m], \mathbb{C})$. If W is a $\mathbb{Q}[\mu_m] \otimes_{\mathbb{Q}} \mathbb{C}$ -module, we write $W = \bigoplus_{\tau \in \mathcal{T}} W_{\tau}$, where W_{τ} denotes the subspace of W on which $a \otimes 1 \in \mathbb{Q}[\mu_m] \otimes_{\mathbb{Q}} \mathbb{C}$ acts as $\tau(a)$. We fix an identification $\mathcal{T} = \mathbb{Z}/m\mathbb{Z}$ by defining, for all $n \in \mathbb{Z}/m\mathbb{Z}$,

$$\tau_n(\zeta) := \zeta^n, \text{ for all } \zeta \in \mu_m.$$

Let $m \geq 1$. For $p \nmid m$, we identify $\mathcal{T} = \text{hom}_{\mathbb{Q}}(\mathbb{Q}[\mu_m], \overline{\mathbb{Q}}_p^{\text{un}})$, where $\overline{\mathbb{Q}}_p^{\text{un}}$ is the maximal unramified extension of \mathbb{Q}_p in an algebraic closure. There is a natural action of the Frobenius σ on \mathcal{T} , defined by $\tau \mapsto \tau^{\sigma} := \sigma \circ \tau$. Let \mathfrak{O} be the set of all σ -orbits \mathfrak{o} in \mathcal{T} .

2.2. Newton polygons. Let X be an abelian scheme defined over the algebraic closure \mathbb{F} of \mathbb{F}_p . Then there is a finite field $\mathbb{F}_0/\mathbb{F}_p$, an abelian scheme X_0/\mathbb{F}_0 , and $\ell \in \mathbb{Z}_{\geq 1}$, such that $X \simeq X_0 \times_{\mathbb{F}_0} \mathbb{F}$ and the action of σ^{ℓ} on $H_{\text{cris}}^1(X_0/W(\mathbb{F}_0))$ is linear; here $W(\mathbb{F}_0)$ denotes the Witt vector ring of \mathbb{F}_0 . The *Newton polygon* $\nu(X)$ of X is the multi-set of rational numbers λ such that $\ell\lambda$ are the valuations at p of the eigenvalues of σ^{ℓ} acting on $H_{\text{cris}}^1(X_0/W(\mathbb{F}_0))$; the Newton polygon does not depend on the choice of $(\mathbb{F}_0, X_0, \ell)$.

The p -rank of X is the multiplicity of the slope 0 in $\nu(X)$; it equals $\dim_{\mathbb{F}_p}(\text{hom}(\mu_p, X))$.

If ν_1 and ν_2 are two Newton polygons, the *amalgamate sum* $\nu_1 \oplus \nu_2$ is the disjoint union of the multi-sets ν_1 and ν_2 . We denote by ν^d the amalgamate sum of d copies of ν .

The Newton polygon $\nu(X)$ is typically drawn as a lower convex polygon, with slopes λ occurring with multiplicity m_{λ} , where m_{λ} denotes the multiplicity of λ in the multi-set. The Newton polygon of a g -dimensional abelian variety is symmetric, with endpoints $(0,0)$ and $(2g, g)$, integral break points, and slopes in $\mathbb{Q} \cap [0, 1]$. For convex polygons, we write $\nu_1 \geq \nu_2$ if ν_1, ν_2 share the same endpoints and ν_1 lies below ν_2 .

We denote by ord the Newton polygon $(0, 1)$ and by ss the Newton polygon $(1/2, 1/2)$. For $s, t \in \mathbb{Z}_{\geq 1}$, with $s \leq t/2$ and $\text{gcd}(s, t) = 1$, we write $(s/t, (t-s)/t)$ for the Newton polygon with slopes s/t and $(t-s)/t$, each with multiplicity t .

Suppose Y is a semi-abelian scheme defined over \mathbb{F} . Then Y is an extension of an abelian scheme X by a torus T ; its Newton polygon is $\nu(Y) := \nu(X) \oplus \text{ord}^{\epsilon}$, where $\epsilon = \dim(T)$.

2.3. Cyclic covers of the projective line.

Definition 2.1. Fix integers $m \geq 2$, $N \geq 3$ and an N -tuple of integers $a = (a(1), \dots, a(N))$. Then a is an *inertia type* for m and $\gamma = (m, N, a)$ is a *monodromy datum* if

- (1) $a(i) \not\equiv 0 \pmod{m}$, for each $i = 1, \dots, N$,
- (2) $\text{gcd}(m, a(1), \dots, a(N)) = 1$,

$$(3) \sum_i a(i) \equiv 0 \pmod{m}.$$

For later applications, we sometimes consider a *generalized monodromy datum*, in which we allow $a(i) \equiv 0 \pmod{m}$. In the case that $a(i) = 0$, we set $\gcd(a(i), m) = m$.

Two monodromy data (m, N, a) and (m', N', a') are *equivalent* if $m = m'$, $N = N'$, and the images of a, a' in $(\mathbb{Z}/m\mathbb{Z})^N$ are in the same orbit under $(\mathbb{Z}/m\mathbb{Z})^* \times \text{Sym}_N$.

For fixed m , we work over an irreducible scheme over $\mathbb{Z}[1/m, \zeta_m]$. Let $U \subset (\mathbb{A}^1)^N$ be the complement of the weak diagonal. Consider the smooth projective curve C over U whose fiber at each point $t = (t(1), \dots, t(N)) \in U$ has affine model

$$(2.1) \quad y^m = \prod_{i=1}^N (x - t(i))^{a(i)}.$$

Consider the μ_m -cover $\phi : C \rightarrow \mathbb{P}_U^1$ defined by the function x and the μ_m -action $\iota : \mu_m \rightarrow \text{Aut}(C)$ given by $\iota(\zeta) \cdot (x, y) = (x, \zeta \cdot y)$ for all $\zeta \in \mu_m$.

For a closed point $t \in U$, the cover $\phi_t : C_t \rightarrow \mathbb{P}^1$ is a μ_m -cover, branched at N points $t(1), \dots, t(N)$ in \mathbb{P}^1 , and with local monodromy $a(i)$ at $t(i)$. By the hypotheses on the monodromy datum, C_t is a geometrically irreducible curve of genus g , where

$$(2.2) \quad g = g(m, N, a) = 1 + \frac{1}{2} \left((N-2)m - \sum_{i=1}^N \gcd(a(i), m) \right).$$

Take $W = H^0(C_t, \Omega^1)$ and, under the identification $\mathcal{T} = \mathbb{Z}/m\mathbb{Z}$, let $f(\tau_n) = \dim(W_{\tau_n})$. The *signature type* of ϕ is defined as $\mathfrak{f} = (f(\tau_1), \dots, f(\tau_{m-1}))$. By [25, Lemma 2.7, §3.2],

$$(2.3) \quad \mathfrak{f}(\tau_n) = \begin{cases} -1 + \sum_{i=1}^N \langle \frac{-na(i)}{m} \rangle & \text{if } n \not\equiv 0 \pmod{m} \\ 0 & \text{if } n \equiv 0 \pmod{m}. \end{cases}$$

where, for any $x \in \mathbb{R}$, $\langle x \rangle$ denotes the fractional part of x . The signature type of ϕ does not depend on t ; it determines and is uniquely determined by the inertia type.

2.4. Hurwitz spaces. Let \mathcal{M}_g be the moduli space of smooth curves of genus g in characteristic p . Its Deligne–Mumford compactification $\overline{\mathcal{M}}_g$ is the moduli space of stable curves of genus g . For a Newton polygon ν , let $\mathcal{M}_g[\nu]$ be the subspace whose points represent objects with Newton polygon ν . We use analogous notation for other moduli spaces.

We refer to [1, Sections 2.1-2.2] for a more complete description of Hurwitz spaces for cyclic covers of \mathbb{P}^1 . Consider the moduli functor $\overline{\mathcal{M}}_{\mu_m}$ (resp. $\tilde{\mathcal{M}}_{\mu_m}$) on the category of schemes over $\mathbb{Z}[1/m, \zeta_m]$; its points represent admissible stable μ_m -covers $(C/U, \iota)$ of a genus 0 curve (resp. together with an ordering of the smooth branch points and the choice of one ramified point above each of these). We use a superscript \circ to denote the subspace of points for which C is smooth. By [1, Lemma 2.2], $\overline{\mathcal{M}}_{\mu_m}$ (resp. $\tilde{\mathcal{M}}_{\mu_m}$) is a smooth proper Deligne–Mumford stack and $\overline{\mathcal{M}}_{\mu_m}^\circ$ (resp. $\tilde{\mathcal{M}}_{\mu_m}^\circ$) is open and dense within it.

For each irreducible component of $\tilde{\mathcal{M}}_{\mu_m}$, the monodromy datum $\gamma = (m, N, a)$ of the μ_m -cover $(C/U, \iota)$ is constant. Conversely, the substack $\tilde{\mathcal{M}}_{\mu_m}^\gamma$ of points representing μ_m -covers with monodromy datum γ is irreducible, [12, Corollary 7.5], [33, Corollary 4.2.3].

On $\overline{\mathcal{M}}_{\mu_m}$, there is no ordering of the ramification points; so only the unordered multi-set $\bar{a} = \{a(1), \dots, a(N)\}$ is well-defined. The components of $\overline{\mathcal{M}}_{\mu_m}$ are indexed by $\bar{\gamma} = (m, N, \bar{a})$. By [1, Lemma 2.4], the forgetful morphism $\tilde{\mathcal{M}}_{\mu_m}^{\gamma} \rightarrow \overline{\mathcal{M}}_{\mu_m}^{\bar{\gamma}}$ is étale and Galois.

Definition 2.2. ⁴ If $\gamma = (m, N, a)$ is a monodromy datum, let $\tilde{Z}(\gamma) = \tilde{\mathcal{M}}_{\mu_m}^{\gamma}$ and let $\overline{Z}(\gamma)$ be the reduced image of $\tilde{\mathcal{M}}_{\mu_m}^{\gamma}$ in $\overline{\mathcal{M}}_g$. We denote the subspace representing objects where C/U is smooth (resp. of compact type⁵, resp. stable) by

$$Z^{\circ}(\gamma) \subset Z(\gamma) \subset \overline{Z}(\gamma) \text{ and } \tilde{Z}^{\circ}(\gamma) \subset \tilde{Z}^c(\gamma) \subset \tilde{Z}(\gamma).$$

By definition, $\overline{Z}(\gamma)$ is a reduced irreducible proper substack of $\overline{\mathcal{M}}_g$. It depends uniquely on the equivalence class of γ . The forgetful morphism $\tilde{Z}(\gamma) \rightarrow \overline{Z}(\gamma)$ is finite and hence it preserves the dimension of any substack.

Remark 2.3. Let $\gamma' = (m, N', a')$ be a generalized monodromy datum. Assume that $a'(N') \equiv 0 \pmod{m}$ and $a'(i) \not\equiv 0 \pmod{m}$ for $1 \leq i < N'$. Consider the monodromy datum $\gamma = (m, N' - 1, a)$, where $a(i) = a'(i)$ for $1 \leq i \leq N' - 1$. Then $\overline{Z}(\gamma') = \overline{Z}(\gamma)$ and $\tilde{Z}(\gamma') = \tilde{Z}(\gamma)_1$, where the subscript 1 indicates that the data includes one marked point on the curve. The fibers of the morphism $\tilde{Z}^{\circ}(\gamma') \rightarrow Z^{\circ}(\gamma')$ are of pure dimension 1.

2.5. Shimura varieties associated to monodromy data. Consider $V := \mathbb{Q}^{2g}$ endowed with the standard symplectic form $\Psi : V \times V \rightarrow \mathbb{Q}$ and $G := \mathrm{GSp}(V, \Psi)$, the group of symplectic similitudes. Let $(G_{\mathbb{Q}}, \mathfrak{h})$ be the Siegel Shimura datum.

Fix $x \in Z(\gamma)(\mathbb{C})$ and let (\mathcal{J}_x, θ) denote the Jacobian of the curve represented by x together with its principal polarization θ . Choose a symplectic similitude

$$\alpha : (H_1(\mathcal{J}_x, \mathbb{Z}), \psi_{\theta}) \rightarrow (V, \Psi)$$

where ψ_{θ} denotes the Riemannian form on $H_1(\mathcal{J}_x, \mathbb{Q})$ corresponding to θ . Via α , the $\mathbb{Q}[\mu_m]$ -action on \mathcal{J}_x induces a $\mathbb{Q}[\mu_m]$ -module structure on V , and the Hodge decomposition of $H_1(\mathcal{J}_x, \mathbb{C})$ induces a $\mathbb{Q}[\mu_m] \otimes_{\mathbb{Q}} \mathbb{C}$ -linear decomposition $V_{\mathbb{C}} = V^+ \oplus V^-$.

We recall the PEL-type Shimura stack $\mathrm{Sh}(\mu_m, \mathfrak{f})$ given in [8]. The Shimura datum of $\mathrm{Sh}(\mu_m, \mathfrak{f})$ given by $(H, \mathfrak{h}_{\mathfrak{f}})$ is defined as

$$H := \mathrm{GL}_{\mathbb{Q}[\mu_m]}(V) \cap \mathrm{GSp}(V, \Psi),$$

and $\mathfrak{h}_{\mathfrak{f}}$ the H -orbit in $\{h \in \mathfrak{h} \mid h \text{ factors through } H\}$ determined by the isomorphism class of the $\mathbb{Q}[\mu_m] \otimes_{\mathbb{Q}} \mathbb{C}$ -module V^+ , i.e., by the integers $\mathfrak{f}(\tau) := \dim_{\mathbb{C}}(V_{\tau}^+)$, for all $\tau \in \mathcal{T}$. Under the identification $\mathcal{T} = \mathbb{Z}/m\mathbb{Z}$, the formula for $\mathfrak{f}(\tau_n)$ is that given in (2.3).

For a Shimura variety $\mathrm{Sh} := \mathrm{Sh}(H, \mathfrak{h}_{\mathfrak{f}})$ of PEL type, we use Sh^* to denote the Baily-Borel (i.e., minimal) compactification and $\overline{\mathrm{Sh}}$ to denote a toroidal compactification (see [19]).

The Torelli morphism $T : \mathcal{M}_g^c \rightarrow \mathcal{A}_g$ takes a curve of compact type to its Jacobian.

Definition 2.4. We say that $Z(\gamma)$ is *special* if $T(Z(\gamma))$ is open and closed in the PEL-type Shimura stack $\mathrm{Sh}(\mu_m, \mathfrak{f})$ given in [8] (see [21, Section 3.3] for details).

⁴This definition is slightly different from the one in our previous papers [22], [21].

⁵A stable curve has *compact type* if its dual graph is a tree. The Jacobian of a stable curve C is a semi-abelian variety; also C has compact type if and only if $\mathrm{Jac}(C)$ is an abelian variety.

If $N = 3$, then $T(Z(\gamma))$ is a point of \mathcal{A}_g representing an abelian variety with complex multiplication and is thus special, [22, Lemma 3.1]. By [25, Theorem 3.6], if $N \geq 4$, then $Z(\gamma)$ is special if and only if γ is equivalent to one of twenty examples in [25, Table 1].

2.6. The Kottwitz set and the μ -ordinary Newton polygon. Let $p \nmid m$ be a rational prime. Then the Shimura datum (H, \mathfrak{h}_f) is unramified at p . We write $H_{\mathbb{Q}_p}$ for the fiber of H at p , and $\mu_{\mathfrak{h}}$ for the conjugacy class of p -adic cocharacters μ_h associated with $h \in \mathfrak{h}_f$.

Following [16]-[18], we denote by $B(H_{\mathbb{Q}_p}, \mu_{\mathfrak{h}})$ the partially ordered set of μ -admissible $H_{\mathbb{Q}_p}$ -isocrystal structures on $V_{\mathbb{Q}_p}$. By [30, Theorem 1.6] (see also [32]), $B(H_{\mathbb{Q}_p}, \mu_{\mathfrak{h}})$ can be canonically identified with the set of Newton polygons appearing on $\text{Sh}(H, \mathfrak{h})$.⁶ We sometimes write $\text{Sh} := \text{Sh}(H, \mathfrak{h})$ and $B = B(\text{Sh}) := B(H_{\mathbb{Q}_p}, \mu_{\mathfrak{h}})$.

Definition 2.5. The μ -ordinary Newton polygon $u := u_{\mu\text{-ord}}$ is the unique maximal element (lowest Newton polygon) of $B(H_{\mathbb{Q}_p}, \mu_{\mathfrak{h}})$.

An explicit formula for u is given below; (see also [21, Section 4.1-4.2]).

2.6.1. Formula for slopes and multiplicities. Let \mathfrak{f} be a signature type. Fix an orbit \mathfrak{o} in \mathcal{T} . We recall the formulas from [25, Section 1.2.5] for the slopes and multiplicities of the \mathfrak{o} -component $u(\mathfrak{o})$ of the μ -ordinary Newton polygon in terms of \mathfrak{f} , following the notation in [9, Section 2.8], [21, Section 4.2].

With some abuse of notation, we replace \mathcal{T} by $\mathcal{T} - \{\tau_0\}$ and \mathfrak{D} by $\mathfrak{D} - \{\{\tau_0\}\}$. Let $g(\tau) := \dim_{\mathbb{C}}(V_{\tau})$. As the integer $g(\tau)$ depends only on the order of τ in the additive group $\mathbb{Z}/m\mathbb{Z}$, and thus only on the orbit \mathfrak{o} of τ , we sometimes write $g(\mathfrak{o}) = g(\tau)$, for any/all $\tau \in \mathfrak{o}$.

Remark 2.6. For all $\tau \in \mathcal{T}$, $\dim_{\mathbb{C}}(V_{\tau^*}^+) = \dim_{\mathbb{C}}(V_{\tau}^-)$, and thus $\mathfrak{f}(\tau) + \mathfrak{f}(\tau^*) = g(\tau)$.

Let $s = s(\mathfrak{o})$ be the number of distinct values of $\{\mathfrak{f}(\tau) \mid \tau \in \mathfrak{o}\}$ in the range $[1, g(\mathfrak{o}) - 1]$. We write these distinct values as

$$g(\mathfrak{o}) > E(1) > E(2) > \cdots > E(s) > 0.$$

Let $E(0) := g(\mathfrak{o})$ and $E(s+1) := 0$. Then $u(\mathfrak{o})$ has exactly $s+1$ distinct slopes, denoted by $0 \leq \lambda(0) < \lambda(1) < \cdots < \lambda(s) \leq 1$. For $0 \leq t \leq s$, the $(t+1)$ -st slope is

$$(2.4) \quad \lambda(t) := \frac{1}{|\mathfrak{o}|} \#\{\tau \mid \mathfrak{f}_i(\tau) \geq E(t)\}.$$

The slope $\lambda(t)$ occurs in $u(\mathfrak{o})$ with multiplicity

$$(2.5) \quad \rho(\lambda(t)) := |\mathfrak{o}|(E(t) - E(t+1)).$$

⁶The term Newton polygon usually refers to the image of an element in B via the Newton map. Here, since we work with PEL-type Shimura varieties of type A and C, the Newton map is injective and hence we do not distinguish between the elements in B and the corresponding Newton polygons.

2.7. Geometry of the Newton polygon strata on Sh . For $b \in B$, let $\text{Sh}[b] := \text{Sh}(H, \mathfrak{h})[b]$ denote the corresponding Newton polygon stratum in Sh . In other words, $\text{Sh}[b]$ is the locally closed substack of Sh parametrizing abelian schemes with Newton polygon b . By Hamacher [14, Theorem 1.1, Corollary 3.12], based on the work of Chai [6], Mantovan [23], and Viehmann [29], and Kottwitz [17, Section 8],⁷ on each irreducible component S of Sh , the substack $S[b]$ is non-empty and equidimensional and

$$(2.6) \quad \text{codim}(S[b], \text{Sh}) = \text{length}(b),$$

where $\text{length}(b) = \max\{n \mid \text{there exists a chain } b = v_0 < v_1 < \cdots < v_n = u, v_i \in B\}$.

The Newton stratification extends to the toroidal and minimal compactifications Sh, Sh^* . In [20, §3.3], the authors studied the Newton stratification on compactifications of PEL-type Shimura varieties at good primes. They proved in this case that all the Newton strata are so called *well-positioned* subschemes [20, Proposition 3.3.9]. In particular, by [20, Definition 2.2.1, Theorem 2.3.2], the set of Newton polygons on (each irreducible component of) $\overline{\text{Sh}}$ is the same as that on Sh and, for any $b \in B$,

$$(2.7) \quad \text{codim}(\overline{\text{Sh}}[b], \overline{\text{Sh}}) = \text{codim}(\text{Sh}[b], \text{Sh}).$$

By the next remark, there exists a μ_m -cover of smooth curves having monodromy datum γ and μ -ordinary Newton polygon u if there exists such a cover of stable curves.

Lemma 2.7. *The following are equivalent: $Z^\circ(\gamma)[u]$ is non-empty; $Z^\circ(\gamma)[u]$ is open and dense in $Z(\gamma)$; and $Z(\gamma)[u]$ is non-empty.*

Proof. This is clear because the Newton polygon is lower semi-continuous, $Z(\gamma)$ is irreducible, and $Z^\circ(\gamma)$ is open and dense in $Z(\gamma)$. \square

Remark 2.8. The Ekedahl–Oort type is also determined for many of the smooth curves in this paper. The reason is that the μ -ordinary Newton polygon stratum in these PEL-type Shimura varieties coincides with the unique open Ekedahl–Oort stratum. Hence one may compute the Ekedahl–Oort type of these smooth curves using [24, Section 1.2.3].

3. CLUTCHING MORPHISMS

3.1. Background. We study a generalization of the clutching morphisms, for curves that are μ_m -covers of \mathbb{P}^1 . The clutching morphisms are the closed immersions [15, 3.9]:

$$\kappa_{i,g-i} : \overline{\mathcal{M}}_{i,1} \times \overline{\mathcal{M}}_{g-i,1} \rightarrow \overline{\mathcal{M}}_g \text{ and } \lambda : \overline{\mathcal{M}}_{g-1,2} \rightarrow \overline{\mathcal{M}}_g,$$

for $1 \leq i \leq g-1$. The image of $\kappa_{i,g-i}$ is the component Δ_i of the boundary of \mathcal{M}_g , whose generic point represents a stable curve that has two components, of genus i and $g-i$, intersecting in one point. The image of λ is the component Δ_0 of the boundary of \mathcal{M}_g , whose points represent stable curves that do not have compact type.

Given a pair of cyclic covers of \mathbb{P}^1 , we analyze the image of a clutching morphism κ , which shares attributes of both $\kappa_{i,g-i}$ and λ . To provide greater flexibility, we include cases when the covers have different degrees or when the two covers are clutched together at several points. As a result, a curve in the image of κ may not have compact type.

⁷Hamacher proved that $\text{Sh}[b]$ is non-empty and equidimensional of expected dimension. Since Hecke translations preserve the Newton polygon strata and act transitively on the irreducible components of Sh , we deduce the same result for $S[b]$. See [17, §8] for a more detailed discussion.

Notation 3.1. Let $\gamma = (m, N, a)$ be a monodromy datum. For an integer $d \geq 1$, consider the induced monodromy data $\gamma^{\dagger d} = (dm, N, da)$, which we sometimes denote γ^\dagger .

If $\phi : C \rightarrow \mathbb{P}^1$ is a μ_m -cover with monodromy datum γ , then $\text{Ind}_m^{dm}(\phi) : \text{Ind}_m^{dm} C \rightarrow \mathbb{P}^1$ is a μ_{dm} -cover with the induced monodromy datum $\gamma^{\dagger d}$. The signature type of $\text{Ind}_m^{dm}(\phi)$ is $\mathfrak{f}^{\dagger d} = \mathfrak{f} \circ \pi_d$, where $\pi_d : \mathbb{Z}/dm\mathbb{Z} \rightarrow \mathbb{Z}/m\mathbb{Z}$ denotes the natural projection. By definition, the μ -ordinary polygon of $\text{Sh}(\mu_{dm}, \mathfrak{f}^{\dagger d})$ is $u^{\dagger d} = u^d$.

3.2. Numerical data and hypothesis (A).

Notation 3.2. Fix integers $m_1, m_2 \geq 2$, $N_1, N_2 \geq 3$. Let $m_3 = \text{lcm}(m_1, m_2)$. For $i = 1, 2$: let $d_i = m_3/m_i$; let $a_i = (a_i(1), \dots, a_i(N_i))$ be such that $\gamma_i = (m_i, N_i, a_i)$ is a (generalized) monodromy datum; and let $g_i = g(m_i, N_i, a_i)$ as in (2.2).

Definition 3.3. A pair of monodromy data $\gamma_1 = (m_1, N_1, a_1)$, $\gamma_2 = (m_2, N_2, a_2)$ as in Notation 3.2 is *admissible* if it satisfies

$$\text{hypothesis(A)} : d_1 a_1(N_1) + d_2 a_2(1) \equiv 0 \pmod{m_3}.$$

Notation 3.4. Assume hypothesis (A) for the pair γ_1, γ_2 . Set $r_1 = \text{gcd}(m_1, a_1(N_1))$, and $r_2 = \text{gcd}(m_2, a_2(1))$. Let $r_0 = \text{gcd}(r_1, r_2)$ and let

$$(3.1) \quad \epsilon = d_1 d_2 r_0 - d_1 - d_2 + 1 \text{ and } g_3 = d_1 g_1 + d_2 g_2 + \epsilon.$$

Note that $d_1 r_1 = d_2 r_2 = d_1 d_2 r_0$ since $\text{gcd}(d_1, d_2) = 1$.

Definition 3.5. If γ_1, γ_2 is an admissible pair of (generalized) monodromy data, we define $\gamma_3 = (m_3, N_3, a_3)$ by $m_3 := \text{lcm}(m_1, m_2)$, $N_3 := N_1 + N_2 - 2$ and the N_3 -tuple a_3 as

$$a_3(i) := \begin{cases} d_1 a_1(i) & \text{for } 1 \leq i \leq N_1 - 1, \\ d_2 a_2(i - N_1 + 2) & \text{for } N_1 \leq i \leq N_1 + N_2 - 2. \end{cases}$$

Lemma 3.6. *The triple γ_3 from Definition 3.5 is a (generalized) monodromy datum. If $\phi_3 : C \rightarrow \mathbb{P}^1$ is a cover with monodromy datum γ_3 , then the genus of C is g_3 as in (3.1).*

Proof. Immediate from Definition 2.1 and (2.2). □

The signature type for ϕ_3 is given in Definition 3.12, see Lemma 3.13.

Remark 3.7. A pair γ_1, γ_2 of non-admissible monodromy data can be modified slightly to produce a pair γ'_1, γ'_2 of admissible generalized monodromy data by marking an extra unramified fiber. Specifically, let

- (1) $\gamma'_1 = (m_1, N_1 + 1, a'_1)$ with $a'_1(i) = a_1(i)$ for $1 \leq i \leq N_1$, and $a'_1(N_1 + 1) = 0$;
- (2) $\gamma'_2 = (m_2, N_2 + 1, a'_2)$ with $a'_2(1) = 0$ and $a'_2(i) = a_2(i - 1)$ for $2 \leq i \leq N_2 + 1$.

This does not change the geometry, because $Z(\gamma'_i) = Z(\gamma_i)$ for $i = 1, 2$ by Remark 2.3.

3.3. Clutching morphisms for cyclic covers.

Notation 3.8. Let $\gamma_1 = (m_1, N_1, a_1)$, $\gamma_2 = (m_2, N_2, a_2)$ be an admissible pair of monodromy data as in Notation 3.2. Let $\gamma_3 = (m_3, N_3, a_3)$ be the monodromy datum from Definition 3.5. For $i = 1, 2, 3$, let $\tilde{Z}_i = \tilde{Z}(\gamma_i)$ be as in Definition 2.2.

Recall that the points of \tilde{Z}_3° represent μ_{m_3} -covers $C \rightarrow \mathbb{P}^1$ with monodromy datum γ_3 , where C is smooth. The next result is more general than related results in the literature.

Proposition 3.9. *If hypothesis (A) (the admissible condition) is satisfied, there is a clutching morphism $\kappa : \tilde{Z}_1 \times \tilde{Z}_2 \rightarrow \tilde{Z}_3$, and the image of κ is in the boundary of \tilde{Z}_3° .*

Proof. For $i = 1, 2$, let $\phi_i : C_i \rightarrow \mathbb{P}^1$ be the μ_{m_i} -cover with N_i ordered and labeled μ_{m_i} -orbits of points which is represented by a point of \tilde{Z}_i . There is a natural inclusion $\mu_{m_i} \subset \mu_{m_3}$. Let $C_i^\dagger = \text{Ind}_{m_i}^{m_3}(C_i)$ be the induced curve and let $\phi_i^\dagger : C_i^\dagger \rightarrow \mathbb{P}^1$ be the induced cover. It has inertia type $a_i^\dagger = d_i a_i = (d_i a_i(1), \dots, d_i a_i(N_i))$.

We define the morphism κ on the pair (ϕ_1, ϕ_2) . There are $d_1 r_1 = d_1 \text{gcd}(m_1, a_1(N_1))$ points of C_1^\dagger above $t_1(N_1)$. By hypothesis (A), this equals $d_2 r_2 = d_2 \text{gcd}(m_2, a_2(1))$, the number of points of C_2^\dagger above $t_2(1)$. Let C_3 be the curve whose components are the d_1 components of C_1^\dagger and the d_2 components of C_2^\dagger , formed by identifying these labeled $d_1 r_1 = d_2 r_2$ points in ordinary double points, in a μ_{m_3} -equivariant way.

Then C_3 is a μ_{m_3} -cover of a tree P of two projective lines. It has N_3 labeled μ_{m_3} -orbits with inertia type a_3 and is thus represented by a point of \tilde{Z}_3 . The admissible condition in Definition 3.3 is exactly the (local) admissible condition on the covers ϕ_1^\dagger and ϕ_2^\dagger at the point(s) above $t_1(N_1)$ and $t_2(1)$. By [10, 2.2], the μ_{m_3} -cover $C_3 \rightarrow P$ is in the boundary of \tilde{Z}_3° if and only if hypothesis (A) is satisfied. \square

The curve C_3 constructed in Proposition 3.9 is a μ_{m_3} -cover of type γ_3 and thus has arithmetic genus g_3 by Lemma 3.6.

Proposition 3.10. *The curve C_3 constructed in the proof of Proposition 3.9 has compact type if and only if $\epsilon = 0$. It has Newton polygon*

$$(3.2) \quad \nu(C_3) = \nu(C_1)^{d_1} \oplus \nu(C_2)^{d_2} \oplus \text{ord}^\epsilon.$$

The term ord^ϵ can be viewed as the *defect* of $\nu(C_3)$. It measures the number of extra slopes of 0 and 1 that arise when C_3 does not have compact type. By Notation 3.4, $\epsilon = 0$ if and only if $r_0 = 1$ and either $d_1 = 1$ or $d_2 = 1$.

Proof. By construction, the dual graph of C_3 is a bipartite graph, with d_1 (resp. d_2) vertices in bijection with the components of C_1^\dagger (resp. C_2^\dagger). In C_3 , each of the components of C_1^\dagger intersects each of the components of C_2^\dagger in r_0 points. After removing $d_1 d_2 (r_0 - 1)$ edges from the dual graph, there is a unique edge between each pair of vertices on opposite sides. After removing another $(d_1 - 1)(d_2 - 1)$ edges from the dual graph, it is a tree. Thus the Euler characteristic of the dual graph of C_3 is $d_1 d_2 (r_0 - 1) + (d_1 - 1)(d_2 - 1)$, which equals ϵ . In particular, C_3 has compact type if and only if $\epsilon = 0$.

By [4, 9.2.8], for some torus T of rank ϵ , there is a short exact sequence

$$0 \rightarrow T \rightarrow \text{Jac}(C_3) \rightarrow \text{Jac}(C_1)^{d_1} \oplus \text{Jac}(C_2)^{d_2} \rightarrow 0.$$

Since $\dim_{\mathbb{F}_p}(\mu_p, T) = \epsilon$, the Newton polygon of $\text{Jac}(C_3)$ is the amalgamate sum (Section 2.2) of the Newton polygon of $\text{Jac}(C_1)^{d_1} \oplus \text{Jac}(C_2)^{d_2}$ and ord^ϵ , which yields (3.2). \square

3.4. The signature. We find the signature f_3 for a cover with monodromy datum γ_3 .

Definition 3.11. Let $d, R \in \mathbb{Z}_{\geq 1}$ with $dR \mid m$. For $n \in \mathbb{Z}/m\mathbb{Z}$, we define $\delta_{d,dR}(n) := 1$ if $dRn \equiv 0 \pmod{m}$ and $dn \not\equiv 0 \pmod{m}$, and $\delta_{d,dR}(n) := 0$ otherwise.

Equivalently, if $\tau \in \mathcal{T}$, then $\delta_{d,dR}(\tau) = 1$ if the order of τ in $\mathbb{Z}/m\mathbb{Z}$ divides dR but not d , and $\delta_{d,dR}(\tau) = 0$ otherwise. Since $\delta_{d,dR}(\tau)$ only depends on the orbit \mathfrak{o} of τ , we also write $\delta_{d,dR}(\mathfrak{o}) := \delta_{d,dR}(\tau)$, for any/all $\tau \in \mathfrak{o}$.

Definition 3.12. Let $\delta := \delta_{d,dr_0} + \delta_{d_1,d_1d_2} - \delta_{1,d_2}$, for $d = d_1d_2$. For $n \in \mathbb{Z}/m_3\mathbb{Z} - \{0\}$, let

$$(3.3) \quad \mathfrak{f}_3(\tau_n) = \mathfrak{f}_1^\dagger(\tau_n) + \mathfrak{f}_2^\dagger(\tau_n) + \delta(n).$$

By definition, $\delta(n) = 1$ if $d_1d_2r_0n \equiv 0 \pmod{m_3}$ and $d_1n \not\equiv 0 \pmod{m_3}$ and $d_2n \not\equiv 0 \pmod{m_3}$, and $\delta(n) = 0$ otherwise.

Lemma 3.13. *If $\phi_3 : C \rightarrow \mathbb{P}^1$ is a cover with (generalized) monodromy datum γ_3 , as defined in Definition 3.5, then the signature type of C_3 is \mathfrak{f}_3 .*

Proof. We use (2.3) to compute \mathfrak{f}_3 .⁸ If $n \equiv 0 \pmod{m_1}$, then $\mathfrak{f}_1^\dagger(\tau_n) = 0$ and $\mathfrak{f}_3(\tau_n) = \mathfrak{f}_2^\dagger(\tau_n)$. If $n \equiv 0 \pmod{m_2}$, then $\mathfrak{f}_2^\dagger(\tau_n) = 0$ and $\mathfrak{f}_3(\tau_n) = \mathfrak{f}_1^\dagger(\tau_n)$. For $n \in \mathbb{Z}/m_3\mathbb{Z}$, with $n \not\equiv 0 \pmod{m_1}$ and $n \not\equiv 0 \pmod{m_2}$, then

$$(\mathfrak{f}_1^\dagger(\tau_n) + 1) + (\mathfrak{f}_2^\dagger(\tau_n) + 1) - (\mathfrak{f}_3(\tau_n) + 1) = \left\langle \frac{-nd_1a_1(N_1)}{m_3} \right\rangle + \left\langle \frac{-nd_2a_2(1)}{m_3} \right\rangle.$$

The right hand side is 0 or 1; it is 0 if and only if $nd_1a_1(N_1) \equiv nd_2a_2(1) \equiv 0 \pmod{m_3}$. \square

3.5. Compatibility with Shimura variety setting.

Notation 3.14. Fix an admissible pair $\gamma_1 = (m_1, N_1, a_1)$, $\gamma_2 = (m_2, N_2, a_2)$ of monodromy data as in Notation 3.2. Consider the monodromy datum γ_3 as in Definition 3.5. In particular, let $m_3 = \text{lcm}(m_1, m_2)$ and let \mathfrak{f}_3 be as in Definition 3.12.

For each $i = 1, 2, 3$, let $Z_i := Z(m_i, N_i, a_i)$, and similarly Z_i°, \tilde{Z}_i^c , etc as in Definition 2.2. Let $\text{Sh}_i := \text{Sh}_i(\mu_{m_i}, \mathfrak{f}_i)$ denote the Shimura substack of \mathcal{A}_g as in Section 2.5. Let \mathcal{X}_i be the universal abelian scheme over Sh_i , $B_i := B(\text{Sh}_i)$ the set of Newton polygons of Sh_i , and u_i the μ -ordinary Newton polygon in B_i from Definition 2.5.

Via the Torelli map T , the clutching morphism $\kappa : \tilde{Z}_1^c \times \tilde{Z}_2^c \rightarrow \tilde{Z}_3$ is compatible with a morphism into the minimal compactification of the Shimura variety

$$\iota : \text{Sh}_1 \times \text{Sh}_2 \rightarrow \text{Sh}_3^*$$

where $\iota(\mathcal{X}_1, \mathcal{X}_2) := \mathcal{X}_1^{d_1} \oplus \mathcal{X}_2^{d_2}$.⁹ If $\epsilon = 0$ then $\text{Im}(\iota)$ lies in Sh_3 and the reader may focus on this case; if $\epsilon \neq 0$, then $\text{Im}(\iota)$ is contained in the boundary $\text{Sh}_3^* - \text{Sh}_3$.

By Proposition 3.10, $\iota(\text{Sh}_1[v_1], \text{Sh}_2[v_2]) \subseteq \text{Sh}_3^*(v_1^{d_1} \oplus v_2^{d_2} \oplus \text{ord}^\epsilon)$, which yields:

Lemma 3.15. *If $v_i \in B_i$, then $v_1^{d_1} \oplus v_2^{d_2} \oplus \text{ord}^\epsilon \in B_3$. In particular, $u_3 \geq u_1^{d_1} \oplus u_2^{d_2} \oplus \text{ord}^\epsilon$.*

In Proposition 4.4, we give a necessary and sufficient condition on the pair of signature types $(\mathfrak{f}_1, \mathfrak{f}_2)$ for the equality $u_3 = u_1^{d_1} \oplus u_2^{d_2} \oplus \text{ord}^\epsilon$ to hold.

⁸Alternatively, one may deduce the formula for \mathfrak{f}_3 geometrically since the extra term δ records the $\mathbb{Z}/m_3\mathbb{Z}$ -action on the dual graph of the curve C_3 constructed in Proposition 3.9.

⁹An abelian variety of dimension less than g_3 with μ_m -action can be viewed as a point on the boundary of Sh_3^* as it comes from the pure part of some semi-abelian variety of dimension g_3 with μ_m -action, which is a point on $\overline{\text{Sh}}_3$. The image of the Torelli map in the minimal compactification is determined by the Torelli map on the irreducible components of the curve and forgets the dual graph structure.

4. THE TORELLI LOCUS AND THE μ -ORDINARY LOCUS OF SHIMURA VARIETIES

In this section, we prove theorems about the intersection of the open Torelli locus with the μ -ordinary Newton polygon stratum in a PEL-type Shimura variety. The main result, Theorem 4.5, provides a method to leverage information from smaller dimension to larger dimension. This provides an inductive method to prove that the open Torelli locus intersects the μ -ordinary stratum for certain types of families.

In Section 4.3, we use the main theorem to establish the existence of smooth curves of arbitrarily large genus with prescribed Newton polygon, see Corollaries 4.7 to 4.10. For the base cases of the inductive method, we can use any instances when the μ -ordinary Newton polygon is known to occur (see Proposition 4.6).

Remark 4.1. The method in this section does not give results for every monodromy datum γ . For example, it is not known whether the μ -ordinary Newton polygon occurs on $Z^\circ(\gamma)$ for all $p \equiv 3, 5 \pmod{7}$ when $\gamma = (7, 4, (1, 1, 2, 3))$. In this case, $\mathfrak{f} = (2, 1, 1, 1, 1, 0)$, $\dim(Z(\gamma)) = 1$, and $\dim(S(\gamma)) = 2$. The three Newton polygons on $S(\gamma)$ are $(1/6, 5/6)$, $(1/3, 2/3)^2$, and ss^6 , which all have p -rank 0. None of the degenerations for this family satisfy hypothesis (B) as defined below.

4.1. Hypothesis (B). We fix an admissible pair $\gamma_1 = (m_1, N_1, a_1)$, $\gamma_2 = (m_2, N_2, a_2)$ of (generalized) monodromy data as in Notation 3.2. We fix a prime p such that $p \nmid m_3 = \text{lcm}(m_1, m_2)$ and work over $\overline{\mathbb{F}}_p$. Recall Notation 3.4 and 3.14. So $\bar{d}_i = m_3/m_i$ and $\mathfrak{f}_i^\dagger := \mathfrak{f}_i \circ \pi_i$ for $i = 1, 2$.

Definition 4.2. The pair of monodromy data γ_1, γ_2 is *balanced* if, for each orbit $\mathfrak{o} \in \mathcal{T} = \text{hom}_{\mathbb{Q}}(\mathbb{Q}[\mu_{m_3}], \mathbb{C})$ and all $\omega, \tau \in \mathfrak{o}$, the values of the induced signature types satisfy:

hypothesis(B) : if $\mathfrak{f}_1^\dagger(\omega) > \mathfrak{f}_1^\dagger(\tau)$ then $\mathfrak{f}_2^\dagger(\omega) \geq \mathfrak{f}_2^\dagger(\tau)$; if $\mathfrak{f}_2^\dagger(\omega) > \mathfrak{f}_2^\dagger(\tau)$ then $\mathfrak{f}_1^\dagger(\omega) \geq \mathfrak{f}_1^\dagger(\tau)$.

Remark 4.3. (1) If $m_1 = m_2$, then hypothesis (B) is symmetric for the pair γ_1, γ_2 .
 (2) If $\gamma_1 = \gamma_2$, then hypothesis (B) is automatically satisfied.
 (3) Hypothesis (B) depends (only) on the congruence of p modulo m_3 . If $p \equiv 1 \pmod{m_3}$, then each orbit \mathfrak{o} in \mathcal{T} has size one and hypothesis (B) is vacuously satisfied.
 (4) Let γ_3 be as in Definition 3.5. If the pair γ_1, γ_2 satisfies hypotheses (A) and (B), then γ_i, γ_3 satisfies hypothesis (B) for $i = 1$ and for $i = 2$.

Proposition 4.4 below gives a geometric interpretation of hypothesis (B). Recall that $\epsilon = d_1 d_2 r_0 - d_1 - d_2 + 1$ from Notation 3.4.

Proposition 4.4. *Let γ_1, γ_2 be an admissible pair of monodromy data. Consider the monodromy datum γ_3 as in Definition 3.5. Then the equality $u_3 = u_1^{d_1} \oplus u_2^{d_2} \oplus \text{ord}^\epsilon$ holds if and only if the pair γ_1, γ_2 is balanced.*

We postpone the proof of Proposition 4.4 to the independent Section 5.

4.2. A first main result. In this subsection, we assume that the pair γ_1, γ_2 is admissible and balanced, meaning that it satisfies hypotheses (A) and (B) as in Definitions 3.3 and 4.2. Let $\gamma_3 = (m_3, N_3, a_3)$ and \mathfrak{f}_3 be as in Definitions 3.5 and 3.12.

The next result provides a partial positive answer to Conjecture 1.1 when $\epsilon = 0$.

Theorem 4.5. *Let γ_1, γ_2 be an admissible, balanced pair of monodromy data. If $Z_1^\circ[u_1]$ and $Z_2^\circ[u_2]$ are both non-empty, then $Z_3^\circ[u_3]$ is non-empty.*

Proof. By Lemmas 3.6 and 3.13, the signature for the monodromy datum γ_3 is given in (3.3). By Proposition 4.4, hypothesis (B) implies that $u_3 = u_1^{d_1} \oplus u_2^{d_2} \oplus \text{ord}^\epsilon$. From the hypothesis, $\tilde{Z}_1^\circ[u_1]$ and $\tilde{Z}_2^\circ[u_2]$ are both non-empty. By Proposition 3.9, the image of κ on $\tilde{Z}_1^\circ[u_1] \times \tilde{Z}_2^\circ[u_2]$ is in \tilde{Z}_3 . By Proposition 3.10, the Newton polygon of a curve C_3 represented by a point in the image of κ is given by $\nu(C_3) = u_1^{d_1} \oplus u_2^{d_2} \oplus \text{ord}^\epsilon$, which is u_3 . Thus $Z_3[u_3]$ is non-empty and applying Lemma 2.7 finishes the proof. \square

4.3. Infinite clutching for μ -ordinary. In this section, we find situations in which Theorem 4.5 can be implemented recursively, infinitely many times, to verify the existence of smooth curves of arbitrarily large genus with prescribed Newton polygon. The required input is a family (or a compatible pair of families) of cyclic covers of \mathbb{P}^1 for which the μ -ordinary Newton polygon at a prime p is known to occur (see Proposition 4.6). Concrete implementations of these results can be found in Section 9.

4.3.1. *Base cases.* We recall instances when the the μ -ordinary Newton polygon u is known to occur for the Jacobian of a (smooth) curve in the family Z .

Proposition 4.6. *The μ -ordinary Newton polygon stratum $Z(m, N, a)[u]$ is non empty if either:*

- (1) $N = 3$; or
- (2) $N \geq 4$ and (m, N, a) is equivalent to one of the twenty examples in [25, Table 1]; or
- (3) u is the only Newton polygon in $B(H_{\mathbb{Q}_p}, \mu_{\mathfrak{h}})$ of maximum p -rank and either $N = 4$ or $p \geq m(N - 3)$ or $p \equiv \pm 1 \pmod{m}$.

Proof. (1) When $N = 3$, then Z is 0-dimensional and thus special as in Definition 2.4.
 (2) This is [21, Proposition 5.1].
 (3) For any monodromy datum $\gamma = (m, N, a)$, define

$$(4.1) \quad \beta(\gamma) := \sum_{\tau \in \mathcal{T}} \min_{j \in \mathbb{N}} \{f(\tau^{\sigma^j})\} = \sum_{\mathfrak{o} \in \mathcal{D}} \#\mathfrak{o} \cdot \min\{f(\tau) \mid \tau \in \mathfrak{o}\}.$$

By [5, Equation (1)], $\beta(\gamma)$ is an upper bound for the p -rank of curves in $Z^\circ(\gamma)$. By [5, Theorem 6.1, Propositions 7.7, 7.4, 7.8], if $p \geq m(N - 3)$ or $N = 4$ or $p \equiv \pm 1 \pmod{m}$, then there exists a μ_m -cover $C \rightarrow \mathbb{P}^1$ defined over $\overline{\mathbb{F}}_p$ with monodromy datum γ , for which the p -rank of C equals $\beta(\gamma)$. The p -rank is the multiplicity of 1 as a slope of the Newton polygon. From the formulas (2.4) and (2.5) for the slopes and multiplicities of the μ -ordinary Newton polygon $u = u(\gamma)$, one can check that the p -rank of u equals $\beta(\gamma)$. \square

We refer to [22] for the computation of the μ -ordinary polygon u when $N = 3$, and to Section 10 for the special families of [25] (see [21]).

4.3.2. *Adding slopes 0 and 1.* By implementing Theorem 4.5 recursively, we obtain a method to increase the genus and the multiplicity of the slopes 0 and 1 in the Newton polygon by the same amount. Because of this, in later results we will aim to minimize the multiplicity of $\{0, 1\}$ in the Newton polygon.

Corollary 4.7. *Let $\gamma = (m, N, a)$ be a monodromy datum. Assume that $Z^\circ(\gamma)[u]$ is non-empty. Then for any n in the semi-group generated by $\{m - t : t \mid m\}$, there exists a μ_m -cover $C \rightarrow \mathbb{P}^1$ over $\overline{\mathbb{F}}_p$ where C is a smooth curve with Newton polygon $u \oplus \text{ord}^n$.*

Proof. For $c \in \mathbb{Z}_{\geq 1}$ with $c \leq m - 1$, let $t = \gcd(m, c)$ and consider the monodromy datum $\gamma_1 = (m/t, 3, (c/t, (m - c)/t, 0))$. Note $g_1(\tau) = 0$, and $f_1(\tau) = 0$, for all $\tau \in \mathcal{T}$.

Let $\gamma_2 = (m, N + 1, a')$, where $a'(1) = 0$, and $a'(i) = a(i - 1)$, for $i = 2, \dots, N + 1$. By construction, the pair γ_1, γ_2 is admissible and balanced. By Remark 2.3, $Z^\circ(\gamma_2)[u]$ is non-empty. Set $a_3 = (c, m - c, a(1), \dots, a(N))$; then $\gamma_3 = (m, N + 2, a_3)$ is the monodromy datum from Definition 3.5 for the pair γ_1, γ_2 . By (3.1), $\epsilon = m - t$. By Theorem 4.5, $Z_3^\circ[u_3]$ is non-empty, where $u_3 = u \oplus \text{ord}^{m-t}$. The statement follows by iterating this construction, letting c vary. \square

4.3.3. Single Induction. We consider inductive systems generated by a single monodromy datum γ_1 . The next result follows from the observation that if the pair γ_1, γ_1 is admissible and if $Z_1^\circ[u_1]$ is non-empty, then all hypotheses of Theorem 4.5 are satisfied, and continue to be after iterations.

Corollary 4.8. *Assume there exist $1 \leq i < j \leq N$ such that $a(i) + a(j) \equiv 0 \pmod{m}$.¹⁰ Let $r = \gcd(a(i), m)$. If $Z^\circ(\gamma)[u]$ is non-empty, then there exists a smooth curve over $\overline{\mathbb{F}}_p$ with Newton polygon $u^n \oplus \text{ord}^{(n-1)(r-1)}$, for any $n \in \mathbb{Z}_{\geq 1}$.*

Proof. After reordering the branch points, we can suppose that $i = 1, j = N$. We define a sequence of families $Z^{\times n}$ as follows: let $Z^{\times 1} = Z$; for $n \geq 2$, let $Z^{\times n}$ be the family constructed from the monodromy datum produced by applying Definition 3.5 to the monodromy data of $Z^{\times(n-1)}$ and $Z^{\times 1}$. For $n \in \mathbb{Z}_{\geq 1}$, the pair of monodromy data for $Z^{\times n}$ and $Z^{\times 1}$ satisfies hypotheses (A) and (B). Then $u_n := u^n \oplus \text{ord}^{(n-1)(r-1)}$ is the μ -ordinary Newton polygon for $Z^{\times n}$. The statement follows by applying Theorem 4.5 repeatedly. \square

The first hypothesis of Corollary 4.8 appears restrictive. However, from any monodromy datum γ with $Z^\circ(\gamma)[u]$ non-empty, we can produce a new monodromy datum which satisfies this hypothesis by clutching with a μ_m -cover branched at only two points. As a result, Corollary 4.8 can be generalized to Corollary 4.9 which holds in much greater generality, at the expense of making the defect slightly larger.

Corollary 4.9. *Assume that $Z^\circ(\gamma)[u]$ is non-empty and let t be a positive divisor of m . Then there exists a smooth curve over $\overline{\mathbb{F}}_p$ with Newton polygon $u^n \oplus \text{ord}^{mn-n-t+1}$, for any $n \in \mathbb{Z}_{\geq 1}$.*

Proof. For $t = m$, consider the family $Z^{\times 1}$ with monodromy datum $(m, N + 2, a')$, where $a'(i) = a(i)$ for $1 \leq i \leq N$, $a'(N + 1) = a'(N + 2) = 0$. Then, the statement follows from Corollary 4.8 applied to $Z^{\times 1}$ for $r = m$.

For $t < m$, consider the family $Z^{\times 1}$ with monodromy datum $(m, N + 2, a')$, where $a'(i) = a(i)$ for $1 \leq i \leq N$, $a'(N + 1) = t$ and $a'(N + 2) = m - t$. The μ -ordinary polygon u' of the associated Shimura variety is $u \oplus \text{ord}^{m-t}$. By Corollary 4.7, $(Z^{\times 1})^\circ[u']$ is non empty. Then, the statement follows from Corollary 4.8 applied to $Z^{\times 1}$ for $r = t$, by observing that $(u')^n \oplus \text{ord}^{(n-1)(t-1)} = u^n \oplus \text{ord}^{mn-n-t+1}$. \square

4.3.4. Double Induction. We next consider inductive systems constructed from a pair of monodromy data satisfying hypotheses (A) and (B). For clarity, we state Corollary 4.10 under the simplifying assumption $m_1 = m_2$. Corollary 9.9 contains an example of this result; it also applies to the pair of monodromy data in the proof of Corollary 9.7.

¹⁰This condition implies that $\overline{Z}(\gamma)$ intersects the boundary component Δ_0 of \mathcal{M}_g .

Corollary 4.10. *Let γ_1 and γ_2 be a pair of monodromy data with $m_1 = m_2$ satisfying hypotheses (A) and (B). Let $r = \gcd(m, a_1(N_1))$ and recall Notation 3.14. Assume that $Z_1^\circ[u_1]$ and $Z_2^\circ[u_2]$ are both non-empty. Then there exists a smooth curve over $\overline{\mathbb{F}}_p$ with Newton polygon $u_1^{n_1} \oplus u_2^{n_2} \oplus \text{ord}^{(n_1+n_2-2)(m-1)+(r-1)}$ for any $n_1, n_2 \in \mathbb{Z}_{\geq 1}$.*

Proof. We apply Corollary 4.9 to the family Z_1 (resp. Z_2) with $t = m$. The result is a family $Z_1^{\times n_1}$ (resp. $Z_2^{\times n_2}$) of smooth curves with Newton polygon $u_1^{n_1} \oplus \text{ord}^{(n_1-1)(m-1)}$ (resp. $u_2^{n_2} \oplus \text{ord}^{(n_2-1)(m-1)}$). Since Z_1 and Z_2 satisfy hypotheses (A) and (B), due to the construction in the proof of Corollary 4.9, $Z_1^{\times n_1}$ and $Z_2^{\times n_2}$ also satisfy hypotheses (A) and (B). The result then follows from Theorem 4.5. \square

5. HYPOTHESIS (B) AND THE μ -ORDINARY NEWTON POLYGON

In this section, we prove Proposition 4.4, namely that hypothesis (B) for a pair of signatures (f_1, f_2) is a necessary and sufficient condition for the associated Shimura variety $\text{Sh}_1 \times \text{Sh}_2$ to intersect the μ -ordinary Newton polygon stratum of Sh_3^* . Recall that σ is Frobenius and \mathfrak{D} is the set of orbits of σ in \mathcal{T} .

Notation 5.1. For a σ -orbit $\mathfrak{o} \in \mathfrak{D}$, let $\mathfrak{p}_\mathfrak{o}$ be the prime above p associated with \mathfrak{o} and let $|\mathfrak{o}|$ be the size of the orbit. For each $\tau \in \mathfrak{o}$, the order of τ in $\mathbb{Z}/m\mathbb{Z}$ is constant and denoted $e_\mathfrak{o}$; by definition, $e_\mathfrak{o} \mid m$. Let $\mathbb{Q}[\mu_m]_{\mathfrak{p}_\mathfrak{o}}$ denote the local field which is the completion of K_e along the prime $\mathfrak{p}_\mathfrak{o}$.

Let \mathcal{X} denote the universal abelian scheme over $\text{Sh} = \text{Sh}(H, \mathfrak{h})^{11}$, and $\mathcal{X}[p^\infty]$ the associated p -divisible group scheme. Let $x \in \text{Sh}(\overline{\mathbb{F}}_p)$ and consider the abelian variety $X := \mathcal{X}_x$. Let $\nu = \nu(X)$ be the Newton polygon of X . We omit the proof of the following.

Lemma 5.2. *The $\mathbb{Q}[\mu_m]$ -action of \mathcal{X} induces a $\mathbb{Q}[\mu_m] \otimes_{\mathbb{Q}} \mathbb{Q}_p$ -action on $\mathcal{X}[p^\infty]$. Thus it induces canonical decompositions*

$$\mathcal{X}[p^\infty] = \bigoplus_{\mathfrak{o} \in \mathfrak{D}} \mathcal{X}[\mathfrak{p}_\mathfrak{o}^\infty] \text{ and } \nu = \bigoplus_{\mathfrak{o} \in \mathfrak{D}} \nu(\mathfrak{o}),$$

where, for each $\mathfrak{o} \in \mathfrak{D}$, the group scheme $\mathcal{X}[\mathfrak{p}_\mathfrak{o}^\infty]$ is a p -divisible $\mathbb{Q}[\mu_m]_{\mathfrak{p}_\mathfrak{o}}$ -module and $\nu(\mathfrak{o}) = \nu(X[\mathfrak{p}_\mathfrak{o}^\infty])$ is its Newton polygon.

For each $\nu \in B = B(H_{\mathbb{Q}_p}, \mu_{\mathfrak{h}})$, we write $\nu(\mathfrak{o})$ for its \mathfrak{o} -component, hence $\nu = \bigoplus_{\mathfrak{o} \in \mathfrak{D}} \nu(\mathfrak{o})$. For all $\nu, \nu' \in B$, note that $\nu \leq \nu'$ if and only if $\nu(\mathfrak{o}) \leq \nu'(\mathfrak{o})$, for all $\mathfrak{o} \in \mathfrak{D}$.

The following lemma says that to prove Proposition 4.4, it suffices to consider each σ -orbit \mathfrak{o} in $\mathcal{T} = \mathbb{Z}/m_3\mathbb{Z}$ separately.

Lemma 5.3. *With the same notation and assumption as in Proposition 4.4: the equality $u_3 = u_1^{d_1} \oplus u_2^{d_2} \oplus \text{ord}^\epsilon$ is equivalent to the system of equalities, for every orbit \mathfrak{o} in \mathcal{T} ,*

$$(5.1) \quad u_3(\mathfrak{o}) = u_1^{d_1}(\mathfrak{o}) \oplus u_2^{d_2}(\mathfrak{o}) \oplus \text{ord}^{\epsilon_\mathfrak{o}},$$

where $\epsilon_\mathfrak{o} := |\mathfrak{o}|$ if $e_\mathfrak{o}$ is a divisor of $d_1 d_2 r_0$ but not a divisor of d_1 or d_2 , and $\epsilon_\mathfrak{o} := 0$ otherwise.

¹¹ or more generally the universal semi-abelian variety over its toroidal compactification,

Proof. Note that $\epsilon = d_1 d_2 r_0 - d_1 - d_2 + 1$ is equal to the number of $\tau \in \mathcal{T}$ whose order is a divisor of $d_1 d_2 r_0$ but not a divisor of d_1 or d_2 ; recall $\gcd(d_1, d_2) = 1$. Thus $\epsilon = \sum_{\mathfrak{o} \in \mathfrak{D}} \epsilon_{\mathfrak{o}}$, and the statement follows from Lemma 5.2 and the discussion below the lemma. \square

Proof of Proposition 4.4. Fix an orbit \mathfrak{o} in \mathcal{T} . For $i = 1, 2$, let u_i^\dagger denote the μ -ordinary Newton polygon of $\text{Sh}(\mu_{m_3}, \mathfrak{f}_i^\dagger)$. By definition, $u_i^\dagger = u_i^{d_i}$, and $u_i^\dagger(\mathfrak{o}) = u_i^{d_i}(\mathfrak{o})$. That is, the Newton polygons $u_i^\dagger(\mathfrak{o})$ and $u_i(\mathfrak{o})$ have the same slopes, with the multiplicity of each slope in $u_i^\dagger(\mathfrak{o})$ being d_i times its multiplicity in $u_i(\mathfrak{o})$. Recall the formulas for the slopes and multiplicities of $u(\mathfrak{o})$ from Section 2.6.1.

Reduction to a combinatorial problem. The formulas for the slopes and multiplicities rely only on the signature type \mathfrak{f} viewed as a \mathbb{N} -valued function on \mathcal{T} , and do not require \mathfrak{f} to be a signature associated with a Shimura variety. We regard each signature type \mathfrak{f} as an \mathbb{N} -valued function on \mathfrak{o} , and denote by $u(\mathfrak{o})$ the Newton polygon defined by the data of slopes in (2.4) and multiplicities in (2.5).¹² We use subscripts and superscripts to identify various \mathbb{N} -valued functions and their Newton polygons, for example, \mathfrak{f}_1^\dagger and $u_1^\dagger(\mathfrak{o})$.

Proposition 4.4 follows from the claim below taking $R = r_0$, using Lemma 5.3.

Claim. *Let R be a positive integer which divides $\gcd(m_1, m_2)$. Set $\delta := \delta_{d, dR} + \delta_{d_1, d_1 d_2} - \delta_{1, d_2}$, with $d := m_3 / \gcd(m_1, m_2) = d_1 d_2$ and notations as in Definition 3.11; set $\epsilon_{\mathfrak{o}} := |\mathfrak{o}|$ if $e_{\mathfrak{o}}$ divides dR but not d_1 or d_2 , and $\epsilon_{\mathfrak{o}} := 0$ otherwise.*

Define $\mathfrak{f}_3 := \mathfrak{f}_1^\dagger + \mathfrak{f}_2^\dagger + \delta$. Then, the equality

$$(5.2) \quad u_3(\mathfrak{o}) = u_1^\dagger(\mathfrak{o}) \oplus u_2^\dagger(\mathfrak{o}) \oplus \text{ord}^{\epsilon_{\mathfrak{o}}}$$

holds if and only if the pair $\mathfrak{f}_1, \mathfrak{f}_2$ satisfies hypothesis (B).

Reduction of claim to the case $d = R = 1$. We first prove that if $\mathfrak{f}_3 = \mathfrak{f}_1^\dagger + \delta$ then

$$(5.3) \quad u_3(\mathfrak{o}) = u_1^\dagger(\mathfrak{o}) \oplus \text{ord}^{\epsilon_{\mathfrak{o}}}.$$

Note that the function $\delta(\tau)$ is constant on \mathfrak{o} , with value $\delta(\mathfrak{o})$ equal to 1 if $e_{\mathfrak{o}}$ divides dR but not d_1 or d_2 , and to 0 otherwise. In particular, $\epsilon_{\mathfrak{o}} = \sum_{\tau \in \mathfrak{o}} \delta(\tau)$, $\delta(\mathfrak{o}) = \delta(\mathfrak{o}^*)$, and by Remark 2.6, $g_3(\mathfrak{o}) = g_1^\dagger(\mathfrak{o}) + 2\delta(\mathfrak{o})$.

If $\delta(\mathfrak{o}) = 0$, then $g_3(\mathfrak{o}) = g_1^\dagger(\mathfrak{o})$, and $\mathfrak{f}_3(\tau) = \mathfrak{f}_1^\dagger(\tau)$ for all $\tau \in \mathfrak{o}$. Hence, $u_3(\mathfrak{o}) = u_1^\dagger(\mathfrak{o})$ which agrees with equality (5.3) for $\epsilon_{\mathfrak{o}} = 0$.

If $\delta(\mathfrak{o}) = 1$, then $g_3(\mathfrak{o}) = g_1^\dagger(\mathfrak{o}) + 2$, and $\mathfrak{f}_3(\tau) = \mathfrak{f}_1^\dagger(\tau) + 1$ for all $\tau \in \mathfrak{o}$. In particular, $g_3(\mathfrak{o}) > \mathfrak{f}_3(\tau) \geq 1$, for all $\tau \in \mathfrak{o}$. By (2.4) and (2.5), both 0 and 1 occur as slopes of $u_3(\mathfrak{o})$, with multiplicity respectively $\rho_3(0) = \rho_1(0) + |\mathfrak{o}|$ and $\rho_3(1) = \rho_1(1) + |\mathfrak{o}|$. Each of the slopes λ of $u_1^\dagger(\mathfrak{o})$, with $\lambda \neq 0, 1$, also occurs for $u_3(\mathfrak{o})$, with multiplicity $\rho_3(\lambda) = \rho_1(\lambda)$. We deduce that $u_3(\mathfrak{o}) = u_1^\dagger(\mathfrak{o}) \oplus \text{ord}^{|\mathfrak{o}|}$, which agrees with equality (5.3) for $\epsilon_{\mathfrak{o}} = |\mathfrak{o}|$.

Equality (5.3) is equivalent to the claim for $\mathfrak{f}_2^\dagger(\tau) = 0$, for all $\tau \in \mathfrak{o}$. Indeed, hypothesis (B) for the pair $(\mathfrak{f}_1^\dagger, 0)$ holds trivially, and equality (5.2) for $\mathfrak{f}_3 = \mathfrak{f}_1^\dagger + \delta$ specializes to (5.3).

By (5.3), we deduce that equality (5.2) holds if and only if it holds in the special case of $d = R = 1$, that is if $u_3(\mathfrak{o}) = u_1^\dagger(\mathfrak{o}) + u_2^\dagger(\mathfrak{o})$ when $\mathfrak{f}_3 = \mathfrak{f}_1^\dagger + \mathfrak{f}_2^\dagger$. By definition, hypothesis (B) holds for the pair $(\mathfrak{f}_1^\dagger, \mathfrak{f}_2^\dagger)$ if and only if it holds for $(\mathfrak{f}_1^\dagger + \delta, \mathfrak{f}_2^\dagger)$. Hence, without loss of

¹²the integer $g(\mathfrak{o})$ will be specified in each part of the proof.

generality, we may assume $d = R = 1$ and $f_3 = f_1 + f_2$. In this case, $g_3(\mathfrak{o}) = g_1(\mathfrak{o}) + g_2(\mathfrak{o})$ by Remark 2.6 and the claim reduces to the following:

Specialized claim: Assume $f_3 = f_1 + f_2$. Then, the equality

$$(5.4) \quad u_3(\mathfrak{o}) = u_1(\mathfrak{o}) \oplus u_2(\mathfrak{o}).$$

holds if and only if the pair f_1, f_2 satisfies hypothesis (B).

Converse direction: assume hypothesis (B). We shall prove that the equality (5.4) holds, by induction on the integer $s_3 + 1$, the number of distinct slopes of $u_3(\mathfrak{o})$. More precisely, we shall proceed as follows. First, we shall establish the base case of induction, for $s_3 = 0$. Next, we shall prove the equality of multiplicities

$$(5.5) \quad \rho_3(\lambda) = \rho_1(\lambda) + \rho_2(\lambda)$$

for $\lambda = \lambda_3(0)$ the first (smallest) slope of $u_3(\mathfrak{o})$. Then, we shall assume $s_3 \geq 1$ and show that the inductive hypothesis and equality (5.5) imply equality (5.4), which will complete the argument. In the induction process, $g_i(\mathfrak{o})$ remains unchanged.

Base case: Assume $s_3 = 0$. Then, for all $\tau \in \mathfrak{o}$, either $f_3(\tau) = g_3(\mathfrak{o})$ or $f_3(\tau) = 0$. The equalities $f_3(\tau) = f_1(\tau) + f_2(\tau)$ and $g_3(\mathfrak{o}) = g_1(\mathfrak{o}) + g_2(\mathfrak{o})$ imply that $f_3(\tau) = g_3(\mathfrak{o})$ (resp. $f_3(\tau) = 0$) if and only if $f_i(\tau) = g_i(\mathfrak{o})$ (resp. $f_i(\tau) = 0$) for both $i = 1, 2$. We deduce that both hypothesis (B) and (5.4) hold in this situation.

Equality (5.5). For $i = 1, 2, 3$, let $E_i(\max)$ denote the maximal value of f_i on \mathfrak{o} . By definition, $E_i(\max)$ is equal to either $E_i(0)$ or $E_i(1)$. In the first case, $\lambda_i(0) > 0$; in the second case, $\lambda_i(0) = 0$. We claim that hypothesis (B) implies $E_3(\max) = E_1(\max) + E_2(\max)$. First, note that hypothesis (B) implies that, for $\omega, \tau \in \mathfrak{o}$:

$$(5.6) \quad f_3(\omega) = f_3(\tau) \text{ if and only if } f_i(\omega) = f_i(\tau) \text{ for both } i = 1, 2.$$

For $i = 1, 2, 3$, set $S_i = \{\tau \in \mathfrak{o} \mid f_i(\tau) = E_i(\max)\}$. Then, property (5.6) implies that $S_3 = S_1 \cap S_2$, which in turn implies $E_3(\max) = E_1(\max) + E_2(\max)$.

We claim that hypothesis (B) implies that $S_3 = S_i$ for some $i \in \{1, 2\}$. Without loss of generality, assume that S_2 properly contains S_3 , and let $\tau_0 \in S_2 - S_3$. Then $\tau_0 \notin S_1$. For any $\omega \in S_1$: $f_1(\omega) = E_1(\max) > f_1(\tau_0)$. Thus, by hypothesis (B), $f_2(\omega) \geq f_2(\tau_0) = E_2(\max)$. We deduce that $f_2(\omega) = E_2(\max)$, hence $S_1 \subseteq S_2$.

By the formulas for slopes (2.4), the equality $S_1 = S_3$ implies that $\lambda_1(0) = \lambda_3(0)$, and the inclusion $S_1 \subseteq S_2$ implies $\lambda_1(0) \leq \lambda_2(0)$ (and the equality holds if and only if $S_2 = S_3$).

For $i = 1, 2, 3$, let $E_i(\text{next})$ denote the maximal value of f_i on $\mathfrak{o} - S_3$. For $i = 1, 3$, $E_i(\text{next}) < E_i(\max)$; for $i = 2$, $E_2(\text{next}) \leq E_2(\max)$ and the equality holds if and only if S_2 properly contains S_3 .

As before by property (5.6), we deduce that hypothesis (B) implies $E_3(\text{next}) = E_1(\text{next}) + E_2(\text{next})$. By the formulas for multiplicities (2.5), the two identities, $E_3(\max) = E_1(\max) + E_2(\max)$ and $E_3(\text{next}) = E_1(\text{next}) + E_2(\text{next})$, imply the desired equality (5.5).

Assume $s_3 \geq 1$. Then the polygon $u_3(\mathfrak{o})$ has at least two distinct slopes. Our plan is to introduce auxiliary functions $\tilde{f}_1(\tau), \tilde{f}_2(\tau)$ such that polygon $\tilde{u}_3(\mathfrak{o})$ associated with the function $\tilde{f}_3(\tau) := \tilde{f}_1(\tau) + \tilde{f}_2(\tau)$ has s_3 distinct slopes.

For $i = 1, 2$, define $\tilde{f}_i(\tau) := f_i(\tau)$ for all $\tau \notin S_3$ and $\tilde{f}_i(\tau) := E_i(\text{next})$ for $\tau \in S_3$. Note that $\tilde{f}_2 = f_2$ unless $S_2 = S_3$. By definition, for $i = 1, 2, 3$, $\tilde{E}_i(\max) = E_i(\text{next})$.

For $i = 1, 3$, and for $i = 2$ if $S_2 = S_3$, the polygon $\tilde{u}_i(\mathfrak{o})$ shares the same slopes as $u_i(\mathfrak{o})$ except $\lambda_i(0)$ which no longer occurs. For each $t = 2, \dots, s_i$, the slope $\lambda_i(t)$ occurs in $\tilde{u}_i(\mathfrak{o})$

with multiplicity equal to $\rho_i(\lambda_i(t))$; while the slope $\lambda_i(1)$ occurs in $\tilde{u}_i(\mathfrak{o})$ with multiplicity $\rho_i(\lambda_i(0)) + \rho_i(\lambda_i(1))$. For $i = 2$, if $S_2 \neq S_3$, then $\tilde{u}_2(\mathfrak{o}) = u_2(\mathfrak{o})$.

Note that \tilde{u}_3 has exactly s_3 slopes. Hence, by the inductive hypothesis, we deduce that $\tilde{u}_3(\mathfrak{o}) = \tilde{u}_1(\mathfrak{o}) \oplus \tilde{u}_2(\mathfrak{o})$. This identity, together with (5.5) and the above computation of $\tilde{\rho}_i(\lambda_i(1))$, implies that $u_3(\mathfrak{o}) = u_1(\mathfrak{o}) \oplus u_2(\mathfrak{o})$.

Forward direction: assume (5.4). We shall prove that the pair (f_1, f_2) satisfies hypothesis (B), arguing by contradiction. Supposing hypothesis (B) does not hold, we shall define auxiliary functions \bar{f}_1, \bar{f}_2 obtained by precomposing f_1, f_2 with a permutation of \mathfrak{o} , such that the Newton polygon $\bar{u}_3(\mathfrak{o})$ associated with the function $\bar{f}_3(\tau) := \bar{f}_1(\tau) + \bar{f}_2(\tau)$ is strictly above $u_3(\mathfrak{o})$, i.e., $\bar{u}_3(\mathfrak{o}) < u_3(\mathfrak{o})$. By the formulas for slopes and multiplicities (2.4) and (2.5), we see that precomposing a function f with a permutation of \mathfrak{o} does not change the associated polygon $u(\mathfrak{o})$. Hence, for $i = 1, 2$ we deduce $\bar{u}_i(\mathfrak{o}) = u_i(\mathfrak{o})$. On the other hand, by repeating the permutation process, we eventually end up with a pair (\bar{f}_1, \bar{f}_2) which satisfies hypothesis (B) and by the above argument, hypothesis (B) implies that $\bar{u}_1(\mathfrak{o}) \oplus \bar{u}_2(\mathfrak{o}) = \bar{u}_3(\mathfrak{o})$. Hence $\bar{u}_3(\mathfrak{o}) = u_1(\mathfrak{o}) \oplus u_2(\mathfrak{o})$, which equals $u_3(\mathfrak{o})$ by (5.2) and contradicts the conclusion that $\bar{u}_3(\mathfrak{o}) < u_3(\mathfrak{o})$.

Contradict hypothesis (B). Thus, there exist $\omega_0, \eta_0 \in \mathfrak{o}$ such that

$$f_2(\omega_0) > f_2(\eta_0) \text{ and } f_1(\omega_0) < f_1(\eta_0).$$

Let γ denote the permutation of \mathfrak{o} which switches ω_0 and η_0 , and define $\bar{f}_1(\tau) := f_1(\gamma(\tau))$ and $\bar{f}_2(\tau) := f_2(\tau)$. Set $\bar{f}_3(\tau) := \bar{f}_1(\tau) + \bar{f}_2(\tau)$. Then, $\bar{f}_3(\tau) = f_3(\tau)$ except for $\tau = \omega_0, \eta_0$. Note that $\bar{f}_3(\omega_0) > f_3(\omega_0)$, $\bar{f}_3(\eta_0) < f_3(\eta_0)$, and $\bar{f}_3(\omega_0) > f_3(\eta_0)$ (also, $\bar{f}_3(\eta_0) < f_3(\omega_0)$).

We claim that $\bar{u}_3(\mathfrak{o}) < u_3(\mathfrak{o})$, meaning that $\bar{u}_3(\mathfrak{o})$ and $u_3(\mathfrak{o})$ share the same endpoints (this follows from the equality $\bar{g}_3(\mathfrak{o}) = g_3(\mathfrak{o})$), and that $\bar{u}_3(\mathfrak{o})$ lies strictly above $u_3(\mathfrak{o})$.

We first show that possibly after sharing the first several slopes, $\bar{u}_3(\mathfrak{o})$ admits a slope which is strictly larger than the corresponding one in $u_3(\mathfrak{o})$. Let us consider the value $A = \bar{f}_3(\omega_0)$. Note that $A > \bar{f}_3(\omega_0) \geq 0$. If $A = f_3(\tau)$ for some $\tau \in \mathfrak{o}$, say $A = E_3(t)$ for some $t \in \{0, \dots, s_3\}$. Then by the formulas for slopes and multiplicities (2.4) and (2.5), we deduce that the first t slopes, and their multiplicities, of $\bar{u}_3(\mathfrak{o})$ and $u_3(\mathfrak{o})$ agree, but the $(t+1)$ -st slope of $\bar{u}_3(\mathfrak{o})$ is strictly larger than the $(t+1)$ -st slope of $u_3(\mathfrak{o})$. If $f_3(\tau) \neq A$ for all $\tau \in \mathfrak{o}$, let $t \in \{0, \dots, s_3\}$ be such that $E_3(t) > A > E_3(t+1)$. We deduce that the first t slopes of $\bar{u}_3(\mathfrak{o})$ and $u_3(\mathfrak{o})$ agree, and so do the multiplicities of the first $t-1$ slopes, but the multiplicity of the t -th slope of \bar{u}_3 is strictly smaller than that of $u_3(\mathfrak{o})$.

By similar arguments for the subsequent slopes and multiplicities, $\bar{u}_3(\mathfrak{o})$ never drops strictly below $u_3(\mathfrak{o})$, but it might (and often does) agree with $u_3(\mathfrak{o})$ for large slopes. \square

6. THE TORELLI LOCUS AND THE NON μ -ORDINARY LOCUS OF SHIMURA VARIETIES

In this section, we study the intersection of the open Torelli locus with Newton polygon strata which are not μ -ordinary in PEL-type Shimura varieties. The main result, Theorem 6.11, provides a method to leverage information from smaller dimension to larger dimension. This theorem is significantly more difficult than Theorem 4.5; we add an extra compatibility condition to maintain control over the codimensions of the Newton polygon strata. This is the first systematic result on this topic that we are aware of.

For applications, we find situations in which Theorem 6.11 can be applied infinitely many times; from this, we produce systems of infinitely many PEL-type Shimura varieties for which we can verify that the open Torelli locus intersects certain non μ -ordinary Newton polygon strata. See Corollaries 6.14 to 6.16 and Section 9 for details.

Notation 6.1. Let $\gamma_1 = (m_1, N_1, a_1)$, $\gamma_2 = (m_2, N_2, a_2)$ be an ordered pair of (generalized) monodromy data which satisfies hypothesis (A). Assume that $m_1 | m_2$. Set $d := m_2/m_1$ and $r := \gcd(m_1, a_1(N_1))$. Then, (3.1) specializes to $\epsilon = d(r-1)$ and $g_3 = dg_1 + g_2 + \epsilon$. In particular, $\epsilon = 0$ if and only if $r = 1$.

Remark 6.10 explains why we restrict to the case $m_1 | m_2$.

6.1. Hypothesis (C). To study the Newton polygons beyond the μ -ordinary case, we introduce an extra hypothesis.

Definition 6.2. Given a Newton polygon ν , the *first slope* $\lambda_{1st}(\nu)$ is the smallest slope of ν and the *last slope* $\lambda_{last}(\nu)$ is the largest slope of ν . If ν is symmetric with q distinct slopes, the *middle slope* $\lambda_{mid}(\nu)$ is the $\lfloor \frac{q+1}{2} \rfloor$ -st slope of ν .

Definition 6.3. An ordered pair of monodromy data γ_1, γ_2 is *compatible* if the slopes of the μ -ordinary Newton polygons u_1 and u_2 satisfy:

hypothesis (C): for each orbit $\mathfrak{o} \in \mathfrak{D}$, every slope of $u_1(\mathfrak{o})$ is in the range

$$[0, 1] \setminus (\lambda_{1st}(u_2(\mathfrak{o})), \lambda_{last}(u_2(\mathfrak{o}))).$$

By convention, the condition in the previous line holds for \mathfrak{o} if $u_i(\mathfrak{o})$ is empty for either $i = 1, 2$. If a pair of monodromy data is compatible, then we write $u_1 \ll_{(C)} u_2$.

Remark 6.4. (1) Hypothesis (C) holds for \mathfrak{o} if either $u_1(\mathfrak{o})$ has slopes in $\{0, 1\}$ or $u_2(\mathfrak{o})$ is supersingular. In particular, if u is ordinary or supersingular, then $u \ll_{(C)} u$.

(2) If $\mathfrak{o} = \mathfrak{o}^*$, then hypothesis (C) holds for \mathfrak{o} if and only if $\lambda_{mid}(u_1(\mathfrak{o})) \leq \lambda_{1st}(u_2(\mathfrak{o}))$.

(3) If $u_1 \ll_{(C)} u_2$ then $u_1^d \ll_{(C)} u_2$ for all $d \in \mathbb{Z}_{\geq 1}$.

Remark 6.5. Let γ_1, γ_2 be a pair of monodromy data as in Notation 6.1. By Section 2.6.1,

$$\lambda_{1st}(u_2(\mathfrak{o})) = \frac{1}{|\mathfrak{o}|} \#\{\tau \in \mathfrak{o} \mid f_2(\tau) = g_2(\mathfrak{o})\} \text{ and } \lambda_{last}(u_2(\mathfrak{o})) = \frac{1}{|\mathfrak{o}|} \#\{\tau \in \mathfrak{o} \mid f_2(\tau) > 0\}.$$

Hypothesis (C) holds for \mathfrak{o} if and only if there exists an integer $E(\mathfrak{o}) \in [0, g_1(\mathfrak{o})]$ such that

$$\begin{aligned} \#\{\tau \in \mathfrak{o} \mid f_1^\dagger(\tau) > E(\mathfrak{o})\} &\leq \#\{\tau \in \mathfrak{o} \mid f_2(\tau) = g_2(\mathfrak{o})\}, \text{ and} \\ \#\{\tau \in \mathfrak{o} \mid f_1^\dagger(\tau) \geq E(\mathfrak{o})\} &\geq \#\{\tau \in \mathfrak{o} \mid f_2(\tau) > 0\}. \end{aligned}$$

The next statement follows from Definition 6.3 and Remark 6.5.

Lemma 6.6. *The following are equivalent: $u \ll_{(C)} u$; for each $\mathfrak{o} \in \mathfrak{D}$, the Newton polygon $u(\mathfrak{o})$ has at most two distinct slopes; and, for each $\mathfrak{o} \in \mathfrak{D}$, there exists an integer $E(\mathfrak{o}) \in [0, g(\mathfrak{o})]$ such that $f(\tau) \in \{0, E(\mathfrak{o}), g(\mathfrak{o})\}$ for all $\tau \in \mathfrak{o}$.*

Unlike hypothesis (B), hypothesis (C) does not behave well under induction in general. Lemma 6.7 identifies two instances when it does. We omit the proof.

Lemma 6.7. (1) *If $u_1 \ll_{(C)} u_2$ then $u_1^n \oplus \text{ord}^l \ll_{(C)} u_2$, for any $n, l \in \mathbb{Z}_{\geq 1}$.*

(2) *If $u_1 \ll_{(C)} u_2$ and $u_2 \ll_{(C)} u_2$, then $u_1^n \oplus u_2^m \oplus \text{ord}^l \ll_{(C)} u_2$, for any $n, m, l \in \mathbb{Z}_{\geq 1}$.*

6.2. The significance of hypothesis (C). Hypothesis (C) is sufficient to prove the geometric condition on the Newton polygon stratification in Proposition 6.8 below. We use hypothesis (C) to prove the surjectivity of the map in (6.2).

Proposition 6.8. *Let $\gamma_1 = (m_1, N_1, a_1), \gamma_2 = (m_2, N_2, a_2)$ be an ordered pair of monodromy data as in Notation 6.1. Assume the pair satisfies hypotheses (A), (B), and (C). Consider the monodromy datum γ_3 as in Definition 3.5. Then, for any Newton polygon $v_2 \in B_2$,*

$$(6.1) \quad \text{codim}(\text{Sh}_2[v_2], \text{Sh}_2) = \text{codim}(\overline{\text{Sh}}_3[u_1^d \oplus v_2 \oplus \text{ord}^\epsilon], \overline{\text{Sh}}_3).$$

The following lemma is a reformulation of Proposition 4.4.

Lemma 6.9. *Hypothesis (B) is equivalent to the assumption that (6.1) holds for $v_2 = u_2$.*

In fact, the proof below shows that if hypothesis (B) does not hold, then (6.1) is false for all $v_2 \in B_2$.

Proof of Proposition 6.8. For any Kottwitz set B and any $v \in B$, let $B(v) = \{t \in B \mid t \geq v\}$.

We first prove the case when $\epsilon = 0$. Consider the map $\Sigma : B_2 \rightarrow B_3$, where $t \mapsto u_1^d \oplus t$.

We first note that Σ is an order-preserving injection.¹³ Let $t \in B_2$. For each orbit \mathfrak{o} in \mathfrak{D} , let $q_1 = q_1(\mathfrak{o})$ (resp. $q'_1 = q'_1(\mathfrak{o})$) be the number of distinct slopes of $u_1(\mathfrak{o})$ in $[0, \lambda_{1st}(u_2(\mathfrak{o}))]$ (resp. $[\lambda_{last}(u_2(\mathfrak{o})), 1]$). By hypothesis (C), for the Newton polygon $u_1^d \oplus t$, the first q_1 and the last q'_1 slopes of $u_1(\mathfrak{o})^d \oplus t(\mathfrak{o})$ are the slopes of $u_1(\mathfrak{o})^d$ with the same multiplicities and the rest are the slopes of $t(\mathfrak{o})$ with the same multiplicities. So, if $t \leq t'$ in B_2 , then $u_1^d \oplus t \leq u_1^d \oplus t'$ in B_3 . In particular, the map Σ induces an injection on the ordered sets

$$(6.2) \quad B_2(v_2) \rightarrow B_3(u_1^d \oplus v_2), \quad t \mapsto u_1^d \oplus t.$$

By (2.6), to conclude, it suffices to prove that under hypotheses (B) and (C) the map in (6.2) is also surjective. By Proposition 4.4, hypothesis (B) implies that $u_3 = u_1^d \oplus u_2$. Hence, for a Newton polygon $v \in B_3(u_1^d \oplus v_2)$, then $u_1^d \oplus v_2 \leq v \leq u_3 = u_1^d \oplus u_2$. By the paragraph after Lemma 5.2,

$$u_1(\mathfrak{o})^d \oplus v_2(\mathfrak{o}) \leq v(\mathfrak{o}) \leq u_1(\mathfrak{o})^d \oplus u_2(\mathfrak{o}).$$

By hypothesis (C), the inequalities above imply that $v(\mathfrak{o})$ and $u_1(\mathfrak{o})^d$ share the first q_1 and last q'_1 slopes, with the same multiplicities except for the q_1 -th slope (resp. q_2 -th slope) which may occur with higher multiplicity in the former if $u_1(\mathfrak{o})$ has slope $\lambda_{1st}(u_2(\mathfrak{o}))$ (resp. $\lambda_{last}(u_2(\mathfrak{o}))$). We deduce that each $v \in B_3(u_1^d \oplus v_2)$ is of the form $v = u_1^d \oplus t$ for some $t \in B_2(v_2)$; thus the map in (6.2) is surjective.

If $\epsilon \neq 0$, by (2.7), the same argument still applies, with u_1^d replaced by $u_1^d \oplus \text{ord}^\epsilon$. \square

Remark 6.10. Let γ be a monodromy datum, of signature \mathfrak{f} . For $d > 1$, let γ^\dagger be the induced monodromy datum, of signature \mathfrak{f}^\dagger , as in Notation 3.1. The map $(\cdot)^\dagger : B(\mathfrak{f}) \rightarrow B(\mathfrak{f}^\dagger)$, $v \mapsto v^d$, is injective and order-preserving, but is not surjective in general. If $(\cdot)^\dagger$ is not surjective, then, by (2.6), there exists $v \in B$ such that $\text{codim}(\text{Sh}[v], \text{Sh}) \neq \text{codim}(\text{Sh}^\dagger[v^d], \text{Sh}^\dagger)$. For example, if $\mathfrak{f} = (1)$, and $d = 2$, this happens when $v = (1/2, 1/2)$.

¹³For convenience, we use hypothesis (C) to construct the order-preserving map (6.2); however, this part can be proved without using this hypothesis.

6.3. The second main result. The next result also provides a partial positive answer to Conjecture 1.1 when $\epsilon = 0$.

Theorem 6.11. *Let γ_1, γ_2 be an ordered pair of monodromy data as in Notation 6.1. Assume it satisfies hypotheses (A), (B), and (C). Let $\epsilon = d(r - 1)$. Consider the monodromy datum γ_3 from Definition 3.5. Let $v_2 \in B_2$. If $Z_1^\circ[u_1]$ and $Z_2^\circ[v_2]$ are non-empty, and $Z_2^\circ[v_2]$ contains an irreducible component Γ_2 such that*

$$(6.3) \quad \text{codim}(\Gamma_2, Z_2) = \text{codim}(\text{Sh}_2[v_2], \text{Sh}_2),$$

then $Z_3^\circ[u_1^d \oplus v_2 \oplus \text{ord}^\epsilon]$ is non-empty and contains an irreducible component Γ_3 such that

$$\text{codim}(\Gamma_3, Z_3) = \text{codim}(\text{Sh}_3[u_1^d \oplus v_2 \oplus \text{ord}^\epsilon], \text{Sh}_3).$$

Remark 6.12. As seen in Section 6.5, hypothesis (C) is not a necessary condition and it can occasionally be removed. Specifically, Theorem 6.11 still holds with hypothesis (C) replaced by the weaker (but harder to verify) assumption that (6.1) holds for the given non μ -ordinary Newton polygon $v_2 \in B_2$.

Remark 6.13. If Z_2 is one of Moonen's special families from [25] and $Z_2^\circ[v_2]$ is non-empty, then every irreducible component of $Z_2^\circ[v_2]$ satisfies the codimension condition (6.3).

Proof of Theorem 6.11. By Remark 2.3, without loss of generality, we may assume that the inertia types a_1 and a_2 contain no zero entries if $r < m_1$ and no zero entries other than $a_1(N_1) = a_2(1) = 0$ if $r = m_1$. Set $l := \text{codim}(\text{Sh}_2[v_2], \text{Sh}_2)$. Recall the clutching morphism $\kappa : \tilde{Z}_1 \times \tilde{Z}_2 \rightarrow \tilde{Z}_3$ from Proposition 3.9. Note that

$$(6.4) \quad \dim(Z_3) = N_3 - 3 = (N_1 - 3) + (N_2 - 3) + 1.$$

We distinguish three cases: $\epsilon = 0$, $\epsilon \neq 0$ and $r < m_1$, and $\epsilon \neq 0$ and $r = m_1$.

Assume $\epsilon = 0$. Then (6.4) implies that

$$(6.5) \quad \dim(Z_3) = \dim(Z_1) + \dim(Z_2) + 1 = \dim(\tilde{Z}_1) + \dim(\tilde{Z}_2) + 1.$$

By Notation 3.4, $g_3 = dg_1 + g_2$ and the formula for f_3 is in (3.3). The clutching morphism κ is compatible with the morphism $\iota : \text{Sh}_1 \times \text{Sh}_2 \rightarrow \text{Sh}_3$ given by $\iota(\mathcal{X}_1, \mathcal{X}_2) = \mathcal{X}_1^d \oplus \mathcal{X}_2$. Since the map $\tilde{Z}_i \rightarrow \bar{Z}_i$ is finite, $\dim(\tilde{Z}_i[v_i]) = \dim(\bar{Z}_i[v_i])$ for any $v_i \in B_i$.

By Proposition 6.8, $l = \text{codim}(\text{Sh}_3[u_1^d \oplus v_2], \text{Sh}_3)$. Let $\tilde{\Gamma}_2$ denote the Zariski closure of the preimage of Γ_2 in $\tilde{Z}_2[v_2]$. A priori, $\tilde{\Gamma}_2$ may not be irreducible, in which case we replace it by one of its irreducible components. Then $\dim(\tilde{\Gamma}_2) = \dim(\Gamma_2)$.

Let $W := \kappa(\tilde{Z}_1[u_1], \tilde{Z}_2[v_2])$. Since $W \subseteq \tilde{Z}_3[u_1^d \oplus v_2]$, then $\tilde{Z}_3[u_1^d \oplus v_2]$ is non-empty. By (6.5), $\kappa(\tilde{Z}_1, \tilde{Z}_2)$ has codimension 1 in \tilde{Z}_3 . In addition, W is an open and closed substack of the intersection of $\kappa(\tilde{Z}_1, \tilde{Z}_2)$ and $\tilde{Z}_3[u_1^d \oplus v_2]$. By [31, page 614], every irreducible component of W has codimension at most 1 in the irreducible component of $\tilde{Z}_3[u_1^d \oplus v_2]$ which contains it. Note that $\kappa(\tilde{Z}_1[u_1], \tilde{\Gamma}_2)$ is an irreducible component of W . Let $\tilde{\Gamma}_3$ be the irreducible component of $\tilde{Z}_3[u_1^d \oplus v_2]$ which contains $\kappa(\tilde{Z}_1[u_1], \tilde{\Gamma}_2)$. It follows that $\text{codim}(\kappa(\tilde{Z}_1[u_1], \tilde{\Gamma}_2), \tilde{\Gamma}_3) \leq 1$. So

$$\dim(\tilde{\Gamma}_3) = \begin{cases} \dim(Z_1) + \dim(Z_2) - l & \text{if } \tilde{\Gamma}_3 = \kappa(\tilde{Z}_1[u_1], \tilde{\Gamma}_2), \\ \dim(Z_1) + \dim(Z_2) - l + 1 & \text{otherwise.} \end{cases}$$

On the other hand, for all $b \in B_3$, by (2.6) and the de Jong–Oort purity theorem [7, Theorem 4.1], the codimension of any irreducible component of $\tilde{Z}_3[b]$ in \tilde{Z}_3 is no greater than $\text{length}(b) = \text{codim}(\text{Sh}_3[b], \text{Sh}_3)$. For $b = u_1^d \oplus v_2$, by (6.5), this yields

$$\dim(\tilde{\Gamma}_3) \geq \dim(Z_3) - l = \dim(Z_1) + \dim(Z_2) + 1 - l.$$

We deduce that $\text{codim}(\tilde{\Gamma}_3, \tilde{Z}_3) = l$ and that $\tilde{\Gamma}_3$ strictly contains $\kappa(\tilde{Z}_1[u_1], \tilde{\Gamma}_2)$.

Let $\bar{\Gamma}_3$ denote the image of $\tilde{\Gamma}_3$ via the forgetful map $\tilde{Z}_3 \rightarrow \bar{Z}_3$. Define $\Gamma_3 = \bar{\Gamma}_3 \cap Z_3^\circ$. To finish the proof, we only need to show that Γ_3 is non-empty. Therefore, it suffices to show that $\tilde{\Gamma}_3$ is not contained in the image of any other clutching map from Proposition 3.9. Since $r = 1$, by Proposition 3.10 the points in W represent curves of compact type, thus $\tilde{\Gamma}_3 \cap \tilde{Z}_3^c$ is non empty.

To finish, we argue by contradiction; suppose $\tilde{\Gamma}_3$ is contained in the image of any of the other clutching maps in \tilde{Z}_3^c . This would imply that all points of $\kappa(\tilde{Z}_1[u_1], \tilde{\Gamma}_2)$ represent μ_m -covers of a curve of genus 0 comprised of at least 3 projective lines. This is only possible if all points of either $\tilde{Z}_1[u_1]$ or $\tilde{\Gamma}_2$ represent μ_m -covers of a curve of genus 0 comprised of at least 2 projective lines. This would imply that either $Z_1^\circ[u_1]$ or $\Gamma_2 \subset Z_2^\circ[v_2]$ is empty, which contradicts the hypotheses of the theorem.

Assume $\epsilon \neq 0$ and $r < m_1$. By the same argument as when $\epsilon = 0$, there exists an irreducible component $\tilde{\Gamma}_3$ of $\tilde{Z}_3[u_1^d \oplus v_2 \oplus \text{ord}^\epsilon]$ of codimension l such that $\tilde{\Gamma}_3$ strictly contains $\kappa(\tilde{Z}_1[u_1], \tilde{\Gamma}_2)$. To finish the proof, we only need to show that $\tilde{\Gamma}_3$ is not contained in the boundary of $\tilde{\mathcal{M}}_{\mu_m}$. As before, $\tilde{\Gamma}_3$ is not contained in the image of any of the clutching maps in \tilde{Z}_3^c . Suppose that $\tilde{\Gamma}_3$ is contained in the image of any of the clutching maps not in \tilde{Z}_3^c . By keeping careful track of the toric rank, one can check that this implies that the points of either $\tilde{Z}_1[u_1]$ or $\tilde{\Gamma}_2$ represent μ_m -covers of curves that are not of compact type. This would imply that either $Z_1^\circ[u_1]$ or $\Gamma_2 \subset Z_2^\circ[v_2]$ is empty, which contradicts the hypotheses of the theorem.

Assume $\epsilon \neq 0$ and $r = m_1$. By Remark 2.3, for $i = 1, 2$, the fibers of the forgetful map $f_i : \tilde{Z}_i^\circ \rightarrow Z_i^\circ$ have pure dimension 1. Let $\tilde{\Gamma}'_2$ be an irreducible component of the preimage via f_2 of Γ_2 ; it is in $\tilde{Z}_2^\circ[v_2]$. Let $\tilde{\Gamma}'_1$ be an irreducible component of the preimage via f_1 of $Z_1^\circ[u_1]$; it is in $\tilde{Z}_1^\circ[u_1]$. Then $\dim(\tilde{\Gamma}'_2) = \dim(\Gamma_2) + 1$. Similarly, $\dim(\tilde{\Gamma}'_1) = \dim(Z_1[u_1]) + 1$.

Let $\tilde{\Gamma}_3$ be the irreducible component of $\tilde{Z}_3[u_1^d \oplus v_2 \oplus \text{ord}^\epsilon]$ that contains the image $\kappa(\tilde{\Gamma}'_1, \tilde{\Gamma}'_2)$. As before, $\dim(\tilde{\Gamma}_3) \geq \dim(\kappa(\tilde{\Gamma}'_1, \tilde{\Gamma}'_2)) + 1$. The rest of the proof follows in the same way as when $r < m_1$, by taking $\Gamma_3 = \bar{\Gamma}_3 \cap Z_3^\circ$, where $\bar{\Gamma}_3$ is the image of $\tilde{\Gamma}_3$ via the forgetful map. To obtain the dimension inequality, note that

$$\dim(\kappa(\tilde{\Gamma}'_1, \tilde{\Gamma}'_2)) = 1 + \dim(Z_1[u_1]) + 1 + \dim(\Gamma_2) = 2 + \dim(Z_1) + \dim(Z_2) - l,$$

where $l = \text{codim}(\text{Sh}_2[v_2], \text{Sh}_2)$. In this case, $\dim(Z_i) = N_i - 4$ for $i = 1, 2$. By (6.4),

$$(6.6) \quad \dim(Z_3) = (N_1 - 4) + (N_2 - 4) + 3 = \dim(Z_1) + \dim(Z_2) + 3.$$

On the other hand, by the de Jong–Oort purity theorem [7, Theorem 4.1],

$$\dim(\tilde{\Gamma}_3) \geq \dim(Z_3) - l = \dim(Z_1) + \dim(Z_2) + 3 - l = \dim(\kappa(\tilde{\Gamma}'_1, \tilde{\Gamma}'_2)) + 1. \quad \square$$

6.4. Infinite clutching for non μ -ordinary. This section is similar to Section 4.3, in that we find situations in which Theorem 6.11 can be implemented recursively, infinitely many times, except that we now focus on non μ -ordinary Newton polygons.

Let $\gamma = (m, N, a)$ be a monodromy datum and let $\nu \in B(\gamma)$.

Corollary 6.14. *(Extension of Corollary 4.7) Assume $Z^\circ(\gamma)[\nu]$ is non-empty and contains an irreducible component Γ such that $\text{codim}(\Gamma, Z(\gamma)) = \text{codim}(\text{Sh}[\nu], \text{Sh})$. Then for any n in the semi-group generated by $\{m - t : t \mid m\}$, there exists a μ_m -cover $C \rightarrow \mathbb{P}^1$ over $\overline{\mathbb{F}}_p$ where C is a smooth curve with Newton polygon $\nu \oplus \text{ord}^n$.*

Proof. Let γ_1 be as in the proof of Corollary 4.7. Note that $u_1(\mathfrak{o})$ is empty for all \mathfrak{o} . So the pair γ_1, γ satisfies hypothesis (C), in addition to (A) and (B). The proof is then the same as for Corollary 4.7, replacing Theorem 4.5 with Theorem 6.11. \square

Corollary 6.15. *(Extension of Corollaries 4.8 and 4.9) Let $\epsilon = (n - 1)(r - 1)$ if there exist $1 \leq i < j \leq N$ such that $a(i) + a(j) \equiv 0 \pmod{m}$, and $\epsilon = (n - 1)(m - 1)$ otherwise. Assume $Z^\circ(\gamma)[\nu]$ is non-empty and contains an irreducible component Γ such that $\text{codim}(\Gamma, Z(\gamma)) = \text{codim}(\text{Sh}[\nu], \text{Sh})$. Assume $u \ll_{(C)} u$. Then for any $n \in \mathbb{Z}_{\geq 1}$, there exists a smooth curve with Newton polygon $u^{n-1} \oplus \nu \oplus \text{ord}^\epsilon$.*

Proof. The result is true when $n = 1$ by hypothesis. For $n \geq 2$, we use Corollary 4.8 (resp. Corollary 4.9 with $t = m$) to construct a family $Z^{\times n-1}$ with Newton polygon $u^{n-1} \oplus \text{ord}^{(n-2)(r-1)}$ (resp. $u^{n-1} \oplus \text{ord}^{(n-2)(m-1)}$). The pair of monodromy data of the families $Z^{\times n-1}$ and Z satisfies hypotheses (A) and (B). Since $u \ll_{(C)} u$, by Lemma 6.7 (1), the pair also satisfies hypothesis (C). Hence we conclude by Theorem 6.11. \square

Corollary 6.16. *With notation and hypotheses as in Corollary 4.10, assume furthermore that for some $\nu_2 \in B(\gamma_2)$, $Z_2^\circ[\nu_2]$ is non-empty and contains an irreducible component Γ such that $\text{codim}(\Gamma, Z_2) = \text{codim}(\text{Sh}_2[\nu_2], \text{Sh}_2)$. Also assume that $u_1 \ll_{(C)} u_2$ and $u_2 \ll_{(C)} u_2$. Then there exists a smooth curve with Newton polygon $u_1^{n_1} \oplus u_2^{n_2-1} \oplus \nu_2 \oplus \text{ord}^{(n_1+n_2-2)(m-1)+(r-1)}$.*

Proof. If $n_2 = 1$, we first apply Corollary 4.9 with $t = m$ to produce a family Z_3 with Newton polygon $u_1^{n_1} \oplus \text{ord}^{(n_1-1)(m-1)}$. Note that Z_3 and Z_2 satisfy hypotheses (A) and (B). Since $u_1 \ll_{(C)} u_2$, by Lemma 6.7 (1), Z_3 and Z_2 also satisfy hypothesis (C). Applying Theorem 6.11 produces a smooth curve with Newton polygon $u_1^{n_1} \oplus \nu_2 \oplus \text{ord}^{(n_1-1)(m-1)+(r-1)}$.

For $n_2 \geq 2$, we apply Corollary 4.10 to produce a family Z_4 with Newton polygon $u_1^{n_1} \oplus u_2^{n_2-1} \oplus \text{ord}^{(n_1+n_2-3)(m-1)}$. Since $u_1 \ll_{(C)} u_2$ and $u_2 \ll_{(C)} u_2$, by Lemma 6.7 (2), Z_4 and Z_2 satisfy hypotheses (A), (B), and (C). Applying Remark 3.7, we produce generalized monodromy data for Z_4 and Z_2 by marking an additional unramified fiber. In this situation, the toric rank is $\epsilon' = m - 1$. Applying Theorem 6.11 to the generalized families for Z_4 and Z_2 completes the proof. \square

Corollary 6.16 applies to the pair of monodromy data in the proof of Corollary 9.7.

6.5. An exceptional example. We give an example of a pair of monodromy data, and non μ -ordinary Newton polygon ν_2 , satisfying hypotheses (A) and (B), but not (C), for which (6.1) can be verified directly. Furthermore, as the Kottwitz set B_2 has size 2, this example also shows that hypothesis (C) is sufficient but not necessary for Proposition 6.8 to hold.

Recall that ss is the Newton polygon $(1/2, 1/2)$.

Proposition 6.17. *If $p \equiv 7 \pmod{8}$ is sufficiently large, then there exists a smooth curve over $\overline{\mathbb{F}}_p$ of genus 9 with Newton polygon $ss^7 \oplus \text{ord}^2$.*

Proof. Let $Z_2 = Z(8, 4, (4, 2, 5, 5))$. Then Z_2 is the special family $M[15]$ in [25, Table 1], and the associated Shimura variety Sh_2 has signature type $\mathfrak{f}_2 = (1, 1, 0, 0, 2, 0, 1)$. At any prime $p \equiv 7 \pmod{8}$, the μ -ordinary Newton polygon is $u_2 = \text{ord}^2 \oplus ss^3$ and the basic Newton polygon is $v_2 = ss^5$ [21, Section 6.2].

Let $Z_1 = Z(4, 3, (1, 1, 2))$, which has signature $(1, 0, 0)$. At any prime $p \equiv 7 \pmod{8}$, the μ -ordinary Newton polygon is $u_1 = ss$ [22, Section 4, $m = 4$]. Then $d = 2$ and $r = 2$. By Section 3.1, the induced signature type is $\mathfrak{f}_1^\dagger = (1, 0, 0, 0, 1, 0, 0)$.

The pair of monodromy data for Z_1 and Z_2 satisfies hypothesis (A). Let $p \equiv 7 \pmod{8}$; then it also satisfies hypothesis (B). For the orbit $\mathfrak{o} = \{1, 7\}$, by [21, Example 4.5], $u_1(\mathfrak{o})$ has slopes $1/2$ and $u_2(\mathfrak{o})$ has slopes 0 and 1. Thus the pair does not satisfy hypothesis (C).

The image of $\tilde{Z}_1 \times \tilde{Z}_2$ under the clutching morphism lies in the family \tilde{Z}_3 of curves with monodromy datum $(8, 5, (2, 2, 2, 5, 5))$. The Shimura variety Sh_3 has signature type $\mathfrak{f}_3 = (2, 2, 0, 0, 3, 1, 1)$ and its μ -ordinary Newton polygon is $u_3 = u_1^2 \oplus u_2 \oplus \text{ord}^2 = ss^5 \oplus \text{ord}^4$ by Proposition 4.4. By [21, Section 4.3], there is only one element u_3 in $B(\text{Sh}_3)$ which is strictly larger than $ss^7 \oplus \text{ord}^2 = u_1^2 \oplus v \oplus \text{ord}^2$. From (2.6), we see that the codimension of $\text{Sh}_3[u_1 \oplus v_2 \oplus \text{ord}^2]$ in Sh_3 is 1. Thus, we conclude by Remark 6.12 and Theorem 6.11. \square

7. SUPERSINGULAR CASES IN MOONEN'S TABLE

In [25, Theorem 3.6], Moonen proved there are exactly 20 positive-dimensional special families arising from cyclic covers of \mathbb{P}^1 . In [21, Section 6], we computed all of the Newton polygons v that occur on the corresponding Shimura varieties using the Kottwitz method, see Section 10. Moreover, in [21, Theorem 1.1], we proved that the open Torelli locus intersects each non-supersingular (resp. supersingular) Newton polygon stratum (resp. as long as the family has dimension 1 and p is sufficiently large).

In this section, we extend [21, Theorem 1.1] to include the supersingular Newton polygon strata in the five remaining cases when the dimension of the family is greater than 1, using results from Section 6. Case (5) is note-worthy since it was not previously known that there exists a smooth supersingular curve of genus 6 when $p \equiv 2, 3, 4 \pmod{5}$, see [22, Theorem 1.1] and [21, Theorem 1.1] for related results.

Theorem 7.1. *There exists a smooth supersingular curve of genus g defined over $\overline{\mathbb{F}}_p$ for all sufficiently large primes satisfying the given congruence condition in the following families:*

- (1) $g = 3$, when $p \equiv 2 \pmod{3}$, in the family $M[6]$;
- (2) $g = 3$, when $p \equiv 3 \pmod{4}$, in the family $M[8]$;
- (3) $g = 4$, when $p \equiv 2 \pmod{3}$, in the family $M[10]$;
- (4) $g = 4$, when $p \equiv 5 \pmod{6}$, in the family $M[14]$; and
- (5) $g = 6$, when $p \equiv 2, 3, 4 \pmod{5}$, in the family $M[16]$.

Corollary 7.2. *Let $\gamma = (m, N, a)$ denote the monodromy datum for one of Moonen's special families from [25, Table 1]. Assume $p \nmid m$. Let $v \in \nu(B(\mu_m, \mathfrak{f}))$ be a Newton polygon occurring on $\text{Sh}(\gamma)$ as in Section 2.6. Then v occurs as the Newton polygon of a smooth curve in the family $Z^\circ(\gamma)$, as long as p is sufficiently large when v is supersingular.*

Proof. The proof is immediate from [21, Theorem 1.1] and Theorem 7.1. \square

Proof of Theorem 7.1 in cases (1), (2), (4), and (5). Let γ denote the monodromy datum, let Z denote the special family of curves and let Sh denote the corresponding Shimura variety and suppose that $p \not\equiv 1 \pmod{m}$. Then $\dim(Z) = \dim(\text{Sh}) = 2$, and the basic locus $\text{Sh}[\nu]$ is supersingular with codimension 1 in Sh .

Following [21, Section 5.2], a point of $\text{Sh}[\nu]$ which is not in the image of Z° is the Jacobian of a singular curve of compact type. This point arises from an admissible clutching of points from two families Z_1 and Z_2 . This yields an admissible degeneration of the inertia type, see [21, Definition 5.4]. A complete list of admissible degenerations of the inertia type for Moonen's families can be found in [21, Lemma 6.4]. In each of these cases, there exists an admissible degeneration such that $\dim(Z_1) = 0$ and the μ -ordinary Newton polygon u_1 for Z_1 is supersingular, and $m_1 = m_2$ (so $d = 1$).

In the degenerations from [21, Lemma 6.4], one checks using [21, Sections 6.1-6.2] that Z_2 is a special family with $\dim(Z_2) = 1$ and that Z_2 has exactly two Newton polygons, the μ -ordinary one u_2 and the basic one ν_2 which is supersingular. By [21, Theorem 1.1], for p sufficiently large, $Z_2^\circ[\nu_2]$ is non-empty. Since there are exactly two Newton polygons on Z , we conclude that these are $u = u_1 \oplus u_2$ and $\nu = u_1 \oplus \nu_2$. By Proposition 4.4, the pair of monodromy data for Z_1 and Z_2 satisfies hypothesis (B). The codimension condition in (6.1) is satisfied since the basic locus has codimension 1 in both Z and Z_2 . By Remark 6.12 and Theorem 6.11, there exists a 1-dimensional family of smooth curves in Z with the basic Newton polygon ν , which is supersingular. \square

Proof of Theorem 7.1 in case (3). We use the same notation as in the first 2 paragraphs of the proof of the other cases. The only difference in case (3) is that $\dim(Z) = \dim(\text{Sh}) = 3$ and the basic locus is supersingular with codimension 2 in Sh . In case (3), the only admissible degeneration comes from the pair of monodromy data $\gamma_1 = (3, 3, (1, 1, 1))$ and $\gamma_2 = (3, 5, (2, 1, 1, 1, 1))$. The latter of these is the monodromy datum for the special family $M[6]$. The basic locus $\text{Sh}[\nu]$ has dimension 1. The codimension condition in (6.1) is not satisfied in this situation: $\text{codim}(\text{Sh}_2[\nu_2], \text{Sh}_2) = 1$, while $\text{codim}(\text{Sh}[\nu], \text{Sh}) = 2$.

For p sufficiently large, we claim that the number of irreducible components of $\text{Sh}[\nu]$ exceeds the number that arise from the boundary of Z . Let W be a 1-dimensional family of supersingular singular curves in $Z \setminus Z^\circ$. The only way to construct such a family W is to clutch a genus 1 curve with μ_3 -action together with a 1-dimensional family of supersingular curves in $M[6]$. In other words, W arises as the image under κ of $T_1 \times T_2$, for some component T_1 of $\text{Sh}(3, 3, (1, 1, 1))$ and some component T_2 of the supersingular locus of $M[6]$. The number of choices for T_1 , for the μ_3 -actions, and for the labelings of the ramification points is a fixed constant that does not depend on p .

Thus it suffices to compare the number $s_{M[10]}$ of irreducible components of the supersingular locus in $M[10]$ with the number $s_{M[6]}$ of irreducible components T_2 of the supersingular locus in $M[6]$ when $p \equiv 2 \pmod{3}$. The signature type for $M[10]$ is $(1, 3)$. By [21, Theorem 8.1], the number $s_{M[10]}$ grows with respect to p .

The signature type for $M[6]$ is $(1, 2)$. By [21, Remark 8.2], we see that $s_{M[6]}$ is the same for all odd $p \equiv 2 \pmod{3}$. More precisely, note that $\dim(\text{Sh}_2) = 2 \dim(\text{Sh}_2(\nu_2))$ when $p \equiv 2 \pmod{3}$, that the center of the associated reductive group is connected, and that the supersingular locus is the basic locus. Thus by [34, Remark 1.1.5 (2)], all odd $p \equiv 2 \pmod{3}$ satisfy the hypothesis of [34, Theorem 1.1.4 (1), Proposition 7.4.2], which provides an expression for $s_{M[6]}$ over $\overline{\mathbb{F}}_p$ in terms of objects independent of p .

Hence there exist irreducible components of $\text{Sh}[v]$ which contain the Jacobian of a smooth curve, for p sufficiently large. \square

8. UNLIKELY INTERSECTIONS

In this section, we prove that the non-trivial intersection of the open Torelli locus with the Newton polygon strata found in most of the results of the paper is unexpected.

Recall that ss denotes the Newton polygon $(1/2, 1/2)$.

Definition 8.1. Let ν be a symmetric Newton polygon of height $2g$, and let $\mathcal{A}_g[\nu]$ be its Newton polygon stratum in the Siegel variety \mathcal{A}_g . Then ν satisfies condition (U) if $\dim(\mathcal{M}_g) < \text{codim}(\mathcal{A}_g[\nu], \mathcal{A}_g)$.

Definition 8.2. The open Torelli locus has an *unlikely intersection* with $\mathcal{A}_g[\nu]$ in \mathcal{A}_g if there exists a smooth curve of genus g with Newton polygon ν , and ν satisfies condition (U).

8.1. The codimension of Newton polygon strata in Siegel varieties. We study the codimension of the Newton strata in \mathcal{A}_g . By [26, Theorem 4.1], see also (2.6),

$$(8.1) \quad \text{codim}(\mathcal{A}_g[\nu], \mathcal{A}_g) = \#\Omega(\nu),$$

where $\Omega(\nu) := \{(x, y) \in \mathbb{Z} \times \mathbb{Z} \mid 0 \leq x, y \leq g, (x, y) \text{ strictly below } \nu\}$.

Remark 8.3. By (8.1), if ν is non-ordinary, then $\text{codim}(\mathcal{A}_{ng}[\nu^n], \mathcal{A}_{ng})$ grows quadratically in n . In particular, if $\nu = ss$, then $\text{codim}(\mathcal{A}_n[ss^n], \mathcal{A}_n) = n(n+1)/2 - \lfloor n^2/4 \rfloor > n^2/4$.

Proposition 8.4. Let $\{u_n\}_{n \in \mathbb{N}}$ be a sequence of symmetric Newton polygons. Let $2g_n$ be the height of u_n . Suppose there exists $\lambda \in \mathbb{Q} \cap (0, 1)$ such that the multiplicity of λ as a slope of u_n is at least n for each $n \in \mathbb{N}$. Then $\text{codim}(\mathcal{A}_{g_n}[u_n], \mathcal{A}_{g_n})$ grows at least quadratically in n .

Proof. Let $\nu = (\lambda, 1 - \lambda)$ and let h be the height of ν . By hypothesis, $u_n = \nu^n \oplus v_n$ for some symmetric Newton polygon v_n for each $n \in \mathbb{N}$ and $g_n \geq nh$. Since v_n lies on or above $\text{ord}^{g_n - nh}$, then $u_n = \nu^n \oplus v_n$ lies on or above $\nu^n \oplus \text{ord}^{g_n - nh}$. Hence

$$\text{codim}(\mathcal{A}_{g_n}[u_n], \mathcal{A}_{g_n}) \geq \text{codim}(\mathcal{A}_{g_n}[\nu^n \oplus \text{ord}^{g_n - nh}], \mathcal{A}_{g_n}).$$

By (8.1), or alternatively Proposition 6.8,

$$\text{codim}(\mathcal{A}_{g_n}[\nu^n \oplus \text{ord}^{g_n - nh}], \mathcal{A}_{g_n}) \geq \text{codim}(\mathcal{A}_{nh}[\nu^n], \mathcal{A}_{nh}).$$

Thus $\text{codim}(\mathcal{A}_{g_n}[u_n], \mathcal{A}_{g_n}) \geq \text{codim}(\mathcal{A}_{nh}[\nu^n], \mathcal{A}_{nh})$, which is sufficient by Remark 8.3. \square

8.2. Verifying condition (U). Given a sequence $\{u_n\}_{n \in \mathbb{N}}$ of symmetric Newton polygons of increasing height, we state simple criteria to ensure that all but finitely many of them satisfy condition (U). Let $2g_n$ be the height of u_n .

Proposition 8.5. Assume that g_n grows linearly in n and that there exists $\lambda \in \mathbb{Q} \cap (0, 1)$ such that the multiplicity of λ as a slope of u_n grows linearly in n , for all sufficiently large $n \in \mathbb{N}$. Then, for all sufficiently large n , the Newton polygon u_n satisfies condition (U).

Proof. By Proposition 8.4, $\text{codim}(\mathcal{A}_{g_n}[u_n], \mathcal{A}_{g_n})$ is quadratic in n while $\dim(\mathcal{M}_{g_n}) = 3g_n - 3$ is linear in n by hypothesis. Thus $\dim(\mathcal{M}_{g_n}) < \text{codim}(\mathcal{A}_{g_n}[u_n], \mathcal{A}_{g_n})$ for $n \gg 0$. \square

Proposition 8.6. If there exists $t \in \mathbb{R}_{>0}$ such that the multiplicity of $1/2$ as a slope of u_n is at least $2tg_n$, for all $n \in \mathbb{N}$, then u_n satisfies condition (U) for each $n \in \mathbb{N}$ such that $g_n \geq 12/t^2$.

Proof. By the proof of Proposition 8.4 and Remark 8.3,

$$\text{codim}(\mathcal{A}_{g_n}[u_n], \mathcal{A}_{g_n}) \geq \text{codim}(\mathcal{A}_{\lceil tg_n \rceil}[\text{ss}^{\lceil tg_n \rceil}], \mathcal{A}_{\lceil tg_n \rceil}) > (tg_n)^2/4.$$

So condition (U) for u_n is true when $(tg_n)^2/4 \geq (3g_n - 3)$ and thus when $g_n \geq 12/t^2$. \square

Proposition 8.7. *Let v_1, v_2 be two symmetric Newton polygons, respectively of height $2g \geq 2$, and $2h \geq 0$. Assume v_1 is not ordinary. Then*

- (1) *for all sufficiently large $n \in \mathbb{N}$, the Newton polygon $v_1^n \oplus v_2$ satisfies condition (U);*
- (2) *if $1/2$ occurs as a slope of v_1 with multiplicity $2\delta > 0$, then the Newton polygon $v_1^n \oplus v_2$ satisfies condition (U), for each $n \geq \max\{15g/\delta^2, 9\sqrt{h}/\delta\}$.¹⁴*

Proof. (1) Let $\lambda \in \mathbb{Q} \cap (0, 1)$ be a slope of v_1 , occurring with multiplicity $m_\lambda \geq 1$. Then, for each $n \in \mathbb{N}$, the Newton polygon $u_n = v_1^n \oplus v_2$ has height $2g_n = 2/ng + h$ and slope λ occurring with multiplicity at least $m_\lambda n$. Taking $u_n = v_1^n \oplus v_2$, the sequence $\{u_n\}_{n \in \mathbb{N}}$ satisfies the hypotheses of Proposition 8.5. Hence, part (1) holds.

(2) As for Proposition 8.4, $\text{codim}(\mathcal{A}_{ng}[v_1^n], \mathcal{A}_{ng}) \leq \text{codim}(\mathcal{A}_{ng+h}[v_1^n \oplus v_2], \mathcal{A}_{ng+h})$. Therefore, condition (U) for $v_1^n \oplus v_2$ is implied by the inequality

$$(8.2) \quad \dim(\mathcal{M}_{ng+h}) < \text{codim}(\mathcal{A}_{ng}[v_1^n], \mathcal{A}_{ng}).$$

Following the proof of Proposition 8.6, if the slope $1/2$ occurs in v_1 with multiplicity 2δ , then inequality (8.2) is true if $3/ng + h - 1 \leq (n\delta)^2/4$, which holds for $n \geq N := 6g\delta^{-2}(1 + (1 + \delta^2(h - 1)3^{-1}g^{-2})^{1/2})$. The asserted bound follows by noticing that $N < \max\{6(1 + \sqrt{2})g/\delta^2, 2\sqrt{3}(1 + \sqrt{2})\sqrt{h}/\delta\}$. \square

Remark 8.8. For $g \gg 0$, Proposition 8.7 implies that the non-trivial intersections of \mathcal{T}_g° with $\mathcal{A}_g[v]$ in Corollaries 4.8, 4.9, and 6.15 (resp. 4.10 and 6.16) are unlikely if the μ -ordinary Newton polygon u is not ordinary. (resp. if either u_1 or u_2 is not ordinary).

Remark 8.9. Consider the following refinement of Definition 8.2: a non-empty substack U of $\mathcal{T}_g \cap \mathcal{A}_g[v]$ is an *unlikely intersection* if $\text{codim}(U, \mathcal{M}_g) < \text{codim}(\mathcal{A}_g[v], \mathcal{A}_g)$.

The results in Sections 4.3 and 6.4 yield families Z of cyclic covers of \mathbb{P}^1 such that $Z^\circ[v]$ is non-empty and has the expected codimension in Z . This produces an unlikely intersection as in Remark 8.9 for $g \gg 0$, when the initial Newton polygon u is not ordinary.

9. APPLICATIONS

We apply the results in Sections 4.3 and 6.4 to construct smooth curves of arbitrarily large genus g with prescribed Newton polygon v . By Proposition 8.7, when g is sufficiently large, the curves in this section lie in the unlikely intersection $\mathcal{T}_g^\circ \cap \mathcal{A}_g[v]$.

Notation 9.1. For $s, t \in \mathbb{N}$, with $s \leq t/2$ and $\text{gcd}(s, t) = 1$, we write $(s/t, (t - s)/t)$ for the Newton polygon of height $2t$ with slopes s/t and $(t - s)/t$, each with multiplicity t .

¹⁴This bound is not sharp, but it is written so that the asymptotic dependency on g, δ, h is more clear.

9.1. Newton polygons with many slopes of $1/2$. We obtain examples of smooth curves of arbitrarily large genus g such that the only slopes of the Newton polygons are $0, \frac{1}{2}, 1$. We focus on examples where the multiplicity of $1/2$ is large relative to g .

Corollary 9.2. *Let $m \in \mathbb{Z}_{\geq 1}$ be odd and $h = (m - 1)/2$. Let p be a prime, $p \nmid 2m$, such that the order f of p in $(\mathbb{Z}/m\mathbb{Z})^*$ is even and $p^{f/2} \equiv -1 \pmod{m}$. For $n \in \mathbb{Z}_{\geq 1}$, there exists a μ_m -cover $C \rightarrow \mathbb{P}^1$ defined over $\overline{\mathbb{F}}_p$ where C is a smooth curve of genus $g = h(3n - 2)$ with Newton polygon $\nu = ss^{hn} \oplus \text{ord}^{2h(n-1)}$. If $n \geq 34/h$, then $\text{Jac}(C)$ lies in the unlikely intersection $\mathcal{T}_g^\circ \cap \mathcal{A}_g[\nu]$.*

Proof. Let $C \rightarrow \mathbb{P}^1$ be a μ_m -cover with $\gamma = (m, 3, a)$ where $a = (1, 1, m - 2)$. Without loss of generality, an equation for C is $y^m = x^2 - 1$. By [35, Theorem 6.1], the Newton polygon of C is ss^h . The first claim follows from applying Corollary 4.9 to $Z(m, 3, \gamma)$ with $t = m$. As in the proof of Proposition 8.6, the second claim follows from the inequalities:

$$\text{codim}(\mathcal{A}_{nh}[ss^{nh}], \mathcal{A}_{nh}) \geq (nh)^2/4 + (nh)/2 > \dim(\mathcal{M}_{3nh-2h}) = 9nh - 6h - 3. \quad \square$$

Remark 9.3. The Newton polygons in Corollary 9.2 are μ -ordinary; they do not appear in the literature, but the result also follows from Proposition 4.6(3) if $p \equiv -1 \pmod{m}$ or if $p \geq m(N - 3)$ where N is the (increasingly large) number of branch points.

We highlight the case $m = 3$ below. To our knowledge, for any odd prime p , this is the first time that a sequence of smooth curves has been produced for every $g \in \mathbb{Z}_{\geq 1}$ such that the multiplicity of the slope $1/2$ in the Newton polygon grows linearly in g .

Corollary 9.4. *Let $p \equiv 2 \pmod{3}$ be an odd prime. Let $g \in \mathbb{Z}_{\geq 1}$. There exists a smooth curve C_g of genus g defined over $\overline{\mathbb{F}}_p$, whose Newton polygon ν_g only has slopes $0, \frac{1}{2}, 1$ and such that the multiplicity of the slope $1/2$ is at least $2\lfloor g/3 \rfloor$. If $g \geq 107$, the curve C_g demonstrates an unlikely intersection of the open Torelli locus with the Newton polygon stratum $\mathcal{A}_g[\nu_g]$ in \mathcal{A}_g .*

Proof. If $g = 3n - 2$ for some n , the result is immediate from Corollary 9.2. For $g = 3n - 2 + 2\epsilon$ with $\epsilon = 1$ (resp. $\epsilon = 2$), we apply Corollary 4.7 with $t = 1$ (resp. twice) and obtain a smooth curve with Newton polygon $ss^n \oplus \text{ord}^{2n-2+2\epsilon}$. \square

Working with Moonen's families gives examples of families of curves where the multiplicity of the slope $1/2$ is particularly high relative to the genus.

Corollary 9.5. *Let $p \equiv 4 \pmod{5}$. For $n \in \mathbb{Z}_{\geq 1}$, there exists a smooth curve of genus $g = 10n - 4$ in $Z = Z(5, 5n, (2, 2, \dots, 2))$ over $\overline{\mathbb{F}}_p$ with μ -ordinary Newton polygon $u_n = ss^{4n} \oplus \text{ord}^{6n-4}$.*

For $n \geq 7$, the curves with Newton polygon u_n from Corollary 9.5 lie in the unlikely intersection $\mathcal{T}_g^\circ \cap \mathcal{A}_g[\nu]$.

Proof. When $p \equiv 4 \pmod{5}$, $M[16]$ has μ -ordinary Newton polygon $u_1 = \text{ord}^2 \oplus ss^4$.¹⁵ The claim is immediate from Corollary 4.9. \square

Corollary 9.6. *Under the given congruence condition on p , and with $p \gg 0$, there exists a smooth curve in $Z = Z(m, N, a)$ over $\overline{\mathbb{F}}_p$ with Newton polygon ν and $\text{codim}(Z[\nu], Z) = 1$.*

¹⁵The codimension condition in (6.1) does not hold for $\nu = ss^6$.

construction	(m, N, a)	genus	congruence	Newton Polygon ν
$M[9] + M[9]$	$(6, 6, (1, 1, 4, 4, 4, 4))$	8	$2 \bmod 3$	$ss^4 \oplus ord^4$
$M[9] + M[12]$	$(6, 6, (1, 1, 1, 1, 4, 4))$	9	$2 \bmod 3$	$ss^5 \oplus ord^4$
$M[12] + M[12]$	$(6, 6, (1, 1, 1, 1, 1, 1))$	10	$2 \bmod 3$	$ss^7 \oplus ord^3$
$M[18] + M[18]$	$(10, 6, (3, 3, 6, 6, 6, 6))$	16	$9 \bmod 10$	$ss^{10} \oplus ord^6$
$M[20] + M[20]$	$(12, 6, (4, 4, 7, 7, 7, 7))$	19	$11 \bmod 12$	$ss^{12} \oplus ord^7$

9.2. Newton polygons with slopes $1/3$, $1/4$, and beyond.

Corollary 9.7. *Let $n \in \mathbb{Z}_{\geq 1}$. The following Newton polygons occur for Jacobians of smooth curves over $\overline{\mathbb{F}}_p$ under the given congruence condition on p .*

congruence	ν (μ -ordinary)	ν (non μ -ordinary) for $p \gg 0$
$2, 4 \bmod 7$	$(1/3, 2/3)^n \oplus ord^{6n-6}$	NA
$3, 5 \bmod 7$	$(1/3, 2/3)^{2n} \oplus ord^{6n-6}$	$(1/3, 2/3)^{2n-2} \oplus ss^6 \oplus ord^{6n-6}$
$2, 5 \bmod 9$	$(1/3, 2/3)^{2n} \oplus ss^n \oplus ord^{8n-8}$	$(1/3, 2/3)^{2n-2} \oplus ss^{n+6} \oplus ord^{8n-8}$
$4, 7 \bmod 9$	$(1/3, 2/3)^{2n} \oplus ord^{9n-8}$	$(1/3, 2/3)^{2n-2} \oplus ss^6 \oplus ord^{9n-8}$

We remark that none of the last three lines follows from [5, Theorem 6.1] because there are at least two Newton polygons in $B(\mu_m, f)$ having the maximal p -rank.

Proof. Lines 1, 2, and 3 are obtained from applying both Corollaries 4.9 and 6.15 to the families $(7, 3, (1, 1, 5))$, $M[17]$, and $M[19]$, respectively.

For the last line, let $m = 9$ and $p \equiv 4, 7 \bmod 9$. There are four orbits $\sigma_1 = (1, 4, 7)$, $\sigma_2 = (2, 5, 8)$, $\sigma_3 = (3)$, and $\sigma_4 = (6)$. The μ -ordinary Newton polygon for the family $M[19]$ is $u = (1/3, 2/3)^2 \oplus ord$, and $\nu = ss^6 \oplus ord$ also occurs for a smooth curve in the family. By [21, Section 6.2], for each $\sigma \in \mathfrak{D}$, $u(\sigma)$ has at most 2 slopes, hence hypothesis (C) is satisfied, and we obtain the Newton polygons in line 4 from Corollaries 4.9 and 6.15.¹⁶ \square

Corollary 9.8. *Let $n \in \mathbb{Z}_{\geq 1}$. The following Newton polygons occur for Jacobians of smooth curves over $\overline{\mathbb{F}}_p$ under the given congruence condition on p .*

congruence	ν (μ -ordinary)	ν (non μ -ordinary) for $p \gg 0$
$2, 3 \bmod 5$	$(1/4, 3/4)^n \oplus ord^{4n-4}$	$(1/4, 3/4)^{n-1} \oplus ss^4 \oplus ord^{4n-4}$
$3, 7 \bmod 10$	$(1/4, 3/4)^n \oplus ss^{2n} \oplus ord^{9n-9}$	$(1/4, 3/4)^{n-1} \oplus ss^{2n+4} \oplus ord^{9n-9}$

Proof. The proof follows from Corollaries 4.9 and 6.15 applied to $M[11]$ and $M[18]$. \square

Corollary 9.9. *Let $p \equiv 2, 3 \bmod 5$. For any $n_1, n_2 \in \mathbb{Z}_{\geq 1}$, there exists a smooth curve of genus $g = 6n_1 + 8n_2$ defined over $\overline{\mathbb{F}}_p$ with Newton polygon $(1/4, 3/4)^{n_2+1} \oplus ss^{2n_1} \oplus ord^{4(n_1+n_2-1)}$.*

Proof. We apply Corollary 4.10 to $Z_1 = Z(5, 3, (2, 2, 1))$ and $Z_2 = M[11]$. By [21, Section 6.2] and [22, Section 4], if $p \equiv 2, 3 \bmod 5$, then $u_1 = ss^2$ and $u_2 = (1/4, 3/4)$.¹⁷ \square

Example 9.10. Let m be prime and p have odd order modulo m . The Newton polygon ν_1 for a μ_m -cover with monodromy datum $\gamma = (m, 3, a)$ has no slopes of $1/2$ by [22,

¹⁶Alternatively, applying Corollaries 4.10 and 6.16 produces the Newton polygons $(1/3, 2/3)^{n_1+2n_2} \oplus ord^{8n_1+9n_2-14}$ and $(1/3, 2/3)^{n_1+2n_2-2} \oplus ss^6 \oplus ord^{8n_1+9n_2-13}$ for $n_1, n_2 \in \mathbb{Z}_{\geq 1}$.

¹⁷The pair Z_1 and Z_2 does not satisfy hypothesis (C) and the codimension condition in (6.1) does not hold inductively.

Section 3.2]. Applying Corollary 4.9 to $Z = Z(\gamma)$ with $t = m$ shows that the Newton polygon $\nu_n = \nu_1^n \oplus \text{ord}^{(m-1)(n-1)}$ occurs for a smooth curve over $\overline{\mathbb{F}}_p$, for any $n \in \mathbb{Z}_{\geq 1}$.

Examples of γ and ν_1 can be found in [22, Theorem 5.4]. For example, when $m = 11$, $a = (1, 1, 9)$ and $p \equiv 3, 4, 5, 9 \pmod{11}$, then $\nu_1 = (1/5, 4/5)$. As another example, let $m = 29$, $a = (1, 1, 27)$, and $p \equiv 7, 16, 20, 23, 24, 25 \pmod{29}$, then $\nu_1 = (2/7, 5/7) \oplus (3/7, 4/7)$, yielding another infinite family that cannot be studied using [5, Theorem 6.1].

10. APPENDIX: NEWTON POLYGONS FOR MOONEN'S FAMILIES

For convenience, we provide the full list of Newton polygons on Moonen's special families from [21, Section 6]. These occur for a smooth curve in the family by Corollary 7.2. The label $M[r]$ is from [25, Table 1]. The notation † means **we further assume** $p \gg 0$.

Label	m	a, f	Newton Polygon [congruence on p]
M[1]	2	(1,1,1,1), f = (1)	ord, ss [1 mod 2]
M[2]	2	(1,1,1,1,1,1), f = (2)	ord ² , ord ⊕ ss, ss ² [1 mod 2]
M[3]	3	(1,1,2,2), f = (1, 1)	ord ² [1, 2 mod 3], ss ² [1 mod 3], ss ² [2 mod 3] [†]
M[4]	4	(1,2,2,3) f = (1, 0, 1)	ord ² [1, 3 mod 4] ss ² [1 mod 4], ss ² [3 mod 4] [†]
M[5]	6	(2,3,3,4) f = (1, 0, 0, 0, 1)	ord ² [1, 5 mod 6] ss ² [1 mod 6], ss ² [5 mod 6] [†]
M[6]	3	(1,1,1,1,2) f = (2, 1)	ord ³ [1 mod 3], ord ² ⊕ ss [2 mod 3] ord ⊕ ss ² , (1/3, 2/3) [1 mod 3], ss ³ [2 mod 3] [†]
M[7]	4	(1,1,1,1) f = (2, 1, 0)	ord ³ [1 mod 4], ord ⊕ ss ² [3 mod 4] ord ² ⊕ ss [1 mod 4], ss ³ [3 mod 4] [†]
M[8]	4	(1,1,2,2,2) f = (2, 0, 1)	ord ³ [1 mod 4], ord ² ⊕ ss [3 mod 4] ord ⊕ ss ² , (1/3, 2/3) [1 mod 4], ss ³ [3 mod 4] [†]
M[9]	6	(1,3,4,4) f = (1, 1, 0, 0, 1)	ord ³ [1 mod 6], ord ² ⊕ ss [5 mod 6] ord ⊕ ss ² [1 mod 6], ss ³ [5 mod 6] [†]
M[10]	3	(1,1,1,1,1,1) f = (3, 1)	ord ⁴ [1 mod 3], ord ² ⊕ ss ² [2 mod 3] ord ² ⊕ ss ² [1 mod 3], (1/4, 3/4) [2 mod 3] ord ⊕ (1/3, 2/3) [1 mod 3], ss ⁴ [2 mod 3] [†] (1/4, 3/4) [1 mod 3]
M[11]	5	(1,3,3,3) f = (1, 2, 0, 1)	ord ⁴ [1 mod 5], (1/4, 3/4) [2, 3 mod 5], ord ² ⊕ ss ² [4 mod 5] ord ² ⊕ ss ² [1 mod 5], ss ⁴ [2, 3, 4 mod 5] [†]
M[12]	6	(1,1,1,3) f = (2, 1, 1, 0, 0)	ord ⁴ [1 mod 6], ord ⊕ ss ³ [5 mod 6] ord ³ ⊕ ss [1 mod 6], ss ⁴ [5 mod 6] [†]
M[13]	6	(1,1,2,2) f = (2, 1, 0, 1, 0)	ord ⁴ [1 mod 6], ord ² ⊕ ss ² [5 mod 6] ord ² ⊕ ss ² [1 mod 6], ss ⁴ [5 mod 6] [†]
M[14]	6	(2,2,2,3,3) f = (2, 0, 0, 1, 1)	ord ⁴ [1 mod 6], ord ² ⊕ ss ² [5 mod 6] ord ² ⊕ ss ² [1 mod 6], ss ⁴ [5 mod 6] [†] ord ⊕ (1/3, 2/3) [1 mod 6]
M[15]	8	(2,4,5,5), f = (1, 1, 0, 0, 2, 0, 1)	ord ⁵ [1 mod 8], ord ² ⊕ ss ³ [3, 7 mod 8], ord ³ ⊕ ss ² [5 mod 8] ord ³ ⊕ ss ² [1 mod 8], (1/4, 3/4) ⊕ ss [3 mod 8] ord ⊕ (1/4, 3/4) [5 mod 8], ss ⁵ [7 mod 8] [†]
M[16]	5	(2,2,2,2,2) f = (2, 0, 3, 1)	ord ⁶ [1 mod 5], (1/4, 3/4) ⊕ ss ² [2, 3 mod 5], ord ² ⊕ ss ⁴ [4 mod 5] ord ⁴ ⊕ ss ² [1 mod 5], ss ⁶ [2, 3, 4 mod 5] [†] ord ³ ⊕ (1/3, 2/3) [1 mod 5]
M[17]	7	(2,4,4,4), f = (1, 2, 0, 2, 0, 1)	ord ⁶ [1 mod 7], ord ³ ⊕ (1/3, 2/3) [2, 4 mod 7] (1/3, 2/3) ² [3, 5 mod 7], ord ² ⊕ ss ⁴ [6 mod 7] ord ⁴ ⊕ ss ² [1 mod 7], (1/6, 5/6) [2, 4 mod 7], ss ⁶ [3, 5, 6 mod 7] [†]
M[18]	10	(3,5,6,6), f = (1, 1, 0, 1, 0, 0, 2, 0, 1)	ord ⁶ [1 mod 10], (1/4, 3/4) ⊕ ss ² [3, 7 mod 10], ord ² ⊕ ss ⁴ [9 mod 10] ord ⁴ ⊕ ss ² [1 mod 10], ss ⁶ [3, 7, 9 mod 10] [†]
M[19]	9	(3,5,5,5), f = (1, 2, 0, 2, 0, 1, 0, 1)	ord ⁷ [1 mod 9], (1/3, 2/3) ² ⊕ ss [2, 5 mod 9] ord ⊕ (1/3, 2/3) ² [4, 7 mod 9], ord ² ⊕ ss ⁵ [8 mod 9] ord ⁵ ⊕ ss ² [1 mod 9], ss ⁷ [2, 5, 8 mod 9] [†] , ord ⊕ ss ⁶ [4, 7 mod 9]
M[20]	12	(4,6,7,7), f = (1, 1, 0, 1, 0, 0, 2, 0, 1, 0, 1)	ord ⁷ [1 mod 12], ord ³ ⊕ ss ⁴ [5 mod 12] ord ⁴ ⊕ ss ³ [7 mod 12], ord ² ⊕ ss ⁵ [11 mod 12] ord ⁵ ⊕ ss ² [1 mod 12], ord ⊕ (1/4, 3/4) ⊕ ss ² [5 mod 12] ord ² ⊕ ss ⁵ [7 mod 12], ss ⁷ [11 mod 12] [†]

REFERENCES

1. Jeffrey D. Achter and Rachel Pries, *The integral monodromy of hyperelliptic and trielliptic curves*, *Math. Ann.* **338** (2007), no. 1, 187–206. MR 2295509
2. ———, *Monodromy of the p -rank strata of the moduli space of curves*, *Int. Math. Res. Not. IMRN* (2008), no. 15, Art. ID rnn053, 25. MR 2438069
3. ———, *The p -rank strata of the moduli space of hyperelliptic curves*, *Adv. Math.* **227** (2011), no. 5, 1846–1872. MR 2803789
4. Siegfried Bosch, Werner Lütkebohmert, and Michel Raynaud, *Néron models*, *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*, vol. 21, Springer-Verlag, Berlin, 1990. MR 1045822 (91i:14034)
5. Irene I. Bouw, *The p -rank of ramified covers of curves*, *Compositio Math.* **126** (2001), no. 3, 295–322. MR 1834740
6. Ching-Li Chai, *Newton polygons as lattice points*, *Amer. J. Math.* **122** (2000), no. 5, 967–990. MR 1781927
7. A. J. de Jong and F. Oort, *Purity of the stratification by Newton polygons*, *J. Amer. Math. Soc.* **13** (2000), no. 1, 209–241. MR 1703336
8. P. Deligne and G. D. Mostow, *Monodromy of hypergeometric functions and nonlattice integral monodromy*, *Inst. Hautes Études Sci. Publ. Math.* (1986), no. 63, 5–89. MR 849651
9. E. Eischen and E. Mantovan, *p -adic families of automorphic forms in the μ -ordinary setting*, preprint, available on arXiv:1710.01864.
10. Torsten Ekedahl, *Boundary behaviour of Hurwitz schemes*, *The moduli space of curves (Texel Island, 1994)*, *Progr. Math.*, vol. 129, Birkhäuser Boston, Boston, MA, 1995, pp. 173–198. MR MR1363057 (96m:14030)
11. Carel Faber and Gerard van der Geer, *Complete subvarieties of moduli spaces and the Prym map*, *J. Reine Angew. Math.* **573** (2004), 117–137. MR 2084584
12. William Fulton, *Hurwitz schemes and irreducibility of moduli of algebraic curves*, *Ann. of Math. (2)* **90** (1969), 542–575. MR 0260752
13. Darren Glass and Rachel Pries, *Hyperelliptic curves with prescribed p -torsion*, *Manuscripta Math.* **117** (2005), no. 3, 299–317. MR 2154252
14. Paul Hamacher, *The almost product structure of Newton strata in the deformation space of a Barsotti-Tate group with crystalline Tate tensors*, *Math. Z.* **287** (2017), no. 3-4, 1255–1277. MR 3719535
15. F. Knudsen, *The projectivity of the moduli space of stable curves. II. The stacks $M_{g,n}$* , *Math. Scand.* **52** (1983), no. 2, 161–199. MR MR702953 (85d:14038a)
16. Robert E. Kottwitz, *Isocrystals with additional structure*, *Compositio Math.* **56** (1985), no. 2, 201–220. MR 809866
17. ———, *Points on some Shimura varieties over finite fields*, *J. Amer. Math. Soc.* **5** (1992), no. 2, 373–444. MR 1124982
18. ———, *Isocrystals with additional structure. II*, *Compositio Math.* **109** (1997), no. 3, 255–339. MR 1485921
19. Kai-Wen Lan, *Arithmetic compactifications of PEL-type Shimura varieties*, *London Mathematical Society Monographs Series*, vol. 36, Princeton University Press, Princeton, NJ, 2013. MR 3186092
20. Kai-Wen Lan and Benoît Stroh, *Compactifications of subschemes of integral models of Shimura varieties*, preprint, available at <http://www-users.math.umn.edu/~kwlan/articles/cpt-sub.pdf>.
21. Wanlin Li, Elena Mantovan, Rachel Pries, and Yunqing Tang, *Newton polygons arising for special families of cyclic covers of the projective line*, to appear in *Research in Number Theory*, available on arXiv:1805.06914.
22. ———, *Newton polygons of cyclic covers of the projective line branched at three points*, to appear in *Research Directions in Number Theory*, available on arXiv:1805.04598.
23. Elena Mantovan, *On the cohomology of certain PEL-type Shimura varieties*, *Duke Math. J.* **129** (2005), no. 3, 573–610. MR 2169874
24. Ben Moonen, *Serre-Tate theory for moduli spaces of PEL type*, *Ann. Sci. École Norm. Sup. (4)* **37** (2004), no. 2, 223–269. MR 2061781
25. ———, *Special subvarieties arising from families of cyclic covers of the projective line*, *Doc. Math.* **15** (2010), 793–819. MR 2735989
26. Frans Oort, *Newton polygon strata in the moduli space of abelian varieties*, *Moduli of abelian varieties (Texel Island, 1999)*, *Progr. Math.*, vol. 195, Birkhäuser, Basel, 2001, pp. 417–440. MR 1827028

27. ———, *Abelian varieties isogenous to a Jacobian in problems from the Workshop on Automorphisms of Curves*, Rend. Sem. Mat. Univ. Padova **113** (2005), 129–177. MR 2168985 (2006d:14027)
28. Rachel Pries, *Current results on Newton polygons of curves*, to appear as Chapter 6, Questions in Arithmetic Algebraic Geometry, Advanced Lectures in Mathematics Book Series.
29. Eva Viehmann, *Newton strata in the loop group of a reductive group*, Amer. J. Math. **135** (2013), no. 2, 499–518. MR 3038719
30. Eva Viehmann and Torsten Wedhorn, *Ekedahl-Oort and Newton strata for Shimura varieties of PEL type*, Math. Ann. **356** (2013), no. 4, 1493–1550. MR 3072810
31. Angelo Vistoli, *Intersection theory on algebraic stacks and on their moduli spaces*, Invent. Math. **97** (1989), no. 3, 613–670. MR MR1005008 (90k:14004)
32. Torsten Wedhorn, *Ordinariness in good reductions of Shimura varieties of PEL-type*, Ann. Sci. École Norm. Sup. (4) **32** (1999), no. 5, 575–618. MR 1710754
33. Stefan Wewers, *Construction of Hurwitz spaces*, Dissertation, 1998.
34. Liang Xiao and Xinwen Zhu, *Cycles on shimura varieties via geometric satake*, preprint, available on arXiv:1707.05700.
35. Noriko Yui, *On the Jacobian variety of the Fermat curve*, J. Algebra **65** (1980), no. 1, 1–35. MR 578793

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WI 53706, USA
E-mail address: wanlin@math.wisc.edu

DEPARTMENT OF MATHEMATICS, CALIFORNIA INSTITUTE OF TECHNOLOGY, PASADENA, CA 91125, USA
E-mail address: mantovan@caltech.edu

DEPARTMENT OF MATHEMATICS, COLORADO STATE UNIVERSITY, FORT COLLINS, CO 80523, USA
E-mail address: pries@math.colostate.edu

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NJ 08540, USA
E-mail address: yunqing@math.princeton.edu